

## TEEKAY EDUCATIONAL CONSULT

Impacting lives. Email: teekayeduconsults@gmail.com Tel: 08146074387,07087006807

## **1001 PRACTICE CBT QUESTIONS ON CHM 102.**

REDEFINING EDUCATIONAL EQUITY ....

### ADDRESS FROM THE DIRECTORS



This material is dedicated to almighty God, to the entire tutors, students and management of **TEEKAY EDUCATIONAL CONSULT**. We can't but appreciate the relentless effort of **TEEKAYs** towards the production of this material which we believe will assist all first-year students offering organic chemistry in various tertiary institutions most especially student of **FEDERAL UNIVERSITY OYE EKITI**.

We hereby implore and advice all students offering organic chemistry **(CHM 102)** to make judicious use of this material as its going to prepare students ahead their examinations and will definitely bring out the best in them.

# **1001 Likely CBT Questions** in CHM 102

1. Which of the following is not a characteristic of organic compounds?

(a) They usually have low melting points.

*(b)* They usually are only slightly soluble or insoluble in water.

(c) If water soluble\* they seldom conduct an electric current.

(d)Bonds which bind the atoms together are nearly always ionic. Answer. (d

an

2. The element least likely to be found in organic compound is

(a) oxygen

*{b)* sulfur

(c) nitrogen

(d) silicon

Answer. (d)

3- One of the major sources of organic compounds is

(a) Natural gas

(b) Fermentation

- (c) Sea water
- (d) Atmosphere

Answer. A

4.Catenation is a property of the carbon atom which describes its ability to
(a) bond with other carbon atoms
(b) form double and triple bonds
(c) exist in plant and animal form
(d) form bonds in its ground state
Answer. (a)
5.stable organic compounds, carbon will always form
(a) 2 bonds

#### Answer.

(b)

6- Carbon-carbon double bonds consist of:

(a) *one o* bond, one pi bond

- (b) Two sigma bond, one pi bond
- (c) One sigma, two pi bonds
- (d) Two sigma, two pi bon ANSWER (a)

#### Answer. (a)

7. Acetylene has a total of:[a) one *a* bond, two pi bonds

- *(b) two a* bond, four π bonds
- (c) Three o bonds, two п bond
- (d) one a bond, four pi bonds

#### Answer (c)

8. In propene there are

- (s) Eight o bonds and one n bond
- (b) seven o bonds and two n bonds
- (c) six o bonds and three pi bonds (d) ALL

#### Answer. (a)

- 9. In propyne there are
- (a) six o bonds and two n bonds
- [b] seven *a* bond and one n bond
- (c) six n bonds and one n bond
- (d) eight bonds

#### Answer. (a)

- 10. 1-Buten-3-yne has
- (a) six o and four n bonds
- (b) seven  $o \mbox{ and } three \mbox{ } n \mbox{ bonds}$

(c) eight o and two n bonds(d) ALL

#### TEEKAY EDU CONSULT

11. Compound in which carbons use only spa hybrid orbitals for bond formation is
(a) CH3CH2CH3
(b) CH3C-CH
(c) CH3CH=CH 2
(d) CH2=CH—CH=CH2
Answer. (a)
12. Compound in which carbon uses sp3 hybrid orbitals for bond formation is

(a) H - C - H

(b) N C- NHS

(c) H−C − 0H(c) (CH3)3COH

ANSWER (d)



VOTE **ADEDIGBA ISRAEL TAIWO** as THE NEXT FACULTY OF SCIENCE **PRESIDENT**. He is one of our directors at TEEKAY EDUCATIONAL CONSULT

Teekay Page 3	Educational Consult
Answer. (d)	(c) 1.54 A
13. When the carbon atom is sp2 hybridized in a	(d) 1.68 A.
compound, it is bonded to	Answer. {b]
(a) 2 other atoms	20. The carbon-carbon bond lengths in rank of
(b) 4 other atoms	increasing bond length is:
(c) 3 other atoms	(a) triple, double, single
(D)5 other atoms	(b) single, double, triple
Answer. (c)	(c) single, triple, double
14 Compound in which carbons use only sn2	(d) triple, single, double
hybrid orbitals for hand formation is	Answer (a)
	21 Which of the following is the correct order of
$\sim$	bond longths?
	boliu lenguis: (a) $C = C = C = C < CYC$
	$ (a) \cup - \cup \ll \cup = \cup < \cup \land \cup = \cup =$
	(b)
(a)	$(c) \ t - t \gg t - t > t = t$
	[a] $L-L < L-L > L=L$
$\langle \cdot \rangle$	Answer. (a)
	22. Which of the following hydrocarbons has the
	shortest C-C bond length?
<i>(b)</i>	(a) CH2=CH2
(c) CH2=CH-CH=CH2	<i>(b)</i> CH3CH3
(d) CH3CH=C=CH2	(c) HC=CH
Answer. (c)	
15 Which of the following is a planar molecule?	
(a) Formaldehyde	
(h) Acetone	
(c) Formic acid	
(d) Agotic poid	(u) Answor (c)
	Allsweit. (C)
Answer. (a)	23. The carbon-carbon bond length is maximum
16. What is bond angle between the hybrid	In
orbitals in methane?	(a) CH2=CH2
(a) 180°	(b) CH3CH3
<i>(b)</i> 120°	(c) HC-CH
(c)109.5°	
(c) 115.S°	
Answer. (c)	
17. The H—C—C bond angle in ethane is	
(a) 60º	
(a) 109.5°	(d)
(c) 120º	Answer. (b)
(d)118°28s	24. What is the hybridization of the carbon
Answer. (b)	atoms numbered 1 and 2 respectively in the
18 The H-C-H bond angle in ethylene is	following structure?
(a) $60^{\circ}$	
<i>[h</i> ] 120 <sup></sup>	
(a) 00"	
(1) 30	
	$\sim$
Allswer. (D)	$\mathbf{\tau}$
19. What is the bond length of a carbon-carbon	(a) sp3, sp2
double bond?	[b1 sp2, sp2
(a) 1.20 A	[c] sp, sp
(b) 1.34 A	<i>{d} sp2, Answer</i> . (d)
Impacting Lives	

 $\boldsymbol{\wedge}$ 

Teekay Page 4	Educational Consult
25. How many atoms are attached to an atom	32. Which <i>molecule</i> has the greatest dipole
having a <i>sp</i> hybridization?	moment
(a] 0	(a) CH3CI
(b) 1	(b) CH3Br
(c) 2	(c) CH3F
(d) 3	(d) CH3l
Answer. [c)	Answer. (c)
26. Which statement is true?	33. Which of the following compounds have
(a) Resonance hybrids <i>are</i> inherently unstable.	highest dipole moment?
<i>(b)</i> Resonance hybrids are more stable than any	(a) dichloromethane
individual resonance form.	[b] chloroform
(c) Resonance hybrids are averages of all	(c] chloromethane
resonance forms resembling the less	[d] carbon tetrachloride
stable forms.	Answer. (a)
(d) Resonance hybrids are <i>averages</i> of all	34. Homolytic fission of C—C bond leads to the
resonance forms resembling the more	formation of:
stable forms.	(a) Free radicals
Answer. (d)	(b) Carbonium ions
27. Resonance structures of a molecule have	(c) Carbanions
(a) same arrangement of atoms	(d) None of these
<i>(b)</i> different arrangement of atoms	Answer. (a)
(c) same number of paired electrons	35. Homolytic fission of covalent bond between
(d) different number of paired electrons	carbon atoms will produce:
Answer. (a) and (c)	(a) Two carbonium ions
28. Which of the following compounds have	[b] Two molecules
planar molecules?	(c) Free radicals
(a) ethyl alcohol	(d) Carbonium ton and carbanion
(b) formaldehyde	Answer. (c)
(c) diethyl ether	36. The order of stability of carbonium ions is
(d) 1,3-butadiene	(a) tertiary > secondary > primary
Answer. [b) and (d)	[b1 secondary > tertiary > primary
29. Which of the following compounds will show	(c) primary > secondary > tertiary
dipole moment?	(d) primary > tertiary > secondary
(a) <i>cis</i> -1,2-dichloroethylene	Answer(a)
<i>{b</i> ) o-dichlorobenzene	37. The order of stability of carbanions is
(c) <i>trans</i> - 1,2-dichloroethylene	(a) primary > secondary > tertiary
(d) -dichlorobenzene	(b) secondary> tertiary > primary
Answer. (a) and {b)	(c) tertiary > secondary > primary
30. Which molecule has a nonzero dipole	(d) tertiary > primary > secondary
moment?	Answer(a)
(a) C12	38. Which of the following carbonium ions willbe
<i>{b)</i> C02	most stable?
(c) CCI4	(a)+^CH3
(d) CHCI3	(b) CH3'CH2
Answer. (d)	(c) (LH3)2'LH2
31. Which of the following compounds have	(d) LH2 = LHLH2
lowest dipole moment?	Answer (d)
(a) carbon tetrachloride	39. The least stable carbanion is :
(b) chloromethane	( <i>a</i> )( <i>b</i> ))20
(c) aichioromethane	(D) (UH3)3U
(a) chlorotorm	
Answer. (a)	(ajlH3

Teekay Page 5	Educational Consult
Answer. (b)	(a) 6-10 kcals
40. Which alkyl free radical is the most stable?	(b) 60-80 kcals
(a] methyl	(c] 80-100 kcals
(b) primary	(d) 100-120 kcals
(c) secondary	Answer. (a)
(d) tertiary	49. Which describes the bond strength or bond
Answer. (d)	dissociation energy?
41. Which of the following is an electrophile?	(a) energy required to break a bond
(a)CH3 O	(b) energy released when a bond break
сустать справления справления справления справления справления справления справления справления справления спра	(c) energy released when a bond is formed
(c)NH3	[d] [a] and [c]
(d) CH3CH2	Answer. (d)
Answer. (b)	50 Which statement is correct?
42. Which of the following is not an electrophile?	(a) Energy is released when a hond break
(a) NH3	(h) A sigma hand results from attraction of
(h) Br+	nrotons and electrons
(c) AIC13	(c) Energy is released when a bond form
(d) N02+	(d) A carbanion is positively charged
Answer (a)	Answer (c)
43 Which of the following statements is correct	51 Which of the following is incorrect?
regarding nucleonhiles?	(a) Resonance stabilization is the extra stability
(a) They have an overall positive charge	compound gains from having delocalized
(a) They have all overall positive charge	compound gams nom naving delocalized
(c) They have an unpaired electron	chectrons.
(d) They have an unpaired electron	(D) Delocalized electrons destabilize a
(u) They have empty of bitals	(a) The greater the number of relatively stable
Allswei. [0]	(c) The greater the number of relatively stable
(a) AICI2	resonance contributors, the greater is the
	resonance stabilization.
(a) PE2	(a) $(a)$ and $[c]$
	Answer. [D]
$(u) \cup (d)$	52. Which of the following is correct?
Allswei. (u)	(a) curved arrows are always drawn from an
45. Which of the following is not a nucleophne?	electron poor center to an electron rich center.
	[b] A doubled headed arrow means one electron
(0) HSU3—	has been moved.
(C) AICI 3	(c) Curved arrows are always drawn from an
(d) H0—	electron rich center to an electron poor center.
Answer. (C)	(d) A single headed arrow means two electrons
46. Which of the following is not a nucleophile?	have been moved.
	Answer. (c)
(b) BFW	53. What does a "curved" arrow represent?
(c) CH30	[a] that two structures are resonance structures
(d) NH3	(b) the movement of two electrons
Answer. (b)	(c) a link between reactants and products
47. Which of the following is a Lewis acid?	(d) that two structures are equivalent
(a)AlCl3	Answer. (b)
<i>(b)</i> CH30H	54. What is the predicted shape, bond angle, an
(c) NH3	hybridization for CH3•?
(d) CH3OCH3	(a) trigonal planar, 120°, sp2
	(h) trigonal planar, 120°, sp3
Answer. {a)	
48. Which of the following ranges best	(c) trigonal planar, 109.5°, sp2

Teekay	Page 6	Educational Consult
Answer. (a)		62. Which of the following is not normally
55. What is the name given for a specie <i>es</i>		considered to be a nucleophile?
that contains a positively charged carbon atom	?	(a) NH3
(a) carbanion		<i>(b)</i> CH3NH2
<i>(b)</i> carbocation		(c) HC-C: —
(c) methyl radical		(d) CH3CtI2+
(d) free radical		Answer. (d)
Answer. (c)		63. Which of the following is nor a nucleophile?
56. What orbitals overlap to create the II—C	bond	(a) FeBr3
in CH3+?		(b) Br—
(a] <i>sp3-sp3</i>		[c] NH3
<i>{b)</i> sp2—sp3		[d] CH30CH3
(c) <i>s-p</i>		Answer (a)
(d) <i>s-sp2</i>		64. Which of the following is the strongest
Answer. (d)		interaction?
57. The lone-pair electrons of the methyl and	nion	(a) a covalent bond
occupy anorbital.		(b) dipole-dipole interactions
(a) <i>s</i>		(c) hydrogen bonding
(b) <i>p</i>		(d) van der Waals
(c) sp2		Answer. (a)
(d) sp3		65. Which of the following statements is
Answer. (d)		incorrect?
58. An increase in which of the following re-	sults	(a) Electrons move toward positively charged
in a decrease in the rate of the chemical		locations.
reaction?		[b] An electron-rich atom is called an
(a] temperature		electrophile.
(b) concentration		(c) An electrophile is electron loving
(c) collision frequency		(d) A nucleophilic has a pair of electrons it can
(d) energy of activation		share.
Answer. (d)		Answer. [b]
59. The reaction step that has its transition	state	66. Which of the following is a nonpolar
at the highest point on the reaction coordin	ate is	molecule?
the. Called the		(a) HC1
(a) rate-determining step		[b] CH3CI
{ <i>b1</i> activation energy		(c) H2
(c] transition step		(d) NH3
(d) product favored step		Answer c)
Answer. (a)		67. Which is defined as a species that accepts a
60. An electrophile act as a when	it	proton?
reacts with a nucleophilic.		(a) Lewis acid
(a) Bronsted- Lowry base		(b) Lewis base
<i>{b}</i> Arrhenius base		(c) Bronsted-Lowry acid
{c) Lewis acid		(d) Bronsted-Lowry base
(d) Lewis base		Answer. (d)
Allswer. (c)		68. Which statement is correct about acid-base
o1. A nucleophile acts as a When 1	ι	cnemistry?
reacts with an electrophile.		(a) the Larger the <i>Ka</i> , the stronger the acid
(a) DI UIISIEU-LOWFY ACIU (b) Arrhonius hago		(b) a small pka correspond to a large Km
(c) Lowis acid		(d) all of the above
(d) Lowis base		Answer (d)
Answer (d)		Mich of the following statements is correct?
		which of the following statements is correct?
Impacting Lives		

Teekay	Page 7	Educational Consult
(a) The stronger the acid, the weaker its		(d) same functional groups
conjugate base.		Answer. {b}
<i>{h</i> ] The stronger the acid, the stronger its		77. Compounds with the same molecular
conjugate hase		formula but different structural formulas are
(c) The stronger the base the stronger its		called
conjugate hase		(a) Alkovides
(d) The stronger the acid the weaker its		(b) Iso compounds
(u) The subliger the actu, the weaker its		(b) iso compounds
conjugate aciu.		(c) isoliters
Answer. (a)		(a) Ortho compounds
/V. What is the conjugate actu of CH3NH2?		Answer. (c)
(a) CH3NH3+		78. Ethanol [CH3CH20H] and dimethyl ether
{D} CH3NH—		(CH30CH3) are best considered:
(C) NH4+		(a) structural isomers
(d) NH2—		(b) stereoisomer
Answer. (a)		(c) enantiomers
71. What is the conjugate base of CH3NH2?		[d] diastereomers
(a) CH3NH3+		Answer. (a)
<i>{b)</i> CH3NH—		79. The compounds CH3CH20CH 2CH3 and
(c) NH4+		CH3OCi-I2CH2CH3 are
(d) NH2-		(a) Enantiomers
Answer. (b)		(b) Conformational isomers
72. The stronger the acid		(c) Metamers
(a) the less stable its conjugate base		(d) Optical isomers
{bj the larger the pka		Answer. (c)
(c) the weaker its conjugate base		80. Which of the following statements is fa1se
<i>{d)</i> the larger the pH		about tautomer's?
Answer. (c)		(a) Tautomer's are structural isomers
73. Which of the following 1s the strongest a	acid?	(b) Tautomer's arc structural isomers which
(a) HI		exist in dynamic equilibrium
<i>{b)</i> H20		(c) Tautomerism involves movement of atoms
(c) CH4		(d) Tautomer's have independent existence
(d) CH30H		Answer. (d)
Answer. {a)		81. How many isomers are possible for the
74. Which of the following are Lewis bases?	?	compound with molecular formula
(a) NF3		C4H8?
(b) BF3		(a) 2
(c) CH3OCH3		(b) 4
(d a and c		[c] 6
Answer. (d)		(d) 8
75. Two compounds have the same composit	tion	Answer. (c)
and also have the same atoms attached to t	he	82. How many str natural isomers are possible
same atoms, although with different orientat	tions	for C4H9Br?
in space. These compounds are		(a) 2
(a) Identical		[b] 3
(b) Position isomers		(c) 4
(c) Structural isomers		(d)5
(d)Stereoisomers		Answer (c)
Answer. (d)		83 How many isomeric aromatic hydrocarbons
76. The isomers of a substance must have		arc possible for C8H10?
(a) same chemical properties		(a) 3
<i>(b)</i> same molecular weight		(b) 4
(c) same structural formula		(c) 5
	I	
Impacting Lives		

Teekay	Page 8	Educational Consults
( <i>d</i> ) 6		(a) BrCH=CHBr
Answer. {b)		(b) BrCH=CHCl
84. Which compound is not an isomer of	of the	(c)
other three?		[d]
(a) n-Pentane		Answer. (d)
(b) 2.2-Dimethy1propane		92 A molecule is said to be chiral
$\{c\}$ 2 - Methyl butane		(a) if it contains plane of symmetry
(d) 2.3-Dimethylbutane		(b) if it contains Centre of symmetry
Answer. (d)		(c) if it cannot be superimposed on its mirror
85. Alkenes show geometrical isomeris	m due to	image
		(d) if it can be superimposed on its mirror image
(a) Asymmetry		Answer. (c)
(b) Rotation around a single bond		93. Which of the statements is false regarding
(c) Resonance		chiral compounds?
(d) Restricted rotation around a double	bond	(a) rotate the plane of polarized light
Answer. (d)	Donia	(b) have cis and trans isomers
86. Which of the following compounds ex	hibit	(c) exist as enantiomers
geometrical isomerism?		(d) can be detected with a polarimeter
(a) 1-Pentcne		Answer. (b)
(h) 2-Methyl-2-pentene		94 An ontimally active compound
(c) 2-Pentene		(a) must contain at least four carbons
(d) 2-Methyl-2-hutene		(b) when in solution rotate the plane of polarized
Answer (c)		light
87 Which of the following compounds	may exist	(c) must always contain an asymmetric carbon
as cis-trans isomers?	may exist	atom
(a) l-Rutene		(d) in solution always give a negative reading in
(h) 2-Botene		nolarimeter
(c) Cyclopropape		Answer (b)
(d) Acetone		95 Plane-polarized light is affected by
Answer [h]		(a) Identical molecules
88 Geometrical isomerism is shown by	7	(b) All polymors
(a) Lactic acid	V	(b) All polymers
(b) Maloic acid		(d) All hismologulos
(v) Maleic actu		(u) All biolitolecules
(d) 1 1 Dichloroothylono		Allswel. (C)
(u) 1,1-Dicition betriytene		96. It is possible to distinguish between optical
90 Which of the following compounds	chow	(a) by using chemical tests
sometrical icomoriam?	SHOW	(a) by using chemical tests
		(b) by mass spectrometry
(b) CH2CH-CH2		(c) by IK spectroscopy (d) by Delector try
(D) CHOCH=CH Z		(u) by Polarimeu y
(C) (CH3)2NH		07 Ontical icomore that are mirror images are
		solled
Answer. (a)	.11 .1.	(a) Tautamara
90. Which of the following compounds	will show	(d) Idulomens
geometrical isomerism?		(b) Diastereonners
$(a) \cup H2 = \cup H \cup I2$		(c) Ellaliuolliers [d] Motomore
		Luj Metamers
		Allswer. (C)
$\{a\} \cup Z \cup U = \cup B \cup Z$		98. Optical isomers that are not mirror images
Answer. {b)		are called
91. Which of the following compounds w	ill not	(a)Diastereomers
show geometrical isomerism?		(b) Enantiomers
Impacting Lives		

Teekay	Page 9	Educational Consult
©Metamers		(c) 2-Ch1oropropanoic acid
(d)Meso compounds		(d) 3-Chloropropene
Answer. (a)		Answer. (c)
99. Enantiomers have which of the followi	ng	<b>106.</b> Which of the following compounds will be
characteristics?	U	optically active?
(a) rotate ordinary light		(a) Succinic acid
(b) have the same melting point		(b) meso-Tartaric acid
(c) are superimposable mirror images		(c) Lactic acid
(d) react with optically active molecules	at the	(d) Chloroacetic acid
same rate		Answer. (c)
Answer. (b)		<b>107.</b> Which of the following isomeric
100. Which of the following statements is	s false	compounds show optical isomerism?
about enantiomers?		(a) Î-Amino pentane
(a) rotate plane-polarized light		(b) 2-Aminopentane
(b) are superimposable mirror images		(c) 3-Aminopentane
(c) are nonsuperimposable mirror image	es	(d) 2,2-Dimethylpropylamine
(d) have the same melting point		Answer. (b)
Answer. [b]		108 2-Butanol is optimally active because it
101. A meso compound:		contains:
(a) is an achiral molecule which contains cl	hiral	(a) an asymmetric carbon
carbons		(b)j a plane of symmetry
(b) contains a plane of symmetry or a Cen	tre of	(c) a hydroxyl groups
symmetry		(d) a Centre of symmetry
(c) is optically inactive		Answer. (a)
(d) is characterized by all of the above		109. Optical isomerism is shown by
Answer. (d)		(a) n-Butyl chloride
102. What is the possible number of opti	ical	(b) sec-Buty1 chloride
Isomers for a compound containing n dissi	milar	(c) <i>tert-But</i> yl chloride
asymmetric carbon atoms?		(d) Isobutyl chloride
(a) n2		Answer. [b]
<i>(b) 2n</i>		110. Which of the following compounds is an
(c) n + 1		optically active compound?
(d n + 2		СН
Answer. (b)		
103. What is the possible number of opt	ical	$(a)$ $CH_3$ $CH_2$ $COH_3$
isomers for a compound containing 2 dissi	milar	<i>(b)</i> CHCI3
asymmetric carbon atoms?		(c) CH3CH2COOH
(a) 2		(d) <b>CH3CH</b> 2OH
<i>b</i> ) 4		Answer. (a)
(c) 6		
( <i>d</i> ) 8		
Answer. (b)		111. Lactic acid Cr 1 molecule
<b>104.</b> meso-Tartaric acid is		which shows
(a) sometimes optically active		(a) Geometrical isomerism
<i>(b)</i> always optically active		(b) Tautomerism
(c) sometimes optically inactive		(c) Optical isomerism
(d) always optically inactive		(d) metamerism
Answer. (d)		Answer. (c)
<b>105.</b> Which of the following compounds w	vill be	112. How many optical isomers <i>are</i> possible for
optically active?		
(a) Propanoic acid		
<i>{b]</i> 3-Chloropropanoic acid		(0) 4
Impacting Lives		

Teekay	Page 10	Educational Consult
(c) 6		119. Which of the following compounds is never?
(d) 8		chiral?
Answer. (a)		(a) 2,3-dibroinobutane
113. Which of the following represents a		(b) 1,3-dibroinobutane
racemic mixture?		(c) 1,2-dichlorobutane
(a) 75% (R)-2-butanol, 25% (S)-2-butanol		(d) 1,4-dibromobuta se
(b) 25% (R)-2-butano1, 75% (S)-2-biitanol		Answer. (d)
(c) 50% (R)-2-butano1, 50% (S)-2-butanol		<b>120.</b> Which of the following statements is
(d) none of the above		correct concerning a pair of
Answer. (c)		enantiomers?
114. Consider [R]- and (S)-2-butanol. Whi	ich	[a) They rotate the plane of polarized light by
physical property distinguishes the two		exactly the same amount and in opposite
compounds?		directions.
(a) melting point		[b] They rotate the plane of polarized light by
(b) solubility in common solvents		differing amounts and
(c) Rotation of plane-polarized light		in opposite directions.
(d) infrared spectrum		(c) They rotate the plane of polarized light by
Answer. (c)		differing amounts and in the same direction.
155. Which of the following is capable of		(d) They have different melting points.
exhibiting cis-trans isomerism?		Answer. (a)
(a) l-butene		121. Which of the following is <i>not</i> true of
<i>{b)</i> I-pentene		enantiomers?
(c) ethene		(a) They have the same melting point.
(d) 2-butene		(b) They have the same boiling point.
Answer. (d)		(c) They have the same density.
<b>116.</b> Which of the following is a true state	ment?	(d) They have the same specific rotation.
(a) All chiral molecules possess a plane of		Answer. (d) 122 A and P are storeoisomore. They are
(b) All achiral moloculos aromoso		<b>122.</b> A difference of a consumption of the second
(c) Ail molecules which possess a single		one another. Which of the following best
asymmetric center of the S configuration a	are	describes the relationshin?
levorotatory.		between A and B?
(d) A mixture of achiral compounds will b	e	(a) structural isomers
optically inactive.		[b] enantiomers
Answer. (d)		(c) cis-trans isomers
117. Which of the statements below corre	ctly	(d) diastereomers
describes an achiral molecule?		Answer. (b)
(a)The molecule has a nonsuperimposable		<b>123.</b> Which of the statements is correct about
mirror image?		diastereomers?
(b) The molecule exhibits optical activity w	vhen it	(o) They are stereoisomers that arc not
interacts with plane-polarized light.		enantiomers.
(c) The molecule has an enantiomer.		(b) They are a pair of identical isomers.
(d) The molecule might be a meso form.		(c) They are a pair of isomers that are mirror
Answer. (a)		Images.
in a malagula of 2.4.6	resent	(d) An their asymmetric centers are the same.
III a III01ecule 01 2,4,0- trimothyl hontono?		<b>124</b> Which of the following is /are ontically
(a) ()		inactive?
(b) 1		(a) <b>a 50-50 mixture</b> of R and S enantiomers
(c) 2		(b) a meso compound
(d) 3		(c) a racemic mixture
Answer. (a)		(d) all the above
Impacting Lives		

Teekay	Page 11	Educational Consult
Answer (d)		(c) C8H18
<b>125.</b> Hydrocarbons are		(d) C8 H16
(a) Composed of carbon and hydrogen		Answer. (a)
(b) Composed of carbon hydrogen and ovi	ann	<b>133</b> How many isomers are possible for butane?
(c) Composed of carbon and ovugon	ygen	(a) 2
(d) Composed of carbon and nitrogon		(a) 2
(u) composed of carbon and incrogen		
<b>126</b> Hydrocarbons are		(c) 4 $(d)$ 5
(a) in calluble in water		(u) J
(a) insoluble in water		Allswei. (D)
(b) composed of carbon and hydrogen		<b>134.</b> How many isomers are possible for 2-
		methylpropane?
(d) None of these		
Answer. (c)		(b) 3
127. Which of following statements is false	2	(c) 4
about propane?		( <i>d</i> ) 5
(a) all bond angles are 109.5°		Answer (a)
<i>(b)</i> each carbon is sp3 hybridized		<b>135.</b> How many isomers are possible for
(c) the compound is combustible		pentane?
(d) the compound undergoes polymerization	on to	(a) 2
give polypropylene.		(b) 3
Answer. (d)		(c) 4
<b>128.</b> Which of the following is a correct nam	e	(d) 5
according to the IUPAC rules		Answer. {b)
(a] 2-Methylcyclohexane		<b>136.</b> How many isomers are possible for
(b) 2-Ethyl-2-inethy1pentane		hexane?
(c) 3,4-Dimethylpentane		(a) 4
(d) 3-Ethyl-2-merhylpentane		(b) 5
Answer. (d)		(c) 6
129. A tertiary carbon is bonded directly to	):	(d) 7
(a) 2 hydrogens		Answer. [b]
<i>(b) 3</i> carbons		137. How many <i>isomers</i> are possible for
(c) 2 carbons		heptane?
(d)4 carbons		(a} 9
Answer. (b)		<i>[b]</i> 10
<b>130.</b> What <i>type</i> of an alkyl group is an isob	utvl	(c) 11
group?	5	(d)12
(a) primary		Answer (a)
(b) secondary		138 Marsh gas mainly contains
(c) tertiary		(a) CH2=CH2
(d) none of the se		(h) CH4
Answer. [a]		(c) H2S
<b>131</b> . Which molecular formula indicates 2-	-	[d] CHC13
methyl nentane?		Answer {h}
(a) C5H12		139 Which statement is false?
<i>{h i</i> C6H14		(a) Many alkanes are soluble in water
(c) C5H10		(h) All alkanes have a lower density than water
(d C6H12		(c) At room temperature some alkanes are
Answer (h)		liquide some solide some gases
<b>132</b> Which molecular formula indicates 2	2 4-	Iquius, soine soinus, soine gases. (d) All alkanas hurn
trimethyl hevene?	т	
(a) COH20		Answer (a)
	ļ	
Impacting Lives		

~



Please TEEKAY need your Support and VOTE in this election . kindly VOTE when its time to do so. Thanks

Teekay	Page 12	Educational Consult
140. As the molecular weight of alkanes		(c) n-Hexane
increases, how do the boiling point and me	elting	(d) Neopentane
point change?		Answer. (d)
(a) boiling point increases; melting point		<b>146.</b> Methane can be prepared by:
increases.		(a) the reaction of iodomethane with sodium in
(b) boiling point increases; melting point		dry ether
increases.		(b) the reaction of methanol with concentrated
(c) boiling point increases; melting point		H2504
decreases		(c) the reaction of sodium methanoate with
(d) boiling point increases; melting point		soda-lime
increases sequentially for alkalles over for	11	[a] the reaction of sodium ethanoate with soda-
Answer (d)		Inne Answor (d)
<b>141</b> The branching of alkanes that produc	<b>' C</b>	<b>147</b> Which of the following reactions can be
symmetrical structures.	.03	used to prepare alkanes?
(a) raises the boiling point : raises the melt	ing	(a) Corev-House synthesis
point		(b) Williamson synthesis
<i>{b</i> } raises the boiling point : lowers the melti	ing	(c) Friedel-Crafts reactions
point	0	(d) None of these
(c) lowers the boiling point; lowers the me	lting	Answer. (a)
point		148 Which of the following reactions can beused
(d) lowers the boiling point; raises the mel	ting	to prepare alkanes?
point		(a) Wurtz reaction
Answer. (a)		(b) Wolf-kishner reduction
<b>142.</b> How does the melting point of propagation	ne and	(c) Kolbe's electrolysis
ethane compare?		(d) All of these
(p) the melting point of propane is greater.		Answer. (d)
(b) the melting point of ethane is greater.		<b>149.</b> Kolbe's electrolysis of sodium butyrate
(r) the melting points are within 5°C of one		gives
(d) no relationship ovists among those allo	noc	(a) C6H10
and their melting points	uies	(c) C8H18
Answer. [b]		(d) C6H12
<b>143.</b> How do the boiling points of butane.		Answer. (h)
propane, and ethane compare:'		<b>150.</b> Which of the following classes of
(a) the boiling point of ethane is greatest.		compounds is unreactive toward sulfuric
(b) the boiling point of butane is greatest.		acid?
(c) the boiling point of propane is greatest.		(a) Alkanes
(d) there is no relationship among these th	iree	[b ) Alcohols
alkanes and their boiling points.		(c)Alkenes
Answer. (b)		(d) Alkynes
<b>144.</b> Which of the following alkanes will h	ave	Answer. (a)
the highest boiling point?		<b>151.</b> Which of the following compounds does
(a) n-Octane		not dissolve in concentrated H2SU4?
$\{b\}$ isopenialle		$\begin{array}{c} \text{OII walling:} \\ \text{(a) n Hoyano} \end{array}$
(d) Neopentane		(a) n-nexane (b) Diethyl ether
Answer. (a)		(c) 1 -Butene
<b>145.</b> Which of the following alkanes will have	ave	(d) Aniline
the <i>lowest</i> boiling point?		Answer (a)
(a) n-Heptane		<b>152.</b> 2,3-Dimethyl-2-butene undergoes catalytic
<i>{b)</i> isopentane		hydrogenation to give
Impacting Lives		

Teekay	Page 13	Educational Consult
(a) 2,3-Dimethylbutane		158. What product is formed in the free-radical
(b) 2-MethyIpentane		bromination of methane?
(c) 2,2-DimethyIbutane		[a) bromomethane
(d) 3-Methylpentane		(b) dibromo methane
Answer. (a]		(c) tribromomethane
<b>153-</b> Why is the halogenation of alka	nes	(d) all of these
considered a chain reaction?		Answer(d)
(al it occurs quickly.		159. Chlorination of an alkane as compared to
<i>(b)</i> it occurs without the generation of	of	bromination proceeds
intermediates.		(a) at a slower rate
(c) each sten generates the reactive int	ermediate	[b] at a faster rate
that causes the next step to occur	.er meulute	(c) with equal rates
(d) the reaction allows long chains of	f	[d] with equal or different rate depending upon the
halogenated alkanes to be formed		source of alkane
Answer. (c)		Answer (h)
154 The chlorination of methane to	vive CCI4 is	160 How many monochlorinated isomers
an example of	Sive duri is	would result from the reaction of chlorine with
(a) an electrophilic addition		n-hutane in the presence of IIV light?
(h) a free-radical substitution		(a) 2
(c) a nucleonhilic addition		(a) 2
(d) an electrophilic substitution		
Answer (b)		(d) 8
155 In the chlorination of alkanes th	ng first stop	$\Delta nswer(2)$
in which chlorine free radicals	le m st step	161 How many monochlorination products are
are produced is called:		nossible in the reaction of 2.2. dimethylbutane
(a) initiation		with chloring in the process of ultraviolet light?
(b) initiation		(a) 2
(c) propagation		(6) 3
(d) deactivation		(0) $(c)$ $(c)$
Answer (a)		(d) 5
156 Chloring free radicals react with	mothano	(u) J
hv	rinethane	162 A compound of formula C3H8 doos not
(a) donating their free-radical electro	on to	react with broming in CC14 in the dark. The
mothana to form chloromothana		compound could be
<i>(h)</i> abstracting a hydrogen atom from	mothano	(a) Alkano
and producing HCI and a methyl rad	lical	(b) Cycloalkane
(c) forming a carbanion intermediate	a that	(c) Alkene
rapidly dissociates to produce	, tildt	(d) Cycloalkene
chloromethane		Answer (a)
(d) forming a carbonium ion intermed	lista that	163 The combustion of Pentane produces:
rapidly dissociates to form	liate tilat	(a) Pentene
chloromethane		(h) HC1 + H20
Answer [h]		(c) Pentyne
157 Which halogen does not apprecia	hly react	(d) $CO2 + H2O$
with methane in a free-radical substitu	ition	$\frac{1}{2} \frac{1}{2} \frac{1}$
reaction?		164. The combustion of one mole of propage
(a) chlorine		C3H8 produces how many moles of H202
(h) bromine		(a) 2
(c) iodine		(a) 2 (b) 3
(d) fluorine		
Answer (c)		[v] T (d) 5
		(u) J Answor (c)

Teekay Page 14	Educational Consults
165. The major carbon compound formed from	Answer. (c)
the incomplete combustion of a hydrocarbon in	172. Which of the following has the greatest
air is	solubility in CH3CH2CH2CH3?
(a) carbon dioxide	(a) CH30H
<i>(b)</i> carbon monoxide	(b) CH3NH2
(c) water	(c) CH30CH3
(d) alkyl chains	(d) (CH3)3CH
Answer. {b]	Answer. (d)
<i>166.</i> The thermal decomposition of alkanes in	173. Consider the three isomeric alkanes n-
the absence of air is <i>called</i> :	hexane. 2.3-dimethylbutane. and 2- methyl
(a) Combustion	pentane. Which of the following correctly lists
<i>(b)</i> Oxidation	these compounds in order of increasing boiling
(c) Cracking	point?
(d) Hydrogenation	(a) 2.3-dimethylbutane<2-methylpentane< n-
Answer. (c)	hexane
167. LPG (Household cooking gas) is mainly a	(h) 2-methy1pentane $<$ n-hexane $<$ 2.3-
mixture of	dimethylbutane
(a) Methane + Ethane	(c) 2-methylpentane < 2 3-dimethylputane < $n$ -
(b) Acetylene + $0$ ?	hexane
(c) Butane + Isobutane	(d) n-heyane $< 2$ -methy1nentane $< 23$ -
(d) Acetylene + H2	dimethylbutane
Answer (c)	Answer (a)
168 What type of honding is most important in	174 The eclinsed and staggered forms of ethane
CH3CH2CH2CH2CH2CH3?	are said to differ in
(a) ionic	(a) molecular formula
<i>(b)</i> hvdrogen	(b) configuration
(c) covalent	(c) conformation
(d) polar	(d) constitution
Answer. (c)	Answer. (c)
169. What is the name given to a compound	175. Octane number is related to
containing only carbons and	(a) Gasoline
hydrogens and having the maximum number of	(b) Kerosene oil
hydrogens in the molecule?	(c) Diesel oil
(a) alkene	(d) Lubricating oil
(h) unsaturated hydrocarbon	Answer. (a)
(c) saturated hydrocarbon	176. A knocking sound is produced in the engine
(d) alkyne	when the fuel
Answer. (c)	(a) Burns fast
170. Which of the following is the best	[b] contains lubricating oil
description of propane, CH3CH2CH3, at room	(c) Burns slow
temperature?	(d) contains water
(a) liquid, soluble in H2O	Answer. (a)
<i>(b)</i> gas, soluble in gasoline	177. The octane rating of gasoline provides
(c) liquid, soluble in gasoline	information on
(d) gas, soluble in water	(a) its antiknock properties
Answer. (b)	(b) its ignition properties
171. Choose the correct hybridization for the	(c) its percentage proportion of octane
atom indicated in the molecule below.	(d) its percentage content of isooctane
(a) sp	Answer. (a)
(b) sp2	178. In any fuel, the percentage by volume of
(c) sp3	isooctane in a mixture of isooctane and n-
(d) none of these	

Teekay Page 15	Educational Consult
heptane which will knock under same conditions	(d) Lead acetate
as the fuel being tested, is called:	Answer. {b)
(a] Cracking	185. Which of the following methods can be
(b) Iodine number	used to increase the octane rating of gasoline?
(c) Aromatization	(a) Adding branched-chain alkanes
(d) <i>n</i> umber	(b) Adding tetraethyllead
Answer. (d) = octane number	(c) Adding aromatic hydrocarbons
<b>179.</b> Which of the following compounds is	[d] All of these
assigned the Octane Number of zero:	Answer. (d)
(a) n-Octane	<b>186.</b> The carbon atoms involved in the double
<i>{b</i> ) 2,3,3-Trimethylpentane	bond of an alkene are
(c) n-Heptane	<i>{a) sp</i> hybridized
(d) 2,2,4-Trimethylpentane	(b) sp2 hybridized
Answer. (c)	(c) <i>sp3</i> hybridized
180- Which of the following compounds is	(d) None of these
assigned the Octane Number of 100:	Answer (b]
(a] <b>n-Heptane</b>	187. Which of the following compounds will
b) 2,3,3-TrimethyTpentane	show geometrical isomerism?
(c) n-Octane	(a) Propene
(d) 2,2,4-Trimethylpentane	(b) <b>2-Butene</b>
Answer. (d)	(c) Propyne
181. Gasoline with an octane number of 90 is	(d) 2-Butyne
equivalent in knocking	Answer. {b)
characteristics to a mixture of heptane and	188. Which of the following compounds is most
isooctane of the following composition	stable?
	(a) Ethylene
(a) 20% heptane + $80\%$ isooctane	(b) 2,3-Dimethyl- I-butene
(b) 90% o neptane + 10% isooctane	(c) Propylene
(c) $80\%$ neptane + $20\%$ isooctane	(d) 2-Butene
(d) 10% neptane + 90% isooctane	Answer. [D]
Answer. (a)	(a) water
182. A fuel with octane number 90 means it is as	(d) water (b) other alcohol
$g_{000}$ as a mixture of:	(b) ethylacolol (c) ammonia
(a) 90% isoocialle + 10% in-neptane (b) 00% n Hontone + 10% inconstance	(d) carbon totrachlorido
$\{D\}$ 90% II-Reptaire + 10% isooctaire	Answer (d)
(d) 2 Litre of $90\%$ isolocialle + 2 litre of Delizene (d) 2 Litre of $90\%$ hontone + 2 litre of	<b>190</b> Which statements about alkenes and
(u) 2  Little of 90% II-fleptatie + 2 little of bonzone Amure (a)	alkanos of corresponding chain longths is true?
183 Which bydrocarbon molecule with eight	(I) Alkenes have slightly lower melting points
carbons would produce the lowest octane	than alkanes
number (i.e. the most knocking) when used as a	(ii) Alkenes have slightly higher melting points
fuel in automobile?	than alkanes
(a) 2-Methyl-3 3-dimethylpentane	(iii) Alkenes have higher boiling points thanalkanes
<i>(h)</i> n-Octane	(iv) Alkenes have lower boiling points than
(c) 2-lsopropy1pentane	alkanes
(d) 2-Methylheptane	(a) I and III
Answer. (b)	(b) 11 and 1 V
184. Which of the following compounds is used	(c) I and IV
as an antiknock compound?	(d) II and III
(a) Ethyl lithium	Answer (a)
<i>(b)</i> Tetraethyllead	
(c) Ethyl acetate	
Impacting Lives	

**191.** How do *the* melting points of *trans* isomers compare to the cis isomers for alkenes? (a) *cis* isomers have higher melting points (b) *trans* isomers have higher melting points (c) both have similar melting points (d) no consistent trend is observed Answer. [b] **192.** Why do *trans* isomers of alkenes have lower boiling points than *cis* isomers? (a) *trans* isomers have better symmetry *(b) cis* isomers have better symmetry (r) trans isomers are less polar (d) cis isomers are less polar Answer. (c) 193- The major product of acid-catalyzed dehydration of 3-pentanol is: (a) l-Pentene *{b)* 2-Methyl-1-butene (c) 2-Pentene (d) 3-Methyl-1-butene Answer. (c) **194.** Ethylene is obtained from ethyl bromide by (a) Simple heating (b) Hydrolysis (c) Dehydrohalogenation (d) Nucleophilic substitution Answer. (c) **195.** The dehydrohalogenation of 2bromobutane with alcoholic KOH gives mainly (a) 2-Butene (b) 2-Butyne (c) 1-Butene (d) l-Butyne **Answer**. (a) **196.** In the reaction of propene with HCI, H + ion acts as the: (a)electrophile (b) carbonium ion (c) nucleophile (d)carbanion Answer. (a) **197.** When bromine attacks the double bond in propene, which of the Following ions are Formed in the first stage of the attack?

(a) CH<sub>3</sub> – CH – CH<sub>2</sub>Br  $(h) CH_3 - CH_2$ (c) CHP - CH - Cl-t BrBr (d) CHP –  $CH – \overline{CH2}$ Answer. (a) 198. In the reaction of ethylene with H20 in the presence of sulfuric acid, which one adds across the double bond first? (a)H+ (b) H. [c] HO-(d) sulfate ion Answer (a) 199. Why does a H+ ion attacking a carboncarbon double bond add to the carbon with the least number of substituents? (a) the reaction is resonance stabilized. (b) the hybrid geometry favors this process. (c) nucleophilic tends to attack stable centers ofnegative charge. (d) a more stable carbonium ion is generated. Answer. (d) 200. Propene reacts with bromine to form 1,2dibromo propane. This is an example of (a) Nucleophilic addition (b) Electrophilic addition (c) Nucleophilic substitution [d] Electrophilic substitution Answer. {b} 201. Baeyer's reagent is (a) dilute KMnO4 (b) HC1 + ZnC12 (c) Br2 in CCl4 (d) NH2NH2 Answer (a) 202. In the addition of HX to a double bond, the hydrogen goes to the carbon that already has more hydrogens is a statement of (a) Hund's rule [b] Markovnikov's rule [c] Huckel rule (d) saytzeff rule Answer. {b} 203. Markovnikov's addition of HBr is not

applicable to (a)Propene (b) 1-butene

Teekay Page 17	Educational Consult
(c) 1-pentene	Answer. (c)
(d) 2-butene	211. 2-Methylpropene reacts with HBr to give
Answer. (d)	(a) Bert-Butyl bromide
204. In the reaction of CH3CH2CH=CH2 with	<i>(b i</i> isobutane
HCL the H of the HCl will become attached to	(c) n-Butyl bromide
which carbon?	(d) None of these
$(2) C_{-1}$	Answer (a)
$f_h)C_2$	212.2 Butono roacts with HBr to give
$\left( c \right) \left( c$	(a)   Promobutano
	(b) 2.2 Dibromobutane
$\left( u \right) C-4$	[0] 2,5-DIDI OIIIODULAILE
Answer, (a)	[c] 2-Bromobulane
205. Which of the following compounds will	[d] 2,2-Dibromobutane
react most readily with bromine in CCI4?	Answer (c)
(a) CH3CH2CHI	213. Which of the following alkenes reacts with
(b) (CH3)3CH	HBr in the presence of peroxide
(c) CH3CH=CH2	to give <i>an</i> ti-Markovnikov's product?
(d) (CH3)4C	(a) l-Butene
Answer. (c)	(b) 2,3-Dimethyl-2-butene
<b>206,</b> The disappearance of the purple color of	(c) 2-Butene
KMn04 in its reaction with alkene is known as	(d) 3-Hexene
(a) Markovnikov test	Answer. (a)
<i>(b)</i> Grignard test	214. Propene reacts with HBr in the presence of
(c) Baeyer test	a peroxide to give
(d) Wurtz test	(a) n-Propyl bromide
Answer. (c)	<i>[b i</i> Allyl bromide
<b>207.</b> Which of the following will give a negative	(c) isopropyl bromide
test when treated with bromine in carbon	(d) Vinvl bromide
tetrachloride?	Answer. (a)
(a) Butane	<b>215.</b> 2-Methylpropene reacts with HBr in the
(h) 2-Butene	nresence of neroxide to give
(c) 1 3-Butadiene	(a) A primary alkyl bromide
(d) 2-Butyne	(b) A secondary alkyl bromide
Answer (a)	(c) A tertiary alkyl bromide
<b>208</b> Ethylene reacts with HI to give	(d) A viginal dibromida
(a) indepthane	
(a) 1000ethalle	Answer. (a)
(b) 2,2-011000ethalle	<b>210.</b> 2-Methyl-2-butene reacts with HBr in the
(c) Norre of these	presence of peroxide to give
(c) None of these	(a) A primary alkyl bromide
Answer. (a)	(b) A secondary alkyl bromide
209. Which of the following reagents will react	(c)A tertiary alkyl bromide
with propene*	(d) A vicinal dibromide
(a) Hot alkaline KMnO4	Answer. {b)
<i>(b)</i> Sodium metal	<b>217.</b> Hydration of 2-methyl-1-propene (with
(c) Cold dilute HNO3	H2O/H2SO4) gives:
(d) LiAlH4	(a) CH3CH2CH2OH
Answer. (a)	<i>(b)</i> (CH3)3C0H
210. Which of the following compounds will	(c) CH3CH2CH2CH2OH
have zero dipole moment?	(d) (CH3)2CH0H
(a) cis-1,2-dibromoothylenctb)	Answer. (d)
1,1-dibromoethylene	<b>218.</b> A compound reacts with steam in the
(c) <i>trans</i> - 1,2-dibromoethylene	presence of concentrated H2504 to give
(d) all of these	isopropyl alcohol. The compound could be
Impacting Lives	

Teekay	Page 18	Educational Consult
(a) Alkane		[b] 2-methyl-2-butene
<i>{b)</i> Alkyne		(c) 2-butene
(c) Alkene		(d) 2-methy1propene
(d Alene		Answer (b)
Answer. (c)		<b>226.</b> A hydrocarbon, C6H12, on ozonolysis gives
219. Propene reacts with Cl2 in H20 to give	ve	only one product which <i>does</i> not give silver
(a) 1-Chloro-2-propanol		mirror with Tollens' reagent. The hydrocarbon is
<i>{b</i> ) 2-Ch1oro-2-propanol		(a) 2,3-Dimethyl-2-butene
(c) I-Chloro-l-propanol		(b) 2-hexene
(d) 2-Chloro-1-propanol		(c) 2-Methyl-2-pentene
Answer. {a)		[d] 3-Hexene
220. 1-Butene reacts with Cl2 in H20 to give	ve	Answer. (a)
(a) 1-Chloro-2-butanol		227. Combustion of an alkene with sufficient
<i>{b)</i> 2-Chloro-2-butanol		oxygen will produce
(c)1 -Chloro-1-butanol		(a) carbon dioxide and water
(d) 2-Ch1oro-1-butanol		(b) carbon monoxide and water
Answer. (a)		(c) only carbon dioxide
<b>221.</b> Catalytic hydrogenation of 3-methyl-1-		(d) only carbon monoxide
butene gives:		Answer. (a)
(a) Isobutane		228 Which of the following are correct
<i>{b</i> 2,2-Dimethylbutane		statements concerning unsaturated
(c) 2-Methylbutane		hydrocarbons?
(d) 2,3-f1imethylbutane		(a) All unsaturated hydrocarbons are insoluble
Answer. (c)		in water.
<b>222.</b> Which of <i>the</i> following alkenes gives of	only	<i>(b)</i> All unsaturated hydrocarbons are soluble in
acetic acid on oxidation with hot		nonpolar solvents.
concentrated KMnO4.		(c) All unsaturated hydrocarbon is more
(a) Ethylene		dense than water.
(b) l-Butene		(d) (a) & (c)
(c) Propene		Answer. (d)
(d) 2-Butene		<b>229.</b> How many electrons are involved in a
Answer. (d)		carbon-carbon double bond?
<b>223.</b> Which of the following compounds will	l give	(a) 1
only acetaldehyde on		[b] 2
ozonolytic:'		(c) 3
(a) 1-butene		(d) 4
(b)Acetylene		Answer. (d)
(c) 2-Butene		<i>230.</i> Which of the following statements about
(d) Ethylene		ethene, C2H4, is <i>incorrect?</i>
Answer. (c)		(a) The H-C-H bond angles are approximately
<b>224.</b> Which of the following alkenes will give	ea	109.5°.
mixture of acetone and		(b) There is a total of five sigma bonds.
formaldenyde on ozonolysis?		(c) The carbon atoms are $sp2$ hybridized.
(a) 2-butene (b i 2 mothyl 2 hytono		(d) The H—L—H bond angles are approximately
(c) 1 hutone		
(d)2 methylpropene		Answer (C)
(u)2-methylpropene		<b>231.</b> Which of the following statements about
<b>225</b> Which of the following all anos will give	VA 2	(a) All nine atoms lie in the same plane
mixture of acetone and	vea	(a) The compound has a cis and transisoner
acetaldehyde on ozonolysis?		(c) There is a total of eight sigma hands
(a) l-hutene		(d) All the carbon atoms are sn? hybridized
		(u) mi me carbon atoms are sp2 hybridized.
Impacting Lives		

Teekay P	age 19	Educational Consult
Answer. (c)		<b>239.</b> Which of the following compounds have
232- Which of the following is capable of		planar molecules?
exhibiting cis-trans isomerism?		(a) 1,3-Butadiene
a) l-butene		(b) Dimethyl ether
b) l-pentene		(c) l-Butene
c) ethene		(d) Alene
d) 2-butcnc		Answer. (a)
Answer. (d)		<b>240.</b> Which of the following molecular formulas
<b>233.</b> Which of the following is the most stabl	e	will correspond to an alkene with two double
alkene?		bonds?
[a] trans-3-hexene		[a] C4H10
(b) cis-3-hexene		(b) C5H12
(c) 1-hexene		[c] C6H10
d) (Z)-3-hexene		(d) C8H16
Answer. (d)		Answer. (c)
<b>234</b> . Which of the alkyl chlorides listed below		<b>241</b> 1 3-Butadiene reacts with bromine to
indergoes dehydrohalogenation in		mainly give
he presence of a strong base to give 2-nenten	ρ	(a) 3 4-Dibromo-1-butene
is the only alkene product?		(h) A-Bromo-1-butenc
a) 1-chloropentane		(b) 4-Dibhomo-2-butene
(h) 2-chloropentane		(d) 1 Bromo 2 butono
(c) 2 chloropontano		(u) 1-biolilo-2-butelle
d) 1 chloro 2 rnothylbutano		Allswer. (c)
a) 1-chioro-2-rheuryibutane		<b>242.</b> What descriptive term is applied to the
<b>Allswell</b> (C) <b>27</b> What is the major and wat from the asi	4	(g) conjugated diana
<b>235.</b> What is the major product from the action of 2 mothed 2	u-	( <i>u</i> ) conjugated diene
catalyzed hydration of 2-methyl-2-		[b] cumulated diene
Jentene?		(c) Isolated diene
a) 2-methylpentane		(d)alkynyl diene
b) 2-methyl-1-pentanol		Answer. (c)
c) 2-methyl-2-pentanol		<b>243.</b> What descriptive term is applied to the
d) 2-methyl-3-pentanol		type of diene represented by 2,4- hexadiene?
Answer. (c)		(a) conjugated diene
<b>236.</b> 1,2-Butadiene has		(b) cumulated diene
(a) o nly <i>sp</i> hybridized carbon atoms		(c) isolated diene
(b) only sp2 hybridized carbon atoms		(d) alkynyl diene
c) only ip3 hybridized carbon atoms		Answer. (a)
(d) <i>sp, sp2,</i> sp3 hybridized carbon atoms		<b>244.</b> Which of the following statements are
Answer. (d)		correct concerning delocalized
<b>237.</b> How many (sigma) bonds are there in		electrons?
CH2=CH—CH=CH2?		(a) Electrons do not belong to a single atom.
(a) 3		(b) Electrons are not confined to a bond betweentwo
(b) 6		atoms
(c) 9		(c) Electrons are shared by three or more atoms.
(d) 12		(d) (a), b) and (c)
Answer. (c)		Answer. (d)
<b>238.</b> Propadiene, CH2=C=CH2, is		<b>245.</b> What is the hybridization of the central
(a) a planar compound		carbon of allene 1.2-propadiene)?
(b) u cumulated diene		(a) sp
c) an isolated diene		(b) sp2
(d) a conjugated diene		(c) $n3$
Answer (h)		(d) n
		Answer (a)

246. What compound results from the 1,4-(d) Alkynes are generally more reactive than addition of one equivalent of HBR to alkenes. 1,3-butadiene? Answer. (d) (a) l-bromo-1-butene 253. Which of the following describes a triple *(b)* 2-bromo-2-butene bond? (c) 4-bromo-1-butene (a) two *sigma* bonds and two pi bonds (d) 1-bromo-2-butcnc (b) one sigma bond and one pi bond (c) two *sigma* bonds and one *pI* bond Answer. (d) 247. Rank the following dienes in order of (d) onesigma bond and two pi bonds Answer. (d) increasing stability: 254. What two atomic orbitals or hybrid atomic trans-1, 3-pentadiene, cis-1,3-pentadiene, 1,4orbitals overlap to form the carbon-carbon a pentadiene and 1,2-pentadiene. **Answer.** 1,2-pentadiene <1,4-pentadiene < *cis*bond in ethyne? [a1 sp 3—sp2 1,3-pentadiene • trans-1,3-(*b*) sp2—sp2 pentadiene (c) s—s **248.** A triple bond consists of [d] sp—sp *{a*) 2 *sigma* bonds and 1 *pi* bond Answer. (d) *(b)* 3 *sigma* bonds 255. Which of the following statements is not (C) I sigma bond and 2 *pi* bonds true about propyne, HC-CCH3? (d) 3 pi bonds (a) It contains six *sigma* bonds. Answer. (c) (b) It contains three pi bonds. **249.** The bond angles associated with the hybrid (c) The H-C-H bond angle is about 109.S°. orbitals of a carbon involved in a (d) The C-C—C bond angle is 180°. triple bond is Answer. (b) (a) 180º 256 How many distinct terminal alkynes existwith [b] 120º (c) I 09º a molecular formula of C5H8? (d) 45° (a) 1 (b) 2 **Answer**. (*a*) (c) 3 250. How many electrons are involved in a (d) 4 carbon-carbon triple bond? Answer. (b) (a) 1 257. How many distinct internal alkynes exist *{b)* 2 with a molecular formula of C6HI 0? (c) 3(a) 1 (d) 6 (b) 2 Answer. (d) (c) 3 251. Which of the following improperly (d) 4 describes the physical properties of an alkyne? Answer. (c) (a) relatively nonpolar (b) slightly insolublee in water 258 Which is the correct order of decreasing acidity in the following compounds? (c) less dense than water A. H20 B. CH3CH3 C. NH3 D. CH2=CH2 E. HC-CH (d) insoluble in most organic solvents (a) A > E > C > D > BAnswer. (d) (b) A > E > D > B > C252. Which of the following statements correctly (c) E > A > C > B > Ddescribes the general rreactivity of alkynes? (*d*) A > C > E > D > B(a) An alkyne is an electron-rich molecule and **Answer** (a) therefore reacts as a nucleophile. 259. Which of the following compounds is the *{b "the o* bonds of alkynes are higher in energy major product when 1-hcxvne is than the bonds aand are thus non reactive treated with excess HBr? (c) Unlike alkenes, alkynes fail to undergo (a)1,1-dibroinohexane electrophilic addition reactions. (b) I, I -dibromo hexene

(c) 1,2-dibromohexene **266.** Which of the following hydrocarbons has (d) 2,2-dibromohexane acidic hydrogens? [a] l-Butene Answer. (d) 260. The reagent needed to convert 2-butyne to (b) l-Butyne cis-2-butene is : (c) 2-BuLene (a) H2/Pt(d) 2-Biitvne (b) H2/Lindlar's catalyst Answer. (b) **267.** Lindlar's catalyst is (c) Li/NH3 (d) Na/NH3 (a) LiAliI4 Answer. (b) (b) Pd/BaSO4 in Quinoline 261. Among the compounds water, 1-butyne, 2-[c] NH2NH2 butyne, and ethane, which are stronger acids [d] HCl/ZnC12 than ammonia? Answer (b) 268. The higher reactivity of an alkene or (a) l-butyne and ethane alkyne, as com payed to an alkane, is due to (b) water and 4 -butyne (c) water and ethane [a] sigma bonds (d) l-butyne and 2-butyne (b) pi bonds (c) hydrogen bonds **Answer**. *{b)* **262**, What is the product when 3-heptyne reacts (d) None of these with the Lindlar catalyst? Answer. (b) 269. Which of the following statements about (a) 2-methyl-2-heptene *(b)* trans-3-heptene alkenes and alkynes is correct? (c) cis-2-methyl-3-hexene (a) Alkynes are reduced more readily than (d) cis-3-hexene alkenes. (b) Alkynes are reduced *less* readily thanalkenes. Answer. (d) (c) Alkynes and alkenes are reduced with **263.** In the addition of hydrogen bromide to speed, alkynes, which of the following Speices is believed to be an intermediate? (d) Alkynes and alkenes cannot be reduced. Answer. (a) (a) vinyl anion 270. Which of the following sequences regarding *{b )* vinyl cation acid-strength is correct? (c) vinyl radical (a) HC-CH > CH3CH20H > CH3C00H (d) carbene (b) HC-CH > CH3C0OH > CH3CH2OHAnswer. [b] 264. Starting with 2-butene, which of the (c) CH3COOH > DC-CH > CH3CH2OH(d) CH3COO H > CH3CH 20H > HCSCH following is the best method for Answer preparing 271. 1.2-Dichloroethane reacts with excess of 2-butyne? (a] HBr, H2/Ni, Zn/H+ NaNH2 to form *{b)* HBr; Zn/H+; H2/Ni (a) Vinyl chloride {c) Br2/CCI4; 2n/H+; H2/N i (b)Ethylene (c) Ethyl chloride (d) Br2/CCl4; 2NaNH2 (d) Acetylene Answer. (d) Answer. (d) **265.** In the following hydrogenation reactions; 272. Propyne is formed by  $HC \equiv CH \xrightarrow{H_2N_i} CH_2 = CH_2 \xrightarrow{H_2N_i} CH_3CH_3$ (a) Polymerization of acetylene The hybrid Slate of the carbon atom changes from (b) Reaction of acetylene with methane (a)  $sp^3 \rightarrow sp^2 \rightarrow sp$ (c) Reaction of acetylene with methyl chloride (<I) Reaction of sodium acetylic with methyl (b) sp  $\rightarrow$  sp<sup>2</sup>  $\rightarrow$  sp<sup>3</sup> chloride (c)  $sp^2 \rightarrow sp^3 \rightarrow sp$ Answer. (d) (d) sp<sup>3</sup>  $\rightarrow$  sp  $\rightarrow$  sp<sup>2</sup> Answer. (b)



#### KINDLY VOTE FOR **TEEKAY** AS THE NEXT FACULTY OF SCIENCE **PRESIDENT**

Teekay	Page 23	Educational Consult
273. Which of the following compounds of	on	280. 2-Butyne undergoes catalytic
hydrolysis gives acetylene?		hydrogenation in the presence of Landler's
(a] CaC2		catalyst to give
(b) Mg2C3		(a) 2-Butene
(c) AI4C3		(b) Butane
(d) Cu2Cl2		(c) I -Butene
Answer. [a]		(d) 2-Mcthylpropenc
274. Which of the following compounds	on	Answer. (a)
hydrolysis gives propyne?		281. Addition of two moles of IICI to propyne
(a) CaC2		gives:
<i>{b</i> ) Mg2C3		[a] 2,2-Dichloropropane
(c) Ai4C3		(b) 1,3-Dichloropropane
(d) Cu2Cl2		(c] 1,2-Dichloropropane
Answer. (c)		(d) None of these
275- which one of the following compou	nds will	Answer. (a)
(a) decolorizes dilute cold		<b>282.</b> Propyne reacts with aqueous H2SO4 in the
KMn04; (b) decolorizes bromine water; a	and (r)	presence of HgS04 to form
gives a white precipitate with ammoniac	al	(a) acetone
AgNO3 solution.		(b) l-propanol
(o) 1-Hexene		[c] acetaldehyde
(b) 1-Hexyne		(d) 2-propanol
(c) 2-Hexene		Answer. (a)
(d) 2-Hexyne		<b>283.</b> Acetylene reacts with water in the presence
Answer. (b)		of sulfuric acid and mercuric
276. Ethylene and acetylene can be		sulfate to give
distinguished by using		(a) Acetone
(s) Bromine in CCI4		(b) Acetic acid
(b) Tollens' reagent		(c) Formaldehyde
(b) Baeyer's reagent		(d) Acetaldehyde
(d) Phenylhydrazine		Answer. (d)
Answer. [b]		284. Which of the following compounds will
277. I-Butyne can be distinguished from 2	2-	react with ammoniacal silver
butyne by using		nitrate?
(a) potassium permanganate		(a) l-Butene
<i>{b)</i> bromine in CCI4		(b) I-Butyne
(c) Tollens' reagent		(c) 2-Butene
(d) chlorine in CC14		(d) 2-Butyne
Answer. (c)		Answer (b)
278. I-Butyne reacts with		<b>285.</b> Which of the following compounds does
(a) NaNI-12 (b) $D_{11}^{11} U_{12}^{12} D_{14}^{12}$ and $U_{22}^{12} D_{14}^{12}$		not reach with ammoniacal AgNU3
(d) All of these		[a] Acetylene
(d) All of these		(b) I-Butyne
Answer. (u)	o hu	(c) Propyne (d) 2 Putrme
279. Propyre can be converted to proper	le by	(u) 2-Butylle
using (a) H2 + Landlar's catalyst		Allower (U) 296 Which of the following compounds reserve
(a) $HZ + Lander S catalyst(b) NH2N H 2$		with ammoniacal suprous chloride
(c) H2 + Pt catalvet		to give a procipitato*
$(C_{J})^{11} = T + Calalyst$ N H 2NH2 $\pm V \cap H$		(a) 1 Butono
$\frac{1}{2} \frac{1}{2} \frac{1}$		$(a)  l_{\text{Butype}}$
		(D) - Duly lie (c) 2-Butono
		(c) 2-Dutene

Teekay	Page 24	Educational Consult
t4 2-Butyne		[b] cyclohexane < cyclopentane < cyclobutane <
Answer. {b)		cyclopropane
<b>287.</b> Ozonolysis of 2-butyne gives		(c] cyclopentane < cyclobutane < cyclopentane <
(a) Formic acid		cyclopropane
<i>{b)</i> Propanoic acid		(d) cyclopentane < cyclopropane < cyclobutane •
(c) Acetic acid		cyclohexane
(d) Butanoic acid		Answer. (b)
Answer. (c)		294. Which of the following has two equatorial
288. Which alkyne yields propanoic acid as	sthe	alkyl substituents in its most stable
only product upon treatment with		conformation?
ozone followed by hydrolysis?		[a] 1,1-dimethylcyclohexane
(a) l-Butyne		(b) cis-1,2-dimethy1cyclohexane
(b) 2-Hexyne		(c] cis-1,3-diethylcyclohexane
(c) 1-Pentyne		(d) crs-1,4-diethylcyclohexane
(d) 3-Hexyne		Answer. (c)
Answer. (d)		295. Which one of the following is not a metal
289. When acetylene is passed through he	ot iron	catalyst for the hydrogenation of an
tube at 400°C, it gives		alkene?
(a) Benzene		(a) Pd
(b) Toluene		(b) Pt
(c) o-Xylene		(c) Na
(d) Mesitylene		(d) Ni
Answer. (a)		Answer. (t)
290. When propyne is passed through hot	: iron	296. What is(are) the product(s) in the Pd-
tube at 400°C, it gives		catalyzed hydrogenation of 1,2 - dimethyl
(a] Benzene		cyclopentene?
(b) Toluene		(a) troor-1,2-dFnethylcyclopentane
<i>{c)</i> m-Xylene		(b) eis-1,2-dlmcthy1cyclopentanc
(d) Mesitylene		(c) a mixture of <i>trans</i> and cix-1,2-
Answer. (d)		dimethyl cyclopentane
291. The monomer for Neoprene is		(d) 1,1-dimethylcyc1opentane
(a) Isoprene		Answer. (b)
<i>{b)</i> acrylonitrile		297. Which of the following is not a possible
(C) Chloroprene		reaction of a carbocation?
(d) 1,3-butadiene		(a) addition of a nucleophile
Answer. [c]		(b) rearrangement to a more stable carbocation
292. A compound, C4H6, reacts with bron	nine	(c) addition of a proton to form an alkane
and forms a white precipitate with ammoni	acal	<i>{d)</i> loss of a <i>§</i> -hydrogen to form an alkene
silver nitrate solution. It reacts with dilut	e	Answer. (c)
H2SO4 in the presence of		298. Addition of HCl to 3-mcthyl-1-pcntenc
mercuric sulfate to form 2-butanone. The		gives two products. One of these is 2-chloro-3-
compound could be		methyl pentane. What is the other product?
(a) l-Butyne		(a) 1-Chloro-3-methylpentane
<i>(b)</i> I-Butene		(b) 3-Chloro-3-methylpentane
(c) 2-Butyne		(c) 3-Chloro-2-methylpentane
(d 2-Butene		(d) 2-Chloro-2-methylpentane
Answer. (a)		Answer (b)
293. Which of the following correctly ran	ks the	299. Predict which of the following alkenes
cycloalkanes in order of		reacts the fastest with HCI?
increasing ring strain per methylene?		(a) CH3CH2CH2CH2CH=CH2
(a) cyclopropane < cyclobutane < cyclohe	exane	(b) cis-CH3CH2CH=CHCH2CH3
<cycloheptane< td=""><td></td><td>(c) trees-CH3CH2CH=CHCH2CH 3</td></cycloheptane<>		(c) trees-CH3CH2CH=CHCH2CH 3
Impacting Lives		

Teekay	Page 25	Educational Consult
(d) (CH3)2C=CHCH2CH3		of the ozone layer in the upper stratosphere?
Answer. (d)		(a) CH4
300. The hydroboration-oxidation r	eaction can	<i>(b)</i> CC12F2
be characterized as theto a	n alkene.	(c) CF4
(a) anti-Markovnikov syn addition of	water	(d) CH2CI2
(b) anti-Markovnikov anti addition of	water	Answer. {c)
(c) Markovnikov syn addition of water		<b>307.</b> Which of the following reagents cannot be
(d) Markovnikov anti addition of wa	ater	used to prepare an alkyl chloride from an
Answer. (a)		alcohol?
301. A compound, CIS H24, is reacted	with	(a) HCl + ZnCl2
excess hydrogen using a metal catalys	t. One	[b] SOC12
equivalent of the compound consum	ned three	(c] NaCl
equivalents of hydrogen. How many	rings did	[d] PCI5
the original compound have:'		Answer. (c)
(a) 1 only		<b>308.</b> 2-Propanol reacts with KBr and
(b) 2 only		concentrated H2504 to give
(c) 3 only		(a) l-Bromopropane
(d) None of these		(b) 1,2-Dibromopropane
Answer. (‹i)		(c) 2-Btomoppopane
<b>302.</b> The reaction of 1-butene with br	omine,	(d) 2,2-Dibromopropane
Br2, in aqueous solution <b>gives</b>		Answer. (c)
Primarily 1-bromo-2-butanol. Ident	ify the	<b>309.</b> The best way to prepare 1,2-
nucleophilic species in the reaction.		Dibromoethane is to treat:
(a] Br2		(a) Ethylene with Br2
(b) Br—		[b] Acetylene with excess of HBr
(c) H20		(c) Ethylene with excess of HBr
(d) H0Br		(d) 1,2-Dtchloroethane with Br2
Answer. (c)		Answer. (o)
303. Which brief statement most ac	curately	<b>310.</b> Alkyl halides undergo
describes why alkenes react the way	v they do?	(a) Electrophilic substitution reactions
(a) C=C double bonds are weak.		<i>(b)</i> Electrophilic addition reactions
(b) A n bond is lost but a stronger	bond is	(c) Nucleophilic substitution reactions
gained.		(d) Nucleophilic addition reactions
(c) c=C double bonds are unstable.		Answer (c)
(d) C=C w bonds are attacked by nuc	leophiles.	<b>311.</b> n-Propyl iodide reacts with sodium
Answer. (a)	_	ethoxide to give :
304. Which of the following poison	ous gas is	(a) CH3CH20CH2CH3
formed when chloroform is exposed	to Light	{b) CH3CH20CH2CH2CH 3
andair?		(c) CH3CH20CH3
(a) Mustard gas		(d) CH30 CH2CH2CH3
(b) Carbon monoxide		Answer. (a)
(c) Phosgene		<b>312.</b> I-Bromobutane reacts with alcoholic KOH
(d) Chlorine		to mainly give
Answer. (c)		(a) I-Butene
305. Freon-12, CCI2F2, is used as a		(b) 2-Butene
(a) Local anesthetic		(c) I-Butanol
(b) Dry-cleaning agent		[a] 2-Butanol
(c) Keirigerant		AIISWEF. {D]
(a) Disinfectant		<b>513.</b> 2-Bromodutane react alcoholic KUH to
Allswer. (C)	ndahaa	mainly give
500. Which of the following compou	nus nas	(a) 1-Butene
been suggested as causing depletion	11.	[D] 2-Butene
Impacting Lives		

Teekay Page 25	Educational Consult
(c) 1-Butanol	(a) Formic acid
(d) 2-hutanoI	(b) Sodium formate
Answer. (b)	(c] Acetic acid
<i>314.</i> Isopropyl bromide reacts with alcoholic	(d) Sodium acetate
KOH to give	Answer. {b}
(a) Propene	<b>321.</b> Which alkyl halides react most readily by
(b) isopropyl alcohol	nucleophilic substitution?
(r) Propane	(a) CH3CH2Cl
(d) n-Propyl alcohol	(b) Cl 13Cli21
Answer. (a)	(c) CH3CH2Br
315. 2,2-Dich1oropropane reacts with aqueous	[d] CH3CH2F
KOH to give	Answer. {b]
(a) 2,2-Propanedio1	<b>322.</b> Which of the following factors influence
(b) Propanal	whether a reaction will proceed by an SN1, SN2,
(c) Acetone	El, or E2 mechanism?
(d) Propene	(a) Structure of the alkyl halide
Answer. (c)	(b) Solvent
316. 1.1-Dichloropropane reacts with aqueous	(c) Concentration of reagents
KOH to give	(d) Native of the nucleophile
(a) 1,1-Propanediol	[e] All of these
{b Acetaldehyde	Answer. (e)
(c) Propanone	<b>323.</b> Which compound reacts most rapidly by an
(d) Propyne	SN1 mechanism.'
Answer (b)	(a) Methyl chloride
317. Which of <i>the</i> following compounds will not	(b) isopropyl chloride
give a white precipitate with alcoholic AgNO3.	(c) Ethyl chloride
(a) Bromobenzene	(d) tertbutyl chloride
(b) l-Bromopropane	Answer. (d)
(c) Ethyl bromide	<b>324.</b> Which of the following compounds would
(d) 2-Bromopropane	react most rapidly in an SN2
Answer. [a]	reaction?
318. Which of the following compounds will give	(a) CH3CH2I
a white precipitate with	(b) CH2=CH-I
alcoholic AgNO3?	(c) (CH3)2CHI
(a) Vinylbenzene	(d) (CH3)3CI
(b) Chlorobenzene	Answer. (a)
(c) Vinyl chloride	<b>325.</b> There are 8 isomers that have the
(d)Allyl chloride	molecular formula C5H11Br. How many of these
Answer. (d)	are tertiary alkyl bromides?
3t9. Carbylamines test involve heating a	(a) 1
mixture of	<i>(b)</i> 2
(a] alcoholic KOH, methyl iodide, and sodium	(c) 3
metal	( <i>d</i> )8
<i>(b)</i> alcoholic KOH, methyl iodide, and primary	Answer. (a)
amine	<b>326.</b> Which of the following statements is
(c) alcoholic KOH, chloroform, and primary	correct for a saturated alkyl halide?
amine	(a)the C—X bond results from overlap of the s
(d) alcoholic KOH, methyl alcohol, and primary	orbital of carbon and the <i>p</i> orbital
amino	of the halogen (X)
Answer. (c)	[b] the C—X bond results from overlap of the p
<b>320.</b> When chloroform is heated with aqueous	orbital of carbon and the p orbital
NaOH, it gives	of the halogen (X)
Impacting Lives	

Teekay Page 26	Educational Consult
(c) the C—X bond results from overlap of the sp3	(b) The rate of reaction depends on the
orbital of carbon and the p	concentration of the alkyl halide.
orbital of the halogen (X)	(c] The rate of reaction of a particular alkyl
(d the C—X bond results from overlap of the <i>sp2</i>	bromide depends on the steric
orbital of carbon and the p	accessibility of the carbon of the C-Br bond.
orbital of the halogen (X)	(d) All alkyl iodides react more rapidly than all
Answer. (c)	alkvl chlorides.
<b>327</b> . Which of the following best describes the	Answer. (d)
carbon-chlorine bond of an alkyl chloride?	<b>333.</b> Which of the following correctly reflects
(a) nonnolar: no dinole	relative stabilities of carbocations?
(h) nolar: 6+ at carbon and 6— at chlorine	[a) $3^{\circ}$ allylic > $2^{\circ}$ > $1^{\circ}$ henzylic
(c) polar: $6-$ at carbon and $6+$ at chlorine	(h) methyl > 2° henzylic > 3°
(d) ionia	(c) hencylic > vinul > 1°
(d) IOIIIC	$ \begin{array}{c} (c) & S & Dellzylic > Villyl > 1 \\ (d) & 2^{0} & alledia > 2^{0} & coincul \\ \end{array} $
<b>Allswer</b> . (D) <b>220</b> Which of the following is a second are aller	$(u) \ge any n \le 2$ viny $(d)$
<b>328.</b> Which of the following is a secondary alkyl	Allswer. (u)
	<b>334.</b> Which of the following species is most
(a) CH3Br	reactive in a SN2 reaction?
$\{D\}$ (CH3)3CBr	(a) CH3CH2CI
(c) (CH3)2CHBr	(b) CH3CH2Br
(d) (CH3)2CHCH2Br	[c] CH3CH2I
Answer. (c)	(d) CH3CH2F
<b>329.</b> How should CH3CHClCH2CH 3 be classified?	Answer. (c)
(a) primary alkyl halide	335. Which of the following is the best leaving
(b) secondary alkyl halide	group?
(d) tertiary alkyl halide	(a) F-
<i>(d)</i> quaternary alkyl halide	(b) Cl-
Answer. (b)s	(c)Br-
<b>330.</b> Which of the following will have the lowest	(d) I—
boiling point?	Answer. (d)
(a) CH3Cl	336. Which of the following is the strongest
<i>(b)</i> CH4	nucleophile in an aqueous solution?
(c) CH2CI2	(a) Ho-
(d) CHCl3	(b) Cl—
Answer. (b)	(c) Br—
<b>331.</b> Which of the following is not correct	(d) 1—
concerning substitution and elimination	Answer. (d)
reactions of alkyl halides	337. Which of the following is the best
(a) The electrophile replaces the leaving group.	nucleophile in water?
(b) Compounds containing electron-donating	$(a) \mid -$
groups bonded to a sp3 hybridized	(b) CH3SCH3
carbon undergo substitution and elimination	(c) CH30CH3
reactions	[d] C]-
(c) The electronegative atom is replaced by	Answer (a)
another atom or group in substitution	338 Which of the following compounds will
reactions	undergo an SN2 reaction most
(d) (a) and $(b)$	readily7
Answer (d)	(a) (CH3)3(CH2)
<b>232</b> Which of the following statements	(h) (CH2)3CCI
concorning SN2 reactions of all all holides is not	
correct?	
(a) The rate of reaction demonds on the	
(a) The rate of reaction depends on the	Answer. (a)
concentration of the nucleophile.	

Teekay	Page 27	Educational Consult
<b>339.</b> Which of <i>the</i> following is the rat	te law for	[b] 2-chloropentane
SN1 mechanisms?		(c) 3-ch1oropentane
(o] Rate = k {Alkyl Halide} {Nucleophile}		(d) 1-chloro-2-methylbutane
(b) Rate = k[Nucleophile]		Answer (a)
(c) Rate = k [Alkyl Halide]		<b>346.</b> Predict the two most likely mechanisms for
(d) Rate = $k1[Alky]$ Hal ide] + $k2[Nuc]$	leophile]	the reaction of 2-iodohexane with sodium
Answer. (c)		ethoxide.
<b>340.</b> Which of the following factors l	has to effect	[a] SN2 and SN1
on the rate of SN1 reactions?		(b) E1 and E2
(A) the nature of the alkyl halide		(c) $5N2$ and $E2$
(h) the nature of the leaving group		[d] El and SN1
(C) the concentration of the alkyl hal	ide	Answer. (c)
(D) the concentration of the nucleon	nile	<b>347</b> Predict the two most likely mechanisms
Answer (d)		which occur when 2-iodobeyane is heated in
<b>341</b> Which 1s the most reactive all w	l halido in a	othanal
SN1 reaction?	i nanue in a	(a) SN2 and SN1
		(a) $SINZ$ and $SINI$
(h) CH2Pr		(D) EI and E2
(D) CHODI		$\{C\}$ SNZ and EZ
(l) CH2Cl		(d) EI and SN1
		Answer. (d)
		<b>348</b> Grignard reagents do not show any
<b>342.</b> Which of the following halides is	most	reaction with
reactive in an E2 reaction with		(a) Alkoxy alkanes
sodium		(b) Alkanones
methoxide?		(c) Alkyl alkanoates
(A) (CH3)3CCH2I		(d) Acy1 halides
(B) (CH3)2CHCHICH3		Answer. (a)
(c) (CH3)2CHCH2Br		349. The 6rfgnard reagent, CH3CH2MgBr, can
(d) (CH3)2CHCH2CI		be used to prepare
Answer. [b]		(a) Ethane
<i>343.</i> Which of <i>the</i> following halides i	s least	<i>{b)</i> 3-Ethyl-3-pentanol
reactive in an E2 reaction with		(c) Propanoic acid
sodium methoxide?		(d) All of these
(a) (CH3)3CCH2I		Answer (D)
<i>{b)</i> (CH3)2CHCHICH3		<b>350</b> . Which is the best reagent to accomplish the
(c) (CH3)2CHCH2 Br		following conversion?
(d) (CH3)2CHCH2CI		(a) Conc. H2SO4
Answer. (a)		<i>{b)</i> Na
344. Dehydrohalogenation of 2-brom	nobutane in	(c) Conc. HCl
the presence of a strong base		(d) Mg, then H2O
proceeds via which of the following	mechanistic	Answer. (d)
pathways?		351. What is the major product of the following
(a) SN1		reaction?
<i>(b)</i> SN2		(A) l-Butanol
(c) El		(b) Butanal
(d E2		(c) 2-Butanol
Answer. (d)		<i>Id j</i> Butanone
345- Which of the alkyl chlorides listed	d below	Answer. (c)
undergoes dehvdrohalogenation in		352. Ethyl magnesium iodide reacts with
the presence of a strong base to give	2-pentene	formaldehyde to give a product which on acid-
as the only alkene product?	- pontono	hydrolysis forms.
(a) l-chloropentane		(a)an aldehyde
(a) i emoropentane		(a jan anuchyuc
Impacting Lives		

Teekay	Page 28	Educational Consult
<i>(b)</i> a primary alcohol		(b) Benzyl alcohol
(c] a ketone		(c) Methyl phenylcarbinol
(d) a secondary alcohol		(d) Benzoic acid
Answer. (b)		Answer (c)
353. Ketones react with Grignard reage	ents to	359. Which of the following is the strongest
form ap addition product which on hydr	rolvsis	base?
gives a	5	(a) HOMgBr
(al Primary alcohol		(b) H2O
<i>(b)</i> Tertiary alcohol		(c) CH3OH
(c) Secondary alcohol		(d) CH3MgBr
(d) Ketal		Answer. (d)
Answer. (b)		360. Which of the following is correct?
354. n-Propyl magnesium bromide on t	reatment	(a) When a Grignard reagent reacts with a
with carbon dioxide and further		ketone, the addition product is a
hydrolysis gives:		primary alcohol.
(a) Acetic acid		(b) When a Grignard reagent reacts with a
<i>(h)</i> Propanoic acid		ketone, the addition product is a
(c) Butanoic acid		secondary alcohol
(d) Formic acid		(c) When a Grignard reagent reacts with a
Answer. (c)		aldehyde the addition product is a tertiary
355 Which of the following compounds	will	alcohol
react with methyl magnesium iodide follo	wed by	(d) None of the above are correct
acid-hydrolysis to give ethyl alcohol	weaby	Answer. (d)
(a) Ethylene		361 Which of the following alcohols can be
(h) Acetaldehyde		prepared by the reaction of methyl formate with
(c) Formaldehyde		excess Grignard reagent?
(d) Acetone		(a) 1-nentaool
Answer. (c)		(b) 2-pentanol
356 Which of the following gives a terti	arv	(c) 3-pentanol
alcohol when treated with Grignard read	gents?	(d) 2-methyl-2-pentanol
	Serrer	Answer. (c)
Ŭ		362. The number of structural isomers of
(a) H-C-H		alcohols with molecular formula
ç,		C3H70H is
		(a) s
$({}) CHI - C - H$		(h) 4
0		(c) 3
()		(d) 2
(d) None of these		Answer (d)
Answer(d)		363. The number of structural isomers of
357 Which of the following compounds	will	alcohols with molecular formula
roact with mothyl magnesium	W111	CAHQOH is
bromide to give text butyl elected?		
(a) Acetyl chlorido		$\begin{bmatrix} a \end{bmatrix} 5$
(a) Acetone		$\begin{bmatrix} D \end{bmatrix}$ 4 $\begin{bmatrix} c \end{bmatrix}$ 3
(c) Isopropyl alcohol		(d) 6
(d) Acetaldebyde		Answor (b)
Answor (b)		364 Why do alcohols have boiling points much
358 Phenyl magnesium bromido roacte	with	higher than hydrocarbons of similar molecular
acotaldohudo to form an addition produ	willi ct which	
undergoes acid-hydrolysis to give		(a) Alcohols have greater van der Waals
(a) Diphonyl carbinol		attraction forces
		מננו מננוטוו וטו נבז.
Impacting Lives		

Teekay Pag	ge 29	Educational Consult
<i>(b)</i> Alcohol molecules have greater molecular		371. Which of the following compounds is the
symmetry.		least soluble in water?
(c) Hydrogen bonds must be broken in the		(a) HOCH2CH2OH
process of volatilization.		(b) CH3CH2CH20H
(d) Alcohols must overcome greater ionic force	es	(c) CH3CH2OH
in the process of volatile Action.		(d) CH3CtI2CH 2CH20H
Answer. (c)		Answer (D
365 The high boiling points of alcohols as		372 Methanol is known as
compared to the corresponding		(a) Rubbing alcohol
alkanes are due to		(b) Grain alcohol
(a) Hydrogen honding		[c] Wood alcohol
(h) Heavy oxygen atom		[d] Denatured alcohol
(c) Water solubility		Answer C
(d) None of those		272 Poctified spirit is
(u) Nolle of these		(a) 100% Ethanol
Allswel. (d) 266 Which of the following has the highest		(h) 000% Ethanol
boiling point?		(b) 10006 Mothanol
bolling point?		(c) 100% Methanol
(a) dietnyl etner		
(b) n-Butyraldenyde		Ansiver. (D
(c) s-propyl chloride		374. Grain alcohol is another name for
(d) n-Butyl alconol		(a) Methyl alconol
Answer. (d)		(b) isopropyl alcohol
367. Which of the following has the highest		(c) Ethylaiconol
boiling point?		(d) n-Propyl alconol
(a) CH3CH2CH2CH2CI		Answer. (c)
(b) HOCH2CH2CH2CH2OH		375. Lucas reagents is
(c) CH3CH 2CH2CH2OH		(a) HCI/NaN02
(d) CH3CH2OCH2CH3		(b) H2/Pd
Answer. (b)		(c) HCI/ZnCI2
368. Which of the following compounds has the	ie	(d) H2/Pd/BaSO4
highest boiling point		Answer. (c)
(a)		376. Lucas test is used to determine the type of
<i>(b)</i> CH3CH2CH2OH		(a) alcohols
(c)		[b) acids
(d) CH3CH2OCH3		(c) amines
Answer. (b)		(d) carbohydrates
369. Which of the following has the highest		Answer. (A)
boiling point?		377. Which of the following gives positive
(a) CH3CH2OH		lodoform test:
(b) СНЗСН2СН 2СН2ОН		(a) 1-Propano1
(c) CH30H		(b) 2-PropanoI
(d) CH3CH2CH2OH		(c) 3-Propanol
Answer. [b]		(d) None of these
370. Rank the following substances in order of	of	Answer. {b)
increasing boiling point (lowest —) highest):		378. Which of the following reagents will
(1) CH3CH2CH2CH2OH, (2) (CH3)2CH0CH3, (3)	)	replace —OH group by a halogen atom?
(СНЗ)ЗСОН		(A) HOCI
(a) $(1) < (2) < (3)$		(B) Br2
(b) $(2) < (1) < (3)$		(C)SOCI2
(C)(2) < (3) < (1)		(D) I2
(d) (3) < (2) < (1)		Answer. (c)
Answer. (c)		
The section I to an		
Impacting Lives		

Teekay	Page30	Educational Consult
379. The compound which reacts most read	ily	386. Ethanol on heating with concentrated
with Lucas reagent is	5	H2SO4 at 170°C gives
(a] CH3CH2CI		(a) ethylene
(b) (CH3)2CH0H		(b)ethyl hydrogen sulfate
(c) CH3CH20		(c) diethyl ether
(d) (CH3)3C0H		(d) diethyl sulfate
Answer. (d)		Answer. (a)
380. Which of the following compounds re	acts	387. The major product of acid-catalyzed
slowest with Lucas reagent at		dehydration of 2-butanol is
room temperature?		(a) 2-Butene
(a) l-butanol		(h) 2-hutyne
{b] 2-propanol		(c] l-Butene
(c) 2-butano1		(d) l-Butyne
(d) 2-methyl-7-propanol		Answer (a)
Answer (a)		388 Which is tile best reagent for carrying out
381 Which of the following compounds w	ill	the following conversion?
react fastest with Lucas reagent?		(a) LiAlH4
(a) l-propanol		(b) Conc H2SO4
(h) 2-methyl-1-propanol		(c) $H^2/N_i$
(c) 2-propanol		$(d) N_2 0 H$
(d) 2-methyl-2-propanol		Answer (b)
Answer (d)		389 Which of the following reaction mixtures
382 Which of the following compounds rea	cts	will give an organic product which
fastest with HBr?		is ionic?
(a) l-propanol		(a) CH3CH20H + PCIS
<i>(h)</i> 2-methyl-1-propanol		(h) CH3CH20H + KBr + H2504
{¢] 2-propanol		(a) CH3CH2OH + Na
(d) 2-methyl-2-propanol		(d) CH3CH2OH + SOCI2
Answer. (d)		Answer (c)
383 When ethanol is treated with sodium	metal	390 Hydroboration-oxidation of propene gives
Soor when eduator is freuted with southin	lifetai	(a) Isopropyl alcohol
(a) Sodium ethoxide is formed		(b) n-Propyl alcohol
<i>(b)</i> The sodium is oxidied		(c) isobutyl alcohol
(c) The acidic hydrogen in reduced		(d) tert-Butyl alcohol
(d) All these occur		answer {h]
Answer. (d)		391 Hydrohoratinn-oxidation of 2-
384. Which of the following alcohols will g	vive a	Methylpropene gives
vellow precipitate of iodoform	,	(a) 2-Methyl-2-propanol
with jodine and dilute NaOH solution?		(h) 1 2 3-Propanetriol
(a) 1-Propanol		(c) 2-Methyl-1-Fronanol
(b) 2- Propanol		(d) 1.2-Propanediol
(r) 1-Butanol		Answer. (c)
(d) 2-Methyl-2-propanol		392. Which of the following compounds will not
Answer. [b]		be <i>easily</i> oxidized?
385. Ethyl alcohol can react with concentry	ated	(a) Primary alcohol
H2S04 to give		(b)Secondary alcohol
(a) Ethylene		(c) Tertiary alcohol
<i>{b}diethyl ether</i>		(d) Aldehyde
(c) Ethyl hydrogen sulfate		Answer. (c)
(d)All of these		393. Which of the following is most resistant to
Ànswer. (d)		oxidation?
		(a) CH3CH20H
	1	

Teekay	Page 31	Educational Consult
<i>{b)</i> (CH3)2CHOH		(b) Carbanions
(c) HOCH2CH20H		(c) free radicals
(d)(CH3)3C0H		(d) Carbines
Answer. (d)		Answer (a)
394. Oxidation of a primary alcoh	ol with	<b>401.</b> The acid-catalyzed dehydration mechanism
produces		for alcohols is best described as
(a) a carboxylic acid		a(n):
<i>(b)</i> an ether		(a) El
(c) a ketone		(b) E2
(d) an ester		(c) SN1
Answer. {a)		[d] SN2
395. Oxidation of a secondary alco	ohol with	Answer. (a)
K2Cr2O7/H+ produces		<b>402.</b> When ethanol is heated with concentrated
(a) a carboxylic acid		H2S04, a gas is produced. Which
(b) a ketone		of the following compounds is formed when this
(c) an aldehyde		gas is treated with bromine in
(d) an ester		CCI4?E2
Answer. [b]		(a) Bromomethane
396, Isopropyl alcohol reacts wi	th acidic sodium	(b) 1,2-Dlbromoethane
dichromate to give		[c] Bromoethane
(a) Acetaldehyde		(d) 1,1,2,2-Tetrabromoethane
(b) Acetic acid		Answer. {b]
(c) Propionic acid		<b>403.</b> What is the functional group in an alcohol?
(d) Acetone		(a) Carbon-carbon double bond
Answer. (d)		<i>[b]</i> NH2
<b>397.</b> Isopropyl alcohol can be co	nverted to	(c) OH
acetone by treatment with		(d) COOH
(a) HCI/ZnCl2		Answer. (c)
(b) Na2Cr2O7/H 2504		<b>404.</b> What is the IUPAC name for the following
(c) NaOH		structure?
(d) LiAlH4		CH3CH(CH3)CH2CH(0H)CH3
Answer. (b)		(a) 4-methyl-2-pentano1
<b>398.</b> Which of the bonds in Ethy	l alcohol	(b) 2-methyl-2-hexanol
(CH3CH20H) will undergo heter	olytic	(c) 2-methyl-4-pentano1
bond fission most readily?		(d) 2-methyl-3-pentanol
(a)0—H		Answer. (a)
(B) C—H		<b>405.</b> Explain why the water molecule has a bent
(c) C—0		shape and a bond angle less than
(dC-C		109.5°.
Answer. (a)		Answer. The electron repulsion between the
399. The mechanism of dehydra	tion of an	two lone pairs of electrons on the
alcohol to give an alkene involve	es	oxygen of water causes the 0-H bonds to
Formation of:		squeeze close together.
(a) Carbonium ions		<b>406.</b> Which of the following compounds does not
{b )Carbanions		have the molecular formula
(C) Free radicals		C4H140?
(d) Carbines		(a) 2-hexanol
Answer. (a)		(b) 3-methyl-2-pentano1
<b>400.</b> The mechanism of dehydra	tion of an	(c) 3-methyl-3-pcntanol
Alcohol to give an ether		(d) cyclohexanol
involvesformation of:		Answer. (d)
(a) Carbonium ions		

Teekav Page 32	Educational Consult	
<b>407</b> . Which of <i>the</i> following is a secondary	nentene?	
alcohol?	(a) 2-methy1pentane	
(o] 3-ethyl-2-methyl-1-pentanol	(h) 2-methyl-1-pentano1	
th) 3-ethyl-2-methyl-2-hentanol	(c) 2-methyl-2-nentanol	
(c) 3-methyl-2-heyanol	(d) 2-methyl-3-pentanol	
(d) 1-hevanol	Answer (c)	
Answer (c)	A14. In rold countries athylana glycol is added to	
And Which of the following is cor-butyl alcohol?	water in car radiators. This	
(a) $C112C1[2C[1,2C]]20[1$	holps to	
(b) CH3CH2CH12CH2CH2	(a) Poduce the viscosity	
	<i>(a)</i> Reduce the viscosity	
	(c) I ower the freezing point	
	[d] Lower the heiling point	
Allower. (D) A110 What is the hybridization of the owner	<i>[a]</i> Lower the boining point	
atom in CH3CH20 H2	Allswei. (C) Als Ethylene ryvile undergoes acid-bydrolysis	
$t_{\bullet}$ gn	to form	
(b) A cn 2	(a) Ethylono glycol	
$\left( c \right) \sin^2 c$	(b) Formic acid	
(c) sp2	(c) Ethyl alcohol	
Answor (di	(d) Acotic acid	
<b>410</b> Where are the two long pairs of electrons of	Answer (2)	
the ovvgen atom in an alcohol	416 Ethylene given undergoes ovidation with	
molecule located?	hot acidic KMn04 to form ·	
(ol in two <i>n</i> orbitals	(a) Formic acid	
th) in two sp orbitals	<i>(b)</i> Formaldehyde	
(c) in two sp2 orbitals	(c) Acetic acid	
(d) in two sp3 orbitals	(d) Acetaldehyde	
Answer. (d)	Answer. (a)	
<b>411.</b> What is the strongest intermolecular force	<b>417.</b> Ethylene glycol reacts with excess of PCI5	
present in liquid ethanol?	to give	
(o) induced dipole-induced dipole	(a) Chloroethane	
<i>tb1</i> dipole-dipole, specifically hydrogen bonding	(b) 1,2-Dichloroethane	
(r) dipole-dipole, but not hydrogen bonding	(c] Hexachloroethane	
tH ion-dipole	[d] 1,3-Dichloroethane	
Answer. [bj	Answer. {b)	
412. Which of the following is the best method	418. 1,2-Ethanediol reacts with anhydrous zinc	
for preparing CH3Br?	chloride to form	
(a)CH30H + Br—	(a) Ethylene	
<i>{b)</i> CH30H + HBr	[b] Acetaldehyde	
(c) CH30H + Br2	(c) Atctylcne	
(d) CH30H + NaBr	(d) Ethyl chloride	
Answer. (b)	Answer. {b )	
413. Which of the following reagents can be	419. When ethylene glycol is heated with	
used to oxidize 1° alcohols to	concentrated HN03, it forms	
aldehydes?	(a) Oxalic acid	
(a) KMn04	(b) Ethylene oxide	
(b) Mn02	(c) Dioxane	
(C) K2Ur2U/	(a) Flieth ylene glycol	
	Answer. (a)	
AllSWEF. (d)	<b>420.</b> Which of the following is used as an	
12. What is the major product from the acid-	anurreeze:	
catalyzeu nyulation of 2-methyl-2-	(a) Euryrene grycor	
Teekay	Page 33	Educational Consult
--	----------	--
<i>{b</i> ) Glycerol		(b) НОСН2СН2СН2СН2ОН
(c) Diethyl ether		(c) CH=CH—CH=CH2
(dj Picric acid		(d) CH3CH2CH2CH2OH
Answer. (a)		Answer (b)
421. When glycerol is heated with oxalic a	icid at	428. Ethers are
260°C, it gives		(a) Lewis acids
(a) 1,2-Propancdio1		(b) Neutral
<i>{b</i> } Vinyl alcohol		(c) Lewis bases
(c) 1,3-Propanediol		(d) Can not be predicted
(d) Allvl alcohol		Answer. (c)
Answer. (d)		<b>429.</b> Ethanol is isomeric with
<b>422.</b> When glycerol is heated with potassi	um	(a) Dimethyl ether
hvdrogen sulfate (KHS04), it forms		(b) Ethanal
(a) Acrolein		(c) Diethyl ether
(b) Acetic acid		(d) Propanone
(c) Allyl alcohol		Answer (a)
(d) Propionic acid		430. Diethyl ether and methyl propyl ether are
Answer. (a)		(a) Conformational isomers
423, Glycerol en warming with a small am	nount	(b) metamcrs
of hydriodic acid gets converted		(c) Geometrical isomers
to		(d) Enantioiners
(a) Propene		Answer. {b j
<i>(b)</i> 3-lodopropene		431. The reaction of a sodium alkoxide with an
(c) 1,3-Diiodopropane		alkyl halide is called
(d) 2-[odopropane		(a) Wurtz-Fittig reaction
Answer. (b)		(b) Perkin reaction
424. When glycerol Is treated with a mixt	ure of	(e) Williamson's synthesis
concentrated HN03 + H2SD4, it		(d) Aldol condensation
forms		Answer. (c)
(a) Nitroethane		432. Ethyl bromide reacts with sodium
<i>(b)</i> l-Nitropropane		methoxide to form
(c) Nitrogiycerine		(a) Diethyl ether
(d 2-Nitropropane		(b) Ethyl methyl ether
Answer. (c)		(c) Dimethyl ether
425. In the manufacture of dynamite, one	of the	(d) n-Propyl alcohol
chemicals used is		Answer. (b)
(a) Glycerol		433. Ethyl alcohol reacts with concentrated
(b) Glycerol triacetate		H2S04 at 140°C to lorm
(c) Glycerol trinitrate		(a) Acetone
(d) Glycerol) triiodide		(b) Ethylene
Answer. (c)		(c) Diethyl ether
<b>426.</b> Which of the following compounds is	s least	[d] Acetic acid
soluble in water?		Answer. (c)
(a) Glycerol		434. Ethers are kept in brown bottles because
{b ] Ethyl alcohol		(a) Brown bottles are cheaper than colorless
(c) Etnylene glycol		clear bottles
(u) Etnyi chioride		( <i>D</i> ) Ethers absorb moisture
Allswel. (U) 427 Compound (A) reacts with sudium m	otal ta	(d) Ethers evaporate readily
form one mole of U2 The	ciai ill	(u) Ethers are oxidized to cxpJ sive peroxides
101  III 0 II e III 0 I e 0 I f 12. I II e compound (A) can be		Answer (a)
(a) CH2CH2CH-CH2		
(a) 01130112011-0112		

Teekay Page34	Educational Consult
435. Ethers are stored in brown bottles. This is	Answer. (c)
because on <i>ex</i> post <i>re</i> to air and light ethers are	442. Diethyl ether reacts with excess of hot
converted to	concentrated HI to form
(a) Peroxides	(a) CH3CH21 + CH3CH20H
(b) Oxonium ions	(b) Only CH3CH2OH
(c) Ozonides	(c) $CH3CH20H + CH2=CH2$
(d) Electrophil es	(d) Only CH3CH21
Anwer. {a)	Answer. (d)
436. Ethers can be freed from peroxides by	443. Which is the best reagent to accomplish the
treatment with	following conversion?
(a) Ferrous salt	CH3CH20CH2CH3 CH3CH2Br
(b) Sodium carbonate	(a) Br2 in CCI4
(c) Ferric salt	(b) NaBr
(d) Sodium bicarbonate	(c) Br2 in H20
Answer. (c)	(d) Cont HBr
437. Which of the following statements is false?	Answer (d)
(a) Diethyl ether has been used as a general	444. Which of the following pairs of compounds
anesthetic	will not form hydrogen bonds
(b) Ethyl alcohol is present in all alcoholic	with each other?
beverages	(a) CH3OH and CH3CH2OH
(c) Methyl alcohol is produced by fermentation	(b) CH3SH and CH3CH2SH
of sugars	(c) CH30CH3 and Cfl3CH 20CH2£H3
(d) Ethylene <b>glycol is a</b> common antifreeze for	(d) CH3CO0H and H2O
automobiles	Answer. (c)
Answer. (c)	445. The compound with the lowest boiling
438. Ethers react with cold concentrated H2SO4	point is
to form	(o) H20
(a) Oxonium salts	(6) CH3CH2OH
(b) Alkenes	(c) CH3OCH3
(c) Aikoxides	(d) CH3CH2CH3
(d) Zwitterions	Answer. (d)
Answer. [a]	446. Which of the following would have the
439. Which of the following reagents readily	highest boiling point?
react with ethyl methyl ether?	(a) 1-Butano1
(a) NaOH	(b) Butane
(b) Conc HI	(c) I -Butene
(c) KMn04	(d) l-Butyne
(d) H20	Answer (a) · Alcohols of the same chain length
Answer. (b)	as alkanes, alkenes, and alkynes
440. Diethyl ether <i>reacts</i> with sodium metal	have higher fioi1ing points due to their ability to
togive	hydrogen bond.
(a] Ethanol	447. Which of the following would have the
(b) Sodium ethoxide	highest boiling point?
(c) Ethanal	(a)Dimethyl ether
(d) Nothing happens	<i>(b)</i> Diethyl ether
Answer. (d)	[c] Ethyl methyl ether
441. Diethyl ether on heating with excess	[d] Diisopropyl ether
concentrated HI gives	Answer. (d). Ethers follow the same trend as
(a) Methyl iodide	alkanes, so diisopropyl ether will have the
(b) Isopropyl iodide	highest boiling point because it has the highest
(c) Ethyl iodide	molecular weight
(a) n-Propyl iodide	[

Teekay	Page35	Educational Consult
448. Which of <i>the</i> following functional	group	[b] Ethylamine
series is ranked according to		(c) 2-Aminoethanol
increasing boiling points?		(d) Acetamide
(a) diethyl ether, ethane, ethanol, ethan	ethiol	Answer (c)
(b) ethane, ethanol, diethyl ether, ethane	ethiol	<b>456.</b> Which of the following is closest to the C—
(c) ethane, diethyl ether, ethanethir>l, et	hanol	0—C bond angle in CH3—0—CH3?
(d) diethyl ether, ethane, cthancthiol, ct	hano1	(a) 180°
Answer. (c)		( <i>b</i> ) 120°
449. The following compounds have ide	entical	(c) 109.5°
molecular weights. Which would have the	ne lowest	(d) 90°
boiling point?		Answer. (c)
(a) l-Methoxypropane		<b>457.</b> When dipentyl ether is treated with excess
(b) l-Butanol		HI, through what type of
(c) 1,1-Dimethylethanol		mechanism(s) does the major product result:
(d) 2-Butanol		(a) SN2
Answer. (a)		(b) SN1
450. Cyclic ethers with three-membered	d ring are	(c) El
called		(d) E2
(a) Lactoncs		Answer. (a)
(b) Oxiranes		458. Which of the following is xot a property of
(c) Alkoxides		thiols (RSH)?
(d) Epoxy resins		(a) They <i>are</i> all solids
Answer. (b)		(b) They can be oxidized to disulfides
451. Ethylene reacts with oxygen in the		[c) They have foul odors
presence of silver catalyst at 300°C to form	n	(d) They are weak acids
(a] Diethyl ether		Answer. (a)
(b) Ethylene <b>glycol</b>		<b>459.</b> Thiols are alcohol analogs In which the
(c) Ethyl alcohol		oxygen has been rcplaced by Sulfur (e.g.,
(d) Ethylene oxide		CH3SH). Given the lact that the S—H bond is less
Answer. (d)		polar than the U—H bond,
<b>452.</b> Ethylene oxide reacts with HBr to g	give	which of the following statements corriparing
(a) I-Bromoeunanon (b) Ethyl hydridd	*	thiols and alcohols is correct?
(a) 2 Promosthanol		(a) Hydrogen bonding forces are weaker in
(d) Ethylono glycol		(h)Undragen hending ferrers are strongen in
Answer [c1		thick
<b>453</b> Acid-catalyzed hydrolysis of ethyler	ne ovide	(c) Hydrogon bonding forces would be the same
vields	ic oxide	(d) No comparison can be made without
(a) CH3CH2OH		additional information
(b) HOCH2CH2OfI		Answer. (a)
(c) CH3CH 2CH20H		<b>460</b> . Which of the following has the lowest
(d) H0CH2CH2CH2OH		boiling point?
Answer. [b]		(a) CH3CH20H
<b>454.</b> The product of the reaction of ethy	lene	(b) CH3CH2CH2SH
oxide with acidic methanol is		(с) НОСН2СН2ОН
(a) CH30CH2CH20H		(d)CH3CH2CH2OH
(b) CH3CH2CH2CH2OH		Answer (b)
(c) HOCH2CH20H		461.Which of the following is least soluble inwater?
(d) CH3CH20CH2CH3		(a) CH30H
Answer. (a)		<i>[b]</i> CH3CH20H
<b>455.</b> Ethylene oxide reacts with ammon	ia to give	(c) CH35H
(a) l-Aminoethanol		

Teekay	Page 36	Educational Consult
(d) HOCH2CH2OH		468. Which of the following will have the highest
Answer(c)		boiling point?
462. n-Buty1 bromide reacts with	NaSH to give	(a) Methanal
(a) CH3CH2 CH2SH	-	(b) Ethanal
(b) CH3SCH3		(c) Propanal
c CH3CH 2CH2CH2SH		(d) Butanal
(d) CH3CH2SCH 2CH3		Answer. (d). The boiling points increase with
Answer. (c)		molecular weight. All of the given compounds
<b>463.</b> The carbon atom of a carbon	yl group is	are aldehydes. Butanal has the highest molecular
(a) <i>sp</i> hybridized		weight.
{b ] sp2 hybridized		<b>469.</b> What property of low-molecular weight
(c) sp3 hybridized		aldehydes and ketones accounts for
(d) None of these		the magnitude of their boiling points?
<b>Answer.</b> (b). Whenever carbon is	bonded to	(a) The ability to form strong H-bonds between
three other atoms or groups, it uses	s p2 hybrid	their molecules.
orbitals to form its bonds.	1 5	(h) The ability of the carbonyl oxygen to form H-
<b>464.</b> Which statement about the c	arbonyl group	honds with other carbonyl
is NOT true?	<b>J D I</b>	groups.
(a) The carhonyl carbon is sp2 hv	bridized.	(c) The ability of the polar carbonyl group to
(b) The bond angles among the three	e atoms	attract other polar molecules.
attached to the carbonyl carbon ar	'e	(d) The ability of the carbonyl group to attract
120'.	-	electrophiles and form honds
(c) The three atoms attached to th	e carbonvl	Answer. (c)
carbon form a nonplanar geomet	rv.	<b>470</b> . Primary alcohols have boiling points
(d) The carbonyl group forms res	onance	thatare the
structures.		corresponding aldehydes.
Answer. (c)		(a) Lower Can
<b>465.</b> Acetone contains		(b) Higher than
(a) nine <i>sigma</i> bonds plus one a b	ond	(c) About the same
<i>{b)</i> ten sigma bonds	$\mathbf{X}$	Answer. [b ]. For compounds of similar
(c) eight sigma bonds plus two n	bonds	molecular weights, boiling points increase with
(d) nine pi bonds plus one <r bond<="" td=""><td></td><td>the degree of hydrogen bonding. <i>Remember :</i></td></r>		the degree of hydrogen bonding. <i>Remember :</i>
Answer. {a)	Ψ.	Alcohols form strong hydrogen bonds and will
<b>466.</b> Formalin is		boil at a higher temperature than the
(a) 10% solution of formaldehyde	in water	corresponding aldehydes.
(6) 20% solution of formaldehyde	in water	471. The melting points of aldehydes and
(c) 40% solution of formaldehyde	e in water	ketones tend to :
(d) 80% solution of formaldehyde	in water	(a) decrease with increasing molecular weight.
Answer. (c)		(b) increase with increasing molecular weight.
467. Which of the following will ha	ve the highest	(c) remain u nchangcil with increasi rig molecular
boiling point?		weight.
(a] Propanone		[d] be unpredictable due to resonance.
<i>(b)</i> 2-Pentanone		answer. (b)
(c) Butanone		472. Which action best accounts for the
(d) 2-Hexanone		solubility of aldehydes and ketones in water?
Answer. (d). Boiling points increas	e with	(a) Polar interactions between solute molecules.
molecular weight. The higher the		(b)H-bondng between solute moiecles.
molecular weight, the higher is the	boiling poi nt	(c)Vanderwnahforces
All of the givcn compou rids arc		(d)H-bondinghctvccnsolutuandsclvcnt
ketones. 2-Hexanone has the high	est molecular	molecules
weight.		Answer. (d)
		<b>473.</b> Ketones are prepared by the oxidation of

Teeka	iy

**Educational Consult** 

**479.** Which of the following reagents will react

readily with both aldehydes and

ketones?

(a) Grignard reagent

(a) Primary alcohol *{b)* Secondary alcohol (c) Tertiary alcohol (d) None of these Answer. (b) 474. Which of the following do you consider to be suitable for obtaining acetone (CH3COCH3)? (a] Heating CH3CH2CH20H with acidic Na2Cr207 *{b]* Passing over heated copper (c) Oxidation of CH3CH2CH3 with concentrated HN03 (d) Heating CH3CH=CH2 with dilute H2504 Answer. (b) 475. Cyclopentanol undergoes oxidation to give -(a) Cyclopentene *{b )* Cyclopentanone (c) Cyclopentane (d) Cyclopentanal **Answer**. (b) **476.** Which statement about the carbonyl group of ketones and aldehydes is true? I. It can attract nucleophiles. II. It can attract electrophiles. II1. It tends to **undergo** addition reactions. IV. It tends to undergo substitution reactions. (a) I and III (b) II and IV (c) I, II, and III (d) I, III, and IV Answer. (c) 477. Which of the following compounds reacts with sodium bisulfite and ammoniacal silver nitrate solution? 0  $(a) CH_3CH_2 - C - OH$   $(b) CH_3CH_2 - C - H$ (c) CH<sub>3</sub>CH<sub>2</sub>CH=CH<sub>2</sub> (d) CH<sub>3</sub>C  $\equiv$  CCH<sub>3</sub> Answer. (b) **478.** Acetone reacts with HCN to form a cyanohydrin. It is an example of (a) Electrophilic addition *{b ]* Electrophilic substitution (c) Nucleophilic addition (d) Nuclcophilic substituition **Answer**. (c)

(b) Fehling's reagent (c) Tollens' reagent (dJ SchifFs reagent Answer. (a) 480. Boiling acetaTdehyde [CH3CHO] reacts with chlorine gas to give CH<sub>3</sub>CH<sub>2</sub>-C-OH 0 н (b) CH<sub>3</sub>CH<sub>2</sub> - C(c) CH3CHCI2 (d) CHC13 Answer(b) 481. The reaction of ethanal with one equivalent of methanol and a trace of an acid will give (a) Acetal *{b]* Hemiacetal (c) Ketal (d) Hemiketal Answer. (b) 482. Acetone undergoes reduction with hydrazine in the presence of NaDH to form propane. This reaction is known as (a) Clemmensen redilction *[b*)Wolf-Kishner reduction (c) Rosenmund reduction (d) Reformatsky reaction Answer.(d) 483. Acetone undergoes reduction with zinc amalgam in HCI to form propane. This reaction is known as (a) Clemmensen reduction (b) Wolf-Kishner reduction (c) Rosenmund reduction (d) Aldol condensation Answer. (a) 484. Cannizzaro reaction is not given by (a) Formaldehyde *{b )* Trimethylacetaldehyde (c) Acctaldeyhyde (d) Benzaldehyde Answer. (c) **485.** When formaldehyde is treated with S0% Na0H solution, it undergoes (a) Cannizzaro reaction

(b) Wurtz reaction

Teekay	Page 38	Educational Consult
(c) Aldol condensation		493. Aldehydes can be distinguished from
(d) Hydrolysis		ketones by using
Answer. (a)		(a) Ammoniacal AgNO3 solution
486. The reduction of a ketone		(b) Phenylhydrazine
(o) always gives a primary alcoh	ol	(c) Saturated NaHSO3 solution
(6) always gives a secondary alco	ohol	(d) Thionyl chloride
(c) always gives a carboxylic acid		Answer. (0)
(d) always gives a ketal		<b>494.</b> The appearance of a silver mirror in
Answer. {b]		Toilets' test indicates the presence of:
487. Reduction of acetaldehyde y	vith H2/Ni	(a) an aldehyde
gives		(b) a ketone
(a) Ethyl alcohol		(c) an alcohol
(b)Acetic acid		(d) an alkene
(c)Ethylene		Answer (a)
(d)Ethane		495 Acetaldehyde on treatment with Totters'
Answer (a)		reagent gives a precipitate of
488 On reduction with LiAlH4 w	which of the	(a) Ag
following compounds could vield	an ontically	$(a) \operatorname{Ag}(b) \operatorname{AgNO3}$
active compound?	anoptically	(c) Cu20
(a)Propagal		(d) none of these
(6) Propanane		Answer (2)
(c) Butanal		496 Which of the following compounds will give
(d) Butanone		a positive test with Folling's
Answer (d)		a positive test with renning s
490 The reduction of cyclobovan	ono with	(a) Formaldohudo
LiAllua (or NoPUA) will give		(b) A cotopo
(a) an alcohol		(b)Acetone (c)E1 $u$ 1 $a$ cotato
(d) all dicollol		(c) ElyIdeeldee
(D) all Olgaliic aciu		(u) necuc aciu
(d) a hamiltatal		AllSWell (d)
		497. Acetaidenyde on treatment with tening s/
Answer. (a)		(a) Cu
490. Aldenydes undergo oxidatio	n with	$(s) \cup u$
KMII04/H+10 give		(D) Cu20
(a) Alcollois		(C) CUU (d) None of these
(D) Actions		(u) None of these
(c) Ketones (d) Asida:		Allswell. (D)
(u) Actus:		Fabling's solution sizing a red
Answer. (a)		Fening's solution giving a red
491. Oxidation of acetaidenyde w	/itm	precipitate?
Nazurzu//H+ gives		
(b) Acetic acid		
(b) Etheral		
(C) Ethanol (d) Apptana		$\bigcup_{i=1}^{n}$
(d) Acetone	/	(d) cH3 C-CH3
Answer(b)		
<b>492.</b> A nyarazone will result from	i the reaction of	499 Whtch of the following compounds will give
nyurazine with		an lodoform test?
(a) a pilenoi $(b)$ an aldebude		an rouoronni test:
(b) an alach al		(a) DEFIZUIC ACIU (b) Ethanol
(c) an acconor (d) an acid		(U) EUIAIIOI (c) Ronzul chlorida
(u) an actu		(d) Methanol
Answer. {DJ		
Impacting Lives		

Answer. (b) 500. Which of the following compounds does not give a positive iodoform test : (a) Ethanol {b] Ethanal (c) methanol (d) Propanone Answer. (c) **501.** Which of the following compounds does not give iodoform test on treatment with 12/NaOH? (a) Ethanol (b) Propanone (c) 2-Propano1 (d) Butanone Answer. (d) **502.** Which of the following compounds will give a positive iodoform test? (a) Benzaldehyde (b) 2-Pentanone (c) 3-Hexanone (d) 3-Pentanone Answer (b) 503. An organic compound (A), C3H80, on oxidation gives (B), C3H60. The compound (A) could be (a) an aldehyde (b) a ketone (c) an alcohol (d) an ester Answer. (c) 504. An organic compound (A), C3H80, on oxidation gives (B), C3H60. The compound (B) could be (a) a carbonyl compound (b) an alcohol (c) a carboxylic acid (d|)an ether Answer. (a) 505. An unknown com pound gavn a positive Tollens' test. Treatment of the unknown with 12/NaOH gave a solid which was identified as iodoform. The unknown was? (a] Ethanal (b) Acetone (c) Propanal (d) Acetophenone Answer. (a) **506.** A compound (A) gave a positive iodoform test, but did not reduce silver nitrate in ammonia solution. Compound (A) could be

Õ Π (a) CH<sub>3</sub> -C Õ b)CH3CH2 C<sup>---</sup> CH<sub>2</sub>CH3 Ο – CH2CH3 CH2 Answer. (d) **507.** Which of the following characterizes the reactions of aldehydes and ketones? (a) electrophilic addition (b) elcctrophi1ic substitution (c) nucleophilic acyl substitution (d) nucleophilic addition: free radical addition Answer (d) 508. Why do aldehydes undergo nucleophilic Addition reactions while esters undergo nucleophilic acyl substitution reactions? (a) The carbonyl carbon of an ester is more electrophilic than that of an aldehyde.

(b) Aldehydes are more sterically hindered than esters.

(c) Once the nucleophile adds to an aldehyde, the tetrahedral Intermediate is too stertcally hindered to eliminate one of the attached groups.

(d) Once the nucleophile adds to an aldehyde, neither H— nor R— can be eliminated since they are strongly basic.

Answer. (d)

509. Which of the following reagents can be used to reduce acetaldehyde to ethyl alcohol? (a) 1. LiAlH4 / 2. H30•

(b) 1. NaBH4 / 2. H30+

(c) H2|Pt

(d) All of these

Answer. (d)

510. What is the hybridization of the carbonyl oxygen in carboxylic acids?

- (a) SP
- *[b]* sp2
- (c] sp3
- [d]s-sp2

# Answer. (b)

511. What is the hybridization and geometry of the carbonyl carbon in carboxylic acids and their derivatives? [a] sp3, tetrahedral

Teekay	Page 40	Educational Consult
<i>(b) sp2,</i> trigonal planar		Answer. (o)
(c) sp2, tetrahedral		519. Which com pound has the highest boiling
(d) sp3, trigonal planar		point?
Answer. (b)		(a ) CH3CH3
512. Organic compound (A), C4H8O, does r	ot	<i>(b)</i> CH30CH3
react with Sodium or PCl5.		(c) CH3CH20H
However, it reduces an alkaline solution of a		(Ы СНЗСООН
copper( ll) salt on heating.		Answer. (d)
Compound (A) could be		520. When carboxylic acids and dicarboxylic
(a) an aldehyde		acids have similar molecular
<i>{b ] a primary alcohol</i>		Weights, how do their melting points compare?
(c) a ketone		[a] Carboxylic acids have greater melting points.
(d) a secondary alcohol		(b) Dicarboxylic acids have greater melting
Answer. (a)		points.
513- Check the incorrect statement :		(c) Both acids have similar melting points.
(a) Acetic acid is present in sour milk		[d] No consistent trend exists.
<i>(b)</i> Formic acid is present in insect bites		Answer. {b)
(c) Tartaric acid is present in grapes		521. The greater acidity of carboxylic acids
(d) Citric acid is a tricarboxylic acid		compared to alcohols arises
Answer. (a)		primarily from :
514. In the conversion of wine to vinegar		(a) the electron-donating effect of the hydroxyl
(a) Ethanol is oxidized to acetic acid		group
<i>(b)</i> Ethanol is reduced to acetic acid		(b) the electron-withdrawing <i>effect</i> of the
(c) Methanol is oxidized to acetic acid		carbonyl oxygen
(d) Methanol is reduced to acetic		(c) the acidic of n-hydrogens of carboxylic acids
acid <b>Answer</b> . { <i>a</i> )		(d) the resonance stability associated with the
515- Vinegar is a		carboxylate ion
(a) 5% solution of acetic arid in water		Answer. (d)
(b) 25% solution of acetic acid in water		622. Which of the following statements is false
(c) 50% solution of acetic acid in water		about the acid-strength of acetic acid?
(d) 40% solution of formic acid in water		(a) Acetic acid is a stronger acid than
Answer. [a]		monocnioroacetic acid,
516. In succinic acid, HOUL(LH2)nLUUH, n	1S	(b) Acetic acid is a stronger acid than propionic
equal to		
		(c) Acetic acid is a weaker acid than
		trichloroacetic acid.
		( <i>u</i> ) Acetic aciu is a weaker aciu than for fine aciu.
Answer (b)		522 Arrange the following compounds in order
517 In succric acid HOOCl(CH2)nCOOH n		of decreasing acidity
is equal to		F
(a) 1		>tooo CHI—CH—cOOH CHI CH—COOH
$\begin{pmatrix} a \\ b \end{pmatrix}$		
$\binom{0}{2}$		(a) $(1) > (2) \gg (3)$
(d) 4		(b) $(3) > (2) \gg (1)$
Answer (d)		(c) $(3) > (1) \bullet (2)$
518. The compound in which hydrogen bon	ding	(d) (2) • (l) > (3)
is not possible is	8	Answer. (b)
(a) CH3OCH3		524. Which of the following is the strongest
(b) H20		
(с) СНЗСН2ОН		(a) Formic acid
(d) CH3C00 H		(b) I FICHIOFOACETIC ACIO
		(כ) אכפדוכ מכום
Impacting Lives		

Teekay	Page 41	Educational Consult
(d) Trifiuoroacetic acid		Dr
Answer. (d)	(, ) Câ	I,- tl -COtOll
525. Which of the following is the str	ongest	F
acid?	.:t) C	H LH L'MH
(a) Butanoic acid	,,,,,,,	
(b) 2-Chlorobutanoic acid	Ansv	ver. (d)
(c) 3-Chlorobutanoic acid	532.	Which of the following will give acetic
(d) 4-Chlorobutanoic acid	acido	n acid-hydrolysis?
Answer. (b)	[a] E	thyl acetate
526. Which of the following is the s	trongest acid (b) A	cetone
in aqueous solution?	(c) M	ethyl propionate
(a) CH3C0OH	(d) L	actic acid
(b) CICH2COOH	Ans	wer (a)
ČC) CH3CH2COOH	533.	Propane nitrile undergoes acid-hydrolysis
(d) CI2CHC00H	to giv	re
Answer. (d)	(a) F	ormic acid
527. Which is the strongest acid?	<i>[b]</i> P	opionic acid
(a) CH3COOH	(c) A	cetic acid
(b) CI2 <b>CHCOOH</b>	(d) B	utyric acid
c) CICH2C0OH	Ansv	ver. {b)
(d) C13CCO0H	534.	The characteristic reaction of carboxylic
Answer. (d)	acids	is:
528. Which of the following compo	unds is most (a) el	ectrophilic addition
acidic?	<i>[b)</i> el	ectrophilic .substitution
(a) CH3C0OH	(c) n	icleophilic addition
<i>[b]</i> C1CH2C00H	(d) n	cleophilic substitution
(c) CH3CH 2C00I4	Ansv	<b>ver.</b> (d)
(d) FCH2COOH	535.	Which of the following compounds will
Answer. (d)	react	with Tollens' reagent to give metallic
529. Which of the following compo	unds is the silve	r?
strongest acid?	(a) F	ormic acid
(a) CH3C0OH	(b) E	thyl alcohol
<i>{b)</i> CH3CH2COOH	(c) A	rctic arid
(c) CF3CO0H	( <i>d</i> ) A	cetone
(d) CH3CH2CH2CO0H	Ansv	ver. (a)
Answer. [c]	536	Vhich of the following compounds entreatment
530. Which of the following compo	unds is least	NaHCO3 will liberate CO2?
acidic?	(a) A	cetic acid
(a) CH3CH2COOH		tnylamine
(b) BrCH2CH2CO0H	(C) A	cetone
Rr		
<sub>(c) CH<sub>3</sub>-CH-rO(H</sub>	Allsv	Ver. (d)
r r	557.	which of the following reagents will
(1) CH <sub>3</sub> -CH-COOH	CONV	ida?
Answer (a)		
531 Which of the following compo	inds is most (h) L	(1/7nCl2
acidic?		1/21/01/2 1/12
(a) CFI3CH2COOH		
(b) <b>BrCH2CH2COO</b> H		
	528	Butvric acid reacts with PC15 to give
	(a) D	anzovi chloride

Теекау	Page 42	Educational Consult
<i>{b)</i> l-Chlorobutane		<b>545.</b> Acetyl chloride undergoes nucleophilic
(c) butyryl chloride		substitution at a faster rate than methyl acetate
(dj l-Chloropropane		because
Answer. (c)		(a) the ester is more sterically hindered than the
<b>539.</b> Acetic acid undergoes re	eduction with	acid chloride
LiAlH4 to give		(b) the acid chloride is more sterically hindered
(a) Ethanol		than the ester
<i>(b)</i> Ethane		(c) the methoxide is a better leaving group than
(c) Ethanal		chloride
(d) Ethyne		(d) chloride is a better leaving group than
Answer. (a)		methoxide
<b>540.</b> Acetic acid reacts with r	nethyl alcohol in	Answer. (d)
the presence of an acid cataly	yst to give	546. Esters and amides are most easily made by
(a) Methyl formate	0	nucleophilic acyl substitution reactions on :
(b) Ethyl formate		(a) acid anhydrides
(c) Methyl acetate		(b) carboxylates
(d) Ethyl acetate		(c) carboxylic acids
Answer. (c)		(d) acid chlorides
<b>541.</b> Calcium acetate on stron	g heating gives	Answer. (d)
(a) Methane + CaCO3	0 00	<b>547.</b> Which of the following conditions will drive
<i>(b)</i> Ethane + CaCO3		the equilibrium of the Fischer esterification
(c) Acetone + $CaCO3$		towards ester formation?
(d) Ethane + CaO		(a) addition of water
Answer. (c)		(b) removal of water as it is formed
<b>542.</b> CH3(CH2)4 is the sodium	salt of hexanoic	(c) addition of alcohol
acid. The water		(d) both $(b)$ and (c)
solubility of this salt 1s :		Answer. (d)
(a) higher than that of hexand	oic acld	548. Which of the following statements
(b) lower than that of hexano	oic acid	describes the first step in the mechanism
(c) completely insoluble		of the aldol condensation?
(d) not predictable		(a) An alpha hydrogen is abstracted by the base
Answer. [a)		to form an enolate anion.
<b>543.</b> Which of the following co	ompounds has the	(b) A nucleophilic base attacks the carbonyl
lowest boiling point?		carbon atom.
(a) l-butanol		(c) The carbonyl oxygen is protonated by the
(b) butanoic acid		base ion.
(c) butanenitrile		(d) The alpha hydrogen is abstracted by an acid
(d) methyl propanoate		to the enolate anion.
Answer. (c)		Answer. (a)
544. Which of the following is	the correct	549. Ammonium acetate on strung heating gives
ranking in decreasing order of	of relative Boiling	(a) Urea
Point of carbonyl containing	compounds?	<i>(b)</i> Formamide
(a) primary amide carboxyli	c acid •• ester -	(c) Uric acid
acyl chloride aldenyde K	etone	(d) Acetamide
$\{b\}$ ester > carboxylic acid >» a	imide - acyl	<b>Answer</b> . (a)
chloride aldehyde - ketone	endia agid > » agtor	<b>550.</b> Which reaction does not yield an ester as
all chlorido amido	oxynt atiu >» estei	one of the products?
(d) carboxylic acid > amido >	sector acul	(a) A calboxylic acid is fieated with an alconol (b) A Crignard reagant is added to a carborrite
chloride aldehude - keton	- colci acyi	correction a carboxyllc
Answer, [a]	~	aciu (c) An acid halide is treated with an alcohol

Teekay	Page43	Educational Consult
(d) An alkyl halide is heated with the salt o	fa	(b) Ethylamine
carboxylic acid		(c) Methyl cyanide
Answer. (b)		[d] None of these
551. Silver acetate reacts with Br2 to form		Answer (c)
methyl bromide, carbon dioxide, and		558. Acetamide is a much weaker base than
AgBr. This is an example of		ammonia. This is because
(a) Wurtz reac:tion		(a) the electron withdrawing effect of the C=O
<i>{b)</i> Etard's reaction		group makes the lone pair of
(c) Hunsdiecker reaction		electrons on nitrogen atom less available for
(d) Perkin reaction		protonation.
Answer. (c)		[b ) the electron withdrawing effect of the C=0
552. Hunsdiecker reaction is used for the		group makes the lone pair of electrons on
preparation of		nitrogen atom more available for protonation.
(a) Alkyl chlorides and bromides		(c) the presence of CH3 group sterically hinders
(b) Alkyl nitrates and nitrites		the protonation of the NH2 group.
(c) Ketenes		[d] an H atom from the nitrogen is less easily lost
(d) Alcohols		from CH3C0NH2 than from
Answer. (a)		NH3.
553. A compound undergoes reduction wit	h	Answer. (a)
LiAlH4. lt also dissolves in aqueous		559. Which of the following statements is <i>False</i>
NaOH from which it can be recovered by		about acetamide?
addition of HCl. The compound is		(a) reacts with Br2/Na0H to form tertiary amine
(a) a carboxylic acid		(b) undergoes acid-hydrolysis to form acetic acid
<i>{b)</i> an ester		[c] undergoes dehydration with P205 to form a
(c) en acid anhydride		nitrile
<i>{d ]</i> an alcohol		(d) is a white crystalline solid, soluble in water
Answer. (a)	ha	Answer. (a)
554. Acta childrades andergo reduction to t	lie	O II
corresponding aldenydes on	C	<b>560.</b> The C—C—0 bond angle $i_n Cl[2-, 2]$
treatment with hydrogen in the presence o	I	is approximately
Pd/BaSU4. This reaction is called :		(a) $60^{e}$
(a) Clemmensen reduction		(b) 109.5°
<i>(b)</i> Rosenmund reduction		(c) 90 <sup>°</sup>
(c) Wolf-Kishner reduction		(d) 120 <sup>e</sup>
Answer. (b)		Answer. (d)
FFF Agetic an hydride is obtained by the		<b>561.</b> Consider a small aliphatic 1° amine1
555. Acetic annyuride is obtained by the		alcohol, ester, carboxylic acid, and
reaction of:		amide, all of similar molecular weight. Which
(a) Acetic acid and water		one is most likely to have the
(c) Acetic acid and diethyl ether		highest boiling point?
(d) A cetic acid and P205		[a] The amide
(d) Acetic acid and 1 205		(b) The alcohol
FEG Acotic aphydride reacts with <b>athanal</b> t	•	(c) The ester
give	5	(d) the carboxylic acid
(a) Methyl acetate		Answer(a) Amides can form very strong H-
(h) Methyl propionates		bonds because of ability of the N to donate
(c) kthv1 acetate		electrons to the carbonyl (more so than OH of a
(d) Ethyl propionate		carboxylic at id) and give the fi considerable
Answer. (c)		negative charge, thereby enhancing the H-bond
<b>557.</b> Acetamide on heating with P205 give	s:	acceptor capability of the molecule. Considering
(a) Methylamine		compounds of similar molecular weight and
Imnacting Lives	I	
impacting hives		

Teekay Page 44	Educational Consult
carbon chain structure, <i>the</i> general trend is	Answer. (d)
(highest to lowest boiling point) : amides,	568. Which of the following reagents will reactwith
carboxylic acids, alcohols, amines,	acetamide to form
aldehydes/ketones/esters, alkynes.	methylamine?
alkenes, and alkanes.	(a) LiAlH4
562 Which of the following acid-derivatives are	(b) $Br^2/NaOH$
most stable?	
(a) Acid chloridae	(d) H2/N;
(a) Estars	
(D) Esters	Answer. {b]
(c) Amides	569. Which of the following compounds does
(d) Anhydrides	not form a salt with dilute HCI?
<b>Answer.</b> (c). Amides are the most stable of the	[a] CH3CH2C0NH2
carboxylic acid derivatives	(b) (CH3CH2)2NH
Because they have the most stable resonance	(c) CH3CH2N H2
forms :	(d)(CH3CH2)3N
0	Answer (a)
	570 Propionamide on heating with a mixture of
R- <sup>1</sup> C ∜ NHS RC <sup>1</sup> NH <sub>2</sub>	bromine and sodium hydroxide gives
	(a) Propana
563. Which of the following will have the highest	(a) Fropalle
boiling point?	(b) Propyralline
(a) Diethyl e1er	(c) Propanol
(b) p-Butane	(d) Ethylamine
(c) Acotamido	Answer. (d)
(d) n Dronylamina	571. What happens when urea is treated with
	hydrazine?
Answer. (c)	(a) Biuret is formed
564. Acetamide can be converted into	(b)Semi carbazide is formed
methylamine by	(c) Carbon dioxide and nitrogen is formed
(a] P205	(d) Acetamide is formed
(b) Br2/NaOH	Answer [h]
(c) PCI5	572 Carbonyl chloride reacts with ammonia to
(d) LiAlH4	give
Answer. (b)	(a) Uroa
565. Acetamide on reduction with LiAlH4 gives	(a) Olea
(a) Acetaldehvde	(D) Acetone
(h) £thy)amine	(c) Acetamide
(c) Acetone	(d) Chloroform
(d) Mothylamino	Answer. (a)
(u) Methylamme	573. Which of the following is not a constituent
Allswei. [0]	of normal urine?
566. Amides undergo hydrolysis under acidic	(a) Alhumin
conditions to give	(b) Urea
(a) Carboxylic acid and amine	(c) Sodium chloride
<i>(b)</i> Carboxylic acid and ammonium ion	(d) Uric acid
(c) Carboxylate ion and ammonium ion	Answer (a)
(d) Carboxylate ion and amine	574. Esters can be formed by nucleonbilic
Answer. (b)	substitution reactions of .
567. By which of the following methods can	substitution reactions of :
CH3CON H2 be converted to CH3CA?	(a) carboxylic acids and alcohols ; acid-catalyzed
(a) $Oxidize with Na2Cr2O7/H+$	(b) acid chlorides and alcohols
(b) Heat with concentrated sulfuric acid	(c) both of the above
$(a)$ Heat to $200^{\circ}$	(d) neither of the above
	Answer. (c)
(u) warm with P206	575. The conversion is called :
Impacting Lives	

Teekay Page 4	5 Educational Consult
(a) Oxidation	(c) Ethyl alcohol
(b) Dehydration	(d) Diethyl ether
(c) Reduction	Answer. (a)
(d) Hydrolysis	583. When ethyl acetoacetate is subjected to
Answer. (c)	ketonic hydrolysis, the ketone
<b>576-</b> The following reaction is <i>ca1led</i> :	obtained is
0 0	(a) Dimethyl ketone
cccv-c-no-b can-c-of cncwn	(h) Methyl ethyl ketone
(a) 5aponification	(c) Diethyl ketone
(b) Condensation	(d) Methyl n-propyl ketone
(c) Elimination	Answer (c)
(d) Esterification	594 Ethyl acotoacotate reacts with
Answer. (a)	phonul hydroging to give
<b>577.</b> Hydrolysis of an ester can be accomplished	(a) Antinumina
by:	(a) AntipyThe
(a) base-promoted hydrolysis	(b) Aspirin
(b)acid-catalward hydrolysis	(c) 4-Metnyl uracli
(c) both of the above	
(d) neither of the above	Answer. (a)
Answor (c)	585. Xeto-enol tautomerism is shown by
E70 Pasis hydrolysis of actors is called	(a) Bentaldehyde
(a) A contribution	(b) Acetone
(d) Acidification	(c) Benzophenone
(b) Fotorification	(d) Acetic acid
(d) Sananification	Answer. (b)
(u) Sapolinication	586. Fatty acids are
Answer. (a)	(a) Unsaturated dicarboxylic acids
579. Base-catalyzed condensation of two ester	(b) Long-chain alkanoic acids
molecules to form an alconol and	(c) Aromatic carboxylic acids
B-keto ester is called	(d) Aromatic dicarboxylic acids
(a) Claisen condensation	Answer. [b]
(b)Corey-House reaction	587. Sodium or potassium salts of fatty acids are
(C)Aldol condensation	called
(d)Transesterification	(a) Proteins
Answer. (a)	[b] Terpenes
580. Ethyl acetoacetate undergoes	(c) Carbohydrates
acid-hydrolysis with dilute HCI to form	(d) Soaps
(a) Acetoacetic acid	Answer (d)
(b)Succinic acid	588. Soap is
(c) Acetic acid	(a) a mixture of salts of fatty acids
(d) Adipic acid	(b) a salt of glycerol
Answer. (a)	(c) a mixture of ethers
<b>581.</b> The ethyl derivative of acetoacetic ester on	[d] a mixture of aromatic ethers
basic-hydrolysis gives	Answer. (a)
(a] Acetic acid	589. A wax is
(b) Acetic acid and propionic acid	(a) a nonpolar solid
(c] Propionic acid	(b) a long-chain alcohol
(d) Acetic acid and n-butyric acid	(c) a triacylglycerol
Answer. (a)	(d)none of these
582. Ethyl acetate on heating with sodium	Answer. (a)
ethoxide gives	590 Fats and oils are
(a) Ethyl acetoacetate	(a) monoesters of alveeral
(b) Sodium acetate	(b) diatars of glycerol

Teekay	Page 46	Educational Consult
(c) triesters of glycerol		(d) is used in the production of lactic acid
(d) diesters of glycol		Answer. {b}
<b>Answer.</b> (c). Fats are solid triesters of gly	cerol.	<b>598.</b> Which of the following compounds will not
Oils are liquid triesters of		be classified as lipids?
glycerol.		(a) Fats
591. Fats differ from waxes in that fats ha	ve :	(b) Waxes
(a) More unsaturation		(c) Soans
(h) Higher melting points		(d) Oils
(c) A glycerol backhone		Answer {h}
(d) Longer fatty acids		<b>599</b> Synthetic detergents can be represented by
Answer (c)		the following general formula
592 Oleic acid is a fatty acid containing		[a] RONa
(2) 12 carbons		$(b) \text{ POSO3N}_2$
(a) 12 carbons $(b)$ 14 carbons		$(b) \operatorname{PCOON}_2$
$\{U\}$ 14 cal bolls		(d) PC00 H
(d)10 carbons		
(u)18 cal bolls		<b>Answei</b> [U] <b>(00</b> The degree of upperturbing of a fat can be
Allswei. $[u]$	10	determined by means of its
593. Both stearic acid and linoleic acid na	Ive 18	(a) iodha numbar
carbons. Linoleic acid is	ጥ .	(a) fourne number
Unsaturated, while stearle acid is saturated	. The	(b) Octaile number
melting point of stearic acid :		(c) Saponification number
(a) is higher than linoleic acid		(a) Melting point
<i>{b]</i> is lower than linoleic acid		Answer. (a)
(c) is same as linoleic acid		<b>601.</b> Which of the following statements is not
(d) cannot predict, insufficient information	on	true about fatty acids?
Answer. {a)		(a) Fatty acids are carboxylic acids with long
594. Liquid offs can be converted to solid	fats by	hydrocarbon side chains
(a) Hydrogenation		(b) The double bonds in unsaturated fatty acids
(b) Saponification		are always conjugated.
(c) Hydrolysis		(c) Most naturally occurring fatty acids contain
(d) Oxidation of double bonds		even numbers of carbons and are
Answer. [a]		unbranched.
595. Partial hydrogenation of vegetable of	ils in	(d) Fatty acids can be saturated or unsaturated.
the presence of Ni catalyst at		Answer. (b)
200°C gives		<b>602.</b> Which of the following best explains why
(a) Vanaspati ghee		the melting points of saturated fats
(b) Margarine		increase with increasing molecular weight?
(c) Both of these		(a) decreased hydrogen bonding
(d) None of these		(b) increased hydrogen bonding
Answer. (a)		(c) dccreased intcrmolocular van der Waal's
596. Alkaline hydrolysis of oils (or fats) is	s called	interactions
		[d] increased intermolecular van der Waal's
(a) 5aponification		interactions
<i>(b)</i> fermentation		Answer. (d)
(c) Diazotization		<b>603.</b> Which of the following terms best describes
(d) Rancidification		the compound below?
Answer. (a)		СН3(СН2)12С02Н
<b>597.</b> Saponification of a fat		(a) a fatty acid
(a) always results in the formation of inso	luble	(b) an oil
soaps		(c) a wax
<i>(b)</i> produces glycerol and soan		(d) a soap
(c) is used in the production of detergents		Answer. (a)
(, ) production of detergende		
Impacting Lives		

Teekay Page 4	7	Educational Consult
<b>604.</b> Which of <i>the</i> following terms best describes		(c) Phosphatidic acid is a phospholipid.
the compound below?		(d) J'he C-2 carbon of glycerol in
CH3(CH2)7CH=CH(CH2)7C02H		phosphoacylglycerols has the R configuration.
(a) an unsaturated fatty acid		Answer (a)
(b) a triglyceride		610. The compound inethylamine, H3C—NH2,
(C) a synthetic detergent		contains a C—N bond. 1 n this bond,
(d)a miccl⊮		which of the following best describes the charge
Answer. (a)		on the nitrogen atom?
<b>605.</b> Which of the following statements best		(a) slightly positive
describes the structure of waxes?		(b) uncharged
(a] long-chain unsaturated carboxylic acids		[c] slightly negative
<i>(b)</i> long-chain saturated carboxylic acidstc)		(d) —1
long-chain esters		Answer (c)
(d) short-chain esters		611. Triethylamine [(CH3CH2)3 N] is a molecule
Answer. (c)		in which the nitrogen atom is
606. Which of the following terms best describes		hybridized and the CNC bond angle is
the compound below?		
CH3(CH2)26C02CH2(CH2)32CH3		(a) <i>sp2</i> , ' >109.5°
(a) a fat		(b) $sp_{2} < 109.5^{\circ}$
(b) a wax		[c] sp3; >109.5°
(c) a terpene		(d) sp3; <109.5°
(d) an unsaturated triglyceride		Answer. (d)
Answer. (b)		<b>612.</b> The N-H bond in the ammonium ion, N H4+,
607. Which of the following statements is not		is formed by the overlap of what
true about triacylglycerols?		two orbitals?
(a) When solids and semisolids at room		(a) sp3—rp3
temperature, they are called fats.		(b) sp3-sp2
(b) When liqu Ids at room temperature, they are		(c) $sp2 - sp2$
called oils.		[d] sp3-s
(c) When hydrolyzed, they produce glycerol and		Answer. (d)
carboxylate salts.		613. What is the hybridization of the nitrogen
(d) Triacylglycerols with low melting points are		atom in CH3NH2?
composed of saturated fatcy		(a) <i>sp</i>
acids, causing them to be liquids at room		[b j sp2
temperature.		(c) sp 3
Answer. (d)		(d) s-sp3
<b>608.</b> Triglycerides which are solids or		Answer. (c)
semisolids at room temperature are called		614. What is the hybridization of the carbon
		atom in CH3NH2?
(a) oil		(a) 3sp3
(b) fat		(b) sp
(c) steroid		<i>(c)</i> sp2
(d) cholesterol		(d) sp3
Answer. (b)		Answer. (d)
<b>609.</b> Which of the following statements is not		615. What type of intermolecular interactions
true about phospholipids?		does (CH3CH2)2 NH undergo?
(a) They are similar to triacylglycerols except		(a) induced dipole-induced dipole
that the middle 0 H group of		(b) dipole-dipole
glycerol reacts with a phosphate rather than		(c) hydrogen bonding
with a fatty acid.		(d) all of these
<i>{b)</i> They constitute a major component of cell		Answer. (d)
membranes.		
Impacting Lives		
r · · · · · · · · · · · · · · · · · · ·		

Teekay	Page 48	Educational Consult
616. Assuming roughly equivalent molecula	r	<b>622.</b> Which of the following is least soluble in
weights, which of the following		water?
would have the highest boiling point?		(a) Methylamine
(a) a tertiary amine		(b) Trimethylamine
(b) a quaternary ammonium salt		(c) Diinethylamine
(c) an alcohol		(d) Aniline
(d) an ether		Answer.(d)
Answer. (b)		<b>623.</b> Which of the following is most soluble in
617. Which of the following statements is fo	ilse	water?
about primary amines?		(a) CH3CH2CH3
(a) They can be prepared by reduction of nit	riles	Ι
with LiAlH4.		(b) $CH3 - CH - CH3$
(b) They do not form salts with acids.		(c) C113C1[2C[12C]]
(c) They react with ice-cold nitrous acid to fe	orm	
nitrogen gas.		Answer (d)
(d) They are basic and soluble in water.		<b>624</b> Which aming is NOT coluble in water?
Answer. (b)		(a) Methylamine
618. Which of <i>the</i> following statements is <i>fa</i>	Ise	(h) Dimothylamino
about secondary amines?		(b) Dimethylamine
(a) They react with chloroform and NaOH to		(d) All aroundton coluble
form isocvanides.		(d) All are water-soluble
<i>(b)</i> They can be prepared by reduction of		Therefore all low molecular
isocyanides.		weight amines are soluble in water
(c) They react with acid chlorides to give		<b>625</b> Consider a $1^{\circ}$ $2^{\circ}$ and $2^{\circ}$ amino all of
substituted amides.		oquivalent molecular weight Which
(d) They react with ice-cold nitrous acid to g	ive	aming is most likely to have the lowest heiling
nitrosamine (yellow oil).		noint?
Answer. [a]		(a) l <sup>o</sup> amine
619. The hybridization of nitrogen in an am	ine is	(h) 2° amine
(a) <i>sp</i>		(c) $3^{\circ}$ amine
(b) sp3		(d) Not enough information to determine
(c) sp2		Answer, (c). The 3° amines do not rate
(d) sp4		hydrogen bonds to one another. The 1°
Answer. (b)		and 2° amines can form hydrogen bonds to
<b>620.</b> Which of the following compounds is a		themselves and therefore have higher
secondary amine?		boiling point than 3° amines.
NH <sub>2</sub>		<b>626.</b> Which of the following will have the highest
		boiling point?
$_{(a)}$ CH <sub>3</sub> -CH-CH <sub>3</sub>		(a) Methylamine
(b) H2NCH2CH2NHZ		(b) Diethylamine
(c) CH 3CH2 N HCP-13		(c) Ethylamine
(d) (CH3)3N		(d) Triethylamine
Answer. (c)		Answer. (d)
<b>621.</b> A sample of pure amine molecules is fo	und	<b>627.</b> Which compound has the highest boiling
to possess no intermolecular Hbonding.		point?
This sample is most likely :		1. CH3CH2CH3 2. CH3CH20H 3. CH3CH2NH2
(a) 1° amine		(a) I
<i>{b)</i> 2° amine		(b) 7
(c) 3° amine		(c) 3
(d) all of these		(d) 2 and 3 have similar boiling points
Answer. (c)		

Teekay Pa	ge 49	Educational Consult
Answer. (b). Amines are fairly polar, but they	r	635. The major product of the following reaction
boil at temperatures lower than		is :
those of alcohols of similar chain length and		CH3CH2Cl+(CH3)2NH□-→?
structure. Amines do have higher		(a) 1° amine
boiling points than alkanes because they posses	SS	$(h) 2^{\circ}$ amine
H-bond donors and acceptors.		(c) 3° amines
628. Which of the following has the highest		(d) amide
boiling point?		Answer. (c)
(a) UH3UH2UH2UH3		636. Aliphatic primary aminns react with cold
$\begin{cases} D \end{bmatrix} \bigcup_{i=1}^{n} \bigcup_{j=1}^{n} \bigcup_{j=1}^{n} \bigcup_{j=1}^{n} \bigcup_{i=1}^{n} \bigcup_{j=1}^{n} \bigcup_{j=1}^{n} \bigcup_{j=1}^{n} \bigcup_{j=1}^{n} \bigcup_{j=1}^{n} \bigcup_{i=1}^{n} \bigcup_{j=1}^{n} \bigcup_{j=1}^{n$		nitrous acid to form
		(a) Alcohols
		(b) Diazonium salts
Answer. (C)		(c) Nitriles
629. Ammes are generally classified as		(d) Nitroalkanes
(a) Weak actus (b) Strong acida		Answer. (a)
(D) Strong actus		637. Which type of amine produces N2 when
(d) Strong bases		treated with H0N02
Answor (c)		[a] Primary
630 Acotamido roacts with Br2/Na0H th giv	0	(b) Secondary
(a) Mothylamino	C	(c) 'Tertiary
(a) Methylannie (b) Uree		(d) Quaternary
(c) Ethylamine		Answer. (a)
(d) Acetyl bromide		638 Methylamine reacts with nitrous acid to
Answer (a)		form
631 Acetonitrile undergoes reduction with		(a) CH3CH3
LiAl H4 to form :		<i>[b]</i> CH30H
(a) Methylamine		[c] CH3N02
(b) Dimethylamine		(d) CH3CH20H
(c) Ethylamine		Answer. {b)
(d) Trimethylamine		639. Which of the following compounds reach
Answer. (c)		with nitrous acid to form an
632. Reduction of an amine will give an		alcohol?
(a) Acid		[a] CH?CH2NH2
<i>{b</i> } Amide		[b] (CH3CH2)2NH
(c) Amine		[c] (CH3CH2)3N
(d) Alcohol		
Answer. (c)		$( ) = NH_2$
633. Which of the following is most basic?		(a)
(a) Ammonia		Answer (b]
(b) Methylamine		640. A nitrile can be made by dehydrating an
(c) Dimethylamine		amide. However, for this reaction to occur, the
(d) Trimethylamine		amide must be :
Answer. (c)		(a) primary
634. Alkyl halides react with ammonia in the		[b) secondary
presence of base to form :		(c) tertiary
(a) primary amines		[d] N-methylated
<i>(b)</i> nitrated alkyl halides		Answer. (a)
(c) amides		641. Which of the following amines react with
(d)quaternary ammonium salts		nitrous acid to form nitrosoamine?
Answer. [a]		(a)Ethylamine
		<i>[U]</i> Theorytamine

Teekay	Page 50	Educational Consult
(c) Diethylamine		(a} ethyl group
(d) lsopropylamine		(b) hydrogen
Answer. (c)		(c] isobutyl group
642. Which of the following does not react	with	(d) isopropyl group
acyl chlorides to form amides		Answer. (d)
(a) ammonia		650. Which of the following <i>describes</i> the side
(b) 1º amine		chain of Leucine?
(c) 2º amine		(a) ethyl group
(d) 3º amine		(b) isopropyl group
Answer. (d)		(c) sec-butyl group
643. Hinsberg's reageni is		[d] isobutyl group
(o) $Pd + BaSO4$		Answer. (d)
tb) p-Toluenesulfonic acid		651. Which of the following amino acids has an
(c) N H2N H2 $*$ KOH		aliphatic R group?
(d) Benzenesulfonic acid		(a) Serine
Answer. (d)		(b) Cysteine
644. Which of the following compounds wi	i <b>11</b>	(c) Asparagine
dissolve in aqueous NaOH alter		(d) Leucine
undergoieg reaction with Hinsberg reagent	-?	Answer. (d)
(a) CH3CH2NH2		652. Which of the following amino acids has an
<i>{b</i> ) (CH3CH2)2NH		aromatic R group?
(c) (CH3 CH2)3N		(a) Serine
(d) (CH3)3N		(h) Cysteine
Answer. (A)		[c] Asparagine
<b>645</b> . Which of <i>the</i> following compounds do	nes	(d) Tyrosine
not react with acetyl chloride?	,65	Answer. (d)
(a) Methylamlne		<b>653.</b> Which of the following amino acids has a
(b) Dimcthylamlnc		sulfur in the R group?
(c) Aniline		(a) Serine
(d) Trimethylamine		(h) Cysteine
Answer (d)		(c) Asnaragine
646 A primary amine can be identified by	using	(d) Tyrosine
(a) H()	using	Answer. (d)
(b) CHCl3		<b>654</b> Which of the following amino acids has a
(c) NaOH		heterocyclic R groun7
(d) CHCl3 + KOH		(a)Glycine
Answer (d)		(h) Threonine
647 Carbylamine test is given by		(c) Proline
(a) Primary amines		(d) Aspartic acid
(b) secondary amines		Answer. (c)
(c) Tertiary amines		<b>655</b> . The alpha-carbon of all the amino acids
(d)None of these		is a chirality center excent for
Answer (a)		is a chirality conter except for
648 Which of the following describes the s	side	(a) Glycine
chain for Alanine?	nuc	(h) Threonine
(a) hydrogen		(c) Proline
(b) ethyl groun		[d] Aspartic acid
(c) methyl group		Answer (a)
(d) isopropyl group		<b>656.</b> Which of the following is correct for
Answer. (c)		anamino acid solution when it is in the
649. Which of the following describes the	side	acidic form?
chain for Valine?		(a) nH > nka
	ļ	

Facebook/I nstagram/PIayStore/Youtube: Baflk Of MCQs

Teekay	Page 51	Educational Consult
<i>{b)</i> pH < pxa		(c) location of the disulfide bridges in the
(c) zwitterionic		peptide chain
(d) $pH = 7$		(d) conformation of the protein backbone
Answer. (b)		Answer (b)
657. When a disulfide linkage is formed,	the	663. Which of the following protein structures
compound containing this new		'does denaturation" destroy?
linkage has been		(a) primary and secondary structures
(a] hydrolyzed		(b) secondary and tertiary structures
<i>(b)</i> dehydrated		(c) tertiary and quaternary structures
(c) electrolyzed		(d) secondary, tertiary, and quaternary
(d) oxidized		structures
Answer. (d)		Answer. (d)
658. Which of the following structures ca	n	664. Which of the following are capable of
polypeptides have?		denaturing proteins.'
(a) primary structure		(a) organic solvents
(b) secondary structure		(b) extreme pH
(c) tertiary structure		(c) heat
(d) all of the these		(d) all of the these
Answer. (d)		Answer. (d)
<b>659.</b> Which of the following is the first st	ep in	665. The monomeric units that make up
the determination of the primary		peptides and protein polyrners are
structure of proteins?		
(a) determining the number and kind of a	mino 🔪	(a) nucleic acids
acids in the peptide		(b) amino acids
tb) reducing the disulfide bridges in the p	rotein	[c) oligosaccharides
(c) protecting the N-terminal of the pepti	de	(d) amylopectins
(d) protecti+tg the <i>C</i> -terminal of the peptic	le	Answer. (b)
Answer. (b)		666 A zwitterion is
<b>660.</b> What are enzymes?		(a) an ion that is positively charged in solution
(a) saccharides that catalyze chemical rea	ictions	(b) an ion that is negatively charged in solution
(b) unsaturated fats that catalyze chemical		(c) a compound that can ionize both as a base
reactions		and an acid.
(c) DNA molecules that catalyte chemical		(d) a carbonydrate with an electrical charge.
(d) protoing that gatalyze shominal reaction	<b>.</b>	Answer (C)
(u) proteins that catalyze themical reaction	JIIS	667. A zwitterion has which of the following
<b>661</b> Which of the following may character	rizo	properties
the "secondary strijsture" of		(u) no net charge $(b)$ a high molting point
nrotoins?		$\{b\}$ a high metung point $(c)$ soluble in water
(a) conformation of the protoin backhone		(d) ail of these
(h) alpha-Heliv		
(r) narallel B-nleated sheet		AllSwell. (d) 669 An aqueous solution of giveing is neutral
d) all of the above		bocause of the formation of
Answer (d)		(a) Carbanian
662 Which of the following is the "quater	narv	(a) Carbanion (b) Zwitterion
structure" of proteins concerned	nai y	(c) Carbonium ions
with?		(d) Free radicals
(s) sequence of amino acids in the pentic	le chain	Answer {h}
(b) description of the way the pentide ch	ains are	669. Which one of the following compounds
arranged with respect to each	•	f'orm zwitterions?
other		(a) carbonyl compounds
		(b) amino acids
	I	

Teekay

(c) phenols

(d) beterocyclic compounds

# Answer. (b)

670. The pH at which the amino acid shows no tendency to migrate when placed in an electric field is known as its :

(a) lsoclectric point

- (b) Dipole moment
- (c) Iodine number
- (d) Wavelength

#### **Answer**. (a)

671. The isoelectric point of a protein is

(a) the pH at which the protein molecule has no charges on its surface.

(b) the pH at which a protein in solution has an equal number of positive and

negative charges.

(c) the electric charge under isothermal conditions.

(d) None of these.

## Answer. (b)

672. Which of the following reactions is suitable for the preparation of a-amino acids?

- (a) Schmidt reaction
- *(b)* Hofmann's degradation of amides
- (c) Strecker'S synthesis
- (b) Reduction of nitro compounds

Answer. (c)

**673.** Glycine is

- (a) NH2CH2CO0H
- (b) NH2CH2CH2CH2CH2NH2
- (c) NO2CH2CH2C00H

(d) BrCH2COOH

## Answer. (a)

**674.** Glycine reacts with nitrous acid tp form : (a) Glycollic acid (b) Diketopiperazine

- (c) Methylalamine
- (d) Ethyl alcohol
- (d) Ethyl alcohol

Answer. (a)

**675.** When glycine is heated, it forms (a) Diketopiperazine

*(b)* Acrylic acid

(c) Butyric acid

(d) bu tyrlactam

# Answer. (a)

676- Glycine is a unique amino acid because it
(a) has no chira) carhon
(b) has a sulfur containing *R* group
(c) cannot form a peptide bond
(d) is an essential amino acid

Answer. (o) **677.** Which of the following organic ions results

when glycine is treated with concentrated HCP? (a)  $\stackrel{+}{NH_3CH_2COOH}$ (b)  $\stackrel{+}{NH_2CH_2COO}$ (c)  $\stackrel{+}{NH}$ , CH,  $\stackrel{-}{OO}$ (d) HOCH<sub>2</sub>COO

Answer. (a) 678. Proteins are (a) polyamides (b) Polymers of ethylene (c] n-Aminocarboxylic acids (d) Polymers of propylene Answer. (a) **679.** The five elements present in most naturally occurring proteins are : (a) C, H, O, P, and S (b) N, C, H, O, and I (c) N, s, C, H, and 0 (d)C.H,O,S,andlAnswer. (c) **680.** The nitrogen content of proteins can be quantitatively determined by (a) Carius method *{b )* Kjeldahl's method (c) Victor Meyer's method (d] Rast method Answer. (b) 681. Which of the following is the main structural feature of proteins? (a) Peptide linkage (b) Ester linkage [c] Ether linkage (d)  $\alpha$ ,  $\beta$ -Linkage Answer. (a)

682. The linear arrangement of amino acid units in proteins is called :

- [*a*] primary structure
- *(b)* secondary structure
- (c) tertiary structure
- (d) quaternary structure
- Answer. (a)
- 683. The primary structure of a protein refers to

(a) whether the protein is fibrous or globular(b) the amino acid sequence in the polypeptide chain

(c) the orientation of the amino acid side chains in *space* 

Teekay	Page 53	Educational Consult
(d) the presence or absence of an a-helix.		(c) Heavy metal salts
Answer. (b)		(d) All of these
684. The $\alpha$ -Helix is a common form of		Answer. (d)
(a)primary structure		692. Ninhydrin test is given by
<i>{b</i> ) Tertiary structure		(a) Carbohydrates
(c) Secondary structure		(b) Proteins
(d) None of these		(c) Alkanes
Answer. (c)		(d) Alkenes
<b>685.</b> The $\alpha$ -Helix is held in a coiled conform	mation	Answer. (b)
partially because of:		693. Which of the following tests is not used for
(a) Optical activity		testing proteins?
<i>(b)</i> Hydrogen bonding		[a] Ninhydrin test
(c) Resonance		(b) Biuret test
(d) Delocalization		(c) Xanthoproteic test
Answer. {b)		(d) Toilets' test
686. The double helical structure of DNA i	s held	Answer (d)
together by		694. A protein solution on warming with
(a) sulfur-sulfur linkages		concentrated HNO3 may turn yellow.
(b) peptide bonding		This test is called :
(c) hydrogen bonding		(a) Xanthoproteic test
(d) glycosidic bonds		(b) Ninhydrin test
Answer. (c)		(c) Biuret test
687. Upon hydrolysis, proteins give		(d) Million's test
(a] Amino acids		Answer. (A)
(b) Hydroxy acids		695. A compound gives a positive Tollens' test
(c) Fatty acids		but negative Ninhydrin test It is
(d) Alcohol		(a) a protein
Answer. [a]		(6) an amino acld
688. Complete hydrolysis of proteins proc	luces :	(c) a monosaccharide
(a) Ammonia and carbon dioxide		(d) pyridine
<i>(b)</i> Urea and uric acid		Answer. (c)
(C)A mixture of amino acids		<b>696.</b> Cycloalkanes have the same molecular
(d)Glycogen and a fatty acid		formula as :
Answer. (c)		(a) Alkanes
689. Digestion of proteins involves		(b) Alkenes
(a)changes in secondary structure only		(c) Alkynes
(b) cleavage of peptide linkages	_	(d) Cycloalkenes
(c) removal of all carbonyl groups in the fo	orm of	Answer (b)
	6 11 110	<b>697.</b> Which of the following compounds will give
(d) removal of all NH2 groups in the form	of NH3	cyclopropane on treatment with
Answer. (b)		sodium in dry ether?
<b>690.</b> Irreversible precipitation of proteins		[a] 1,3-Dibromopropane
caused by heating is called :		(b) 1,1-Dibromopropane
(a) Polymerisation		(c) 1,2-Dibromopropane
(b) Denaturation		(d) 2,2-Dibromopropane
(C) Electrophoresis		Answer (a)
(d) inversion		698. Which of the following cycloalkanes is most
Allswer. (D) 601 Procipitation or coordilation of practo	nc	reactive?
be sourced by	115	(a) Cyclopropane
may be caused by		(b) Cyclonexone
(a) field (b) Changes in pU		(c) Cyclobutane (d) Cyclobartana
		(u) cycloneptane
Impacting Lives		

Teekay	Page 54	Educational Consult
Answer. (a)		(d) 120°
699. Which of the following compounds wi	11	Answer. (a)
react most readily with concentrated		<b>706.</b> Which of the cycloalkanes is not expected
sulfuric acid?		to have ring strain?
(a) Ethane		(a) Cyclopropane
(b) Cyclobexaoe		(b) Cyclobutane
(c) Propane		(c)Cyc oheptane
(d) Cyclohexene		[d] None of these
Answer. (d)		Answer. (c)
700. A compound of formula C6H12 does r	not	<b>707.</b> The bond angle between carbon atoms in
react with concentrated sulfuric acid.		cyclohexane is
The compound could be		(a) 60°
(a) Alkane		(b)90°
<i>(b)</i> Cvcloalkane		(c) 109.6°
(c) Alkene		(d) 120°
(d)Cvcloalkene		Answer (c)
Answer. [b]		<b>708</b> Which of the following statements is false
701. Which of the following molecules will		about cyclohexane?
decolorize bromine in carbon		(a) It 1s a saturated cyclic hydrocarbon
tetrachloride most readily?		(b) All C—C—C bond angles are 109°28'
(a) 1 2-Dimethylcycionronane		(c) It is veiv unstable strained compound
(a) 1,2 Dimetary to propring (b)		(d) It can exist in two conformations which are
(c) 1 2-Dimethylcydobutane		designated as the hoat-form and
(d) Cyclohexane		the chair-form
Answer {a}		Answer. (c)
702 Cyclopropane reacts with concentrate	d HBr	<b>709</b> The most stable conformation of
to give		cyclohexane is the :
(a) l-Bromonronane		(a) Haworth form
(b) Bromocyclopropane		(b) Boat form
(c) 2-Bromonronane		(c) Newman form
(d) 1 2-Dibromonronane		(d) Chair form
Answer [a]		Answer (d)
<b>703</b> Cyclobutane reacts with hydrogen in	tho	<b>710</b> What nercentage of cyclohexane molecule
presence of nickel catalyst at 200°C	uic	is estimated to be in the hoat form
to give		at any given time?
(a) Butano		(2) over 90%
(b) l-Butono		(a) $b \in [h]$ (b) hetween 90% and 90%
(c) 2-Butene		(c) approximately $50\%$
(d) None of these		(d) less than 1%
Answer [2]		Answer (d)
704 Cyclohevanol can be converted into		711 Uriedes are
cyclohevene by heating with		(a) Halogen derivatives of urea
(a) $zn(Hg)$ and HCI		(h) A catul darivatives of urea
(a) Zh(Hg) and HCI (b) Concentrated H2504		(c) Alkyl derivatives of urea
(r) SOC12		(d) None of these
(d) H2 and Ni		Answer (b)
Answer (b)		712 Uric acid on ovidation with alkaling KMn04
Allswell (D)	2	format
cyclobeyane is	11	
$(_{2})100^{\circ}28'$		(a) 010a (b) Barbituric acid
(a) 107 20 (b) 60°		(v) bal bitui it attu
		(d) Coffeine

Teekay P	age 55	Educational Consult
Answer. (c)		<b>720.</b> An aqueous solution of glucose behaves as
713- Diethyl malonate reacts with urea in th	e	an aldehyde because
presence of sodium ethoxide to form		
(a) Uric acid		(a) it is hydrolyzed by water to the free aldehyde
(b) Barbituric acid		[b] it is a ketone, but is in equilibrium with the
c) Phenobarbital		aldehyde form
d) Barbital		(c) glucose is actually a cyclicaldehyde
Answer. (b)		(d) its cyclic hemiacetal, the predominant form,
<sup>7</sup> 14. A carbohydrate composed of three to te	en	is in equilibrium with the free aldehyde form
sugar molecules is called a(n) :		Answer. (d)
(a] disaccharide		720. Which of the following statements best
<i>(b)</i> oligosaccharide		describes the meaning of
(c) polysaccharide		"mutarotation"?
(d) monosaccharide		(a) a rapid exchange between the a and § forms
Answer. {b)		of diastereomeric sugars
<b>715.</b> All chiral D-sugars rotate plane-polariz	ed	(b) a rapid exchange between the D and L forms
ight :		(c) a slow exchange between hydrogen and
a) clockwise		deuterated hydrogen
(b) counterclockwise		(d) a slow change in optical legation to reach an
(c) +20.0°		equilibrium value
d) in a direction that cannot be predicted but		Answer. (d)
must be determined experimentally		<b>721.</b> In solution, glucose exists as
Answer. (d)		(a) the open-chain form <b>only</b>
716. How many chirality centers are there in	1 a	(b) the cyclic hemiacetal form only
2-ketohexose?		(c) the cyclic acetal form only
(a] 2		(d) an equilibrium mixture of the open-chain
(b) 3		form and cyclic hemiacetal forms
(c) 4		Answer. (d)
(d) 5		<b>722.</b> Anomers of D-glucopyranose differ in their
Answer. [b]		stereochemistry at
717. How many stereoisomers are possible f	for a	(a) C1 (b) C2 (c) C3 (d) C4
2-ketohexose?		Answer. (a)
(a] 2		<b>723.</b> Which of the following statements best
(b) 4		describes the meaning of a
(c) 8		"glycoside"?
(d) 16		(a) It is the mirror image of a sugar.
Answer. (c)		(b) It is the hemiacetal of a sugar.
718. Which of the following statements best	;	(c) It is the acetal of a sugar.
describes the difference between amylose and		(d) It is the enantiomer of a sugar.
amylopectin?		Answer. (c)
(a) Amylose is a branched polysaccharide wl	nile	<b>724.</b> Which of the following would give a
amylopectin is a		positive Tollens's test?
chain polysaccharide.		(a) s-D-glucopyranose
(b) Amylose is a straight-chain polysacchari	de	[b] methyl alpha-D-glucopyranoside
while amylopectin is a branched polysaccharic	le.	(c) sucrose
(c) Amylose contains o-1,6-glycosidic linkag	e	(d)methyl n-D-ribofuranoside
which amylopectin does not		Answer (a)
contain.		<b>725.</b> Which of the following is true about
d) Amylose is composed of thousands of D-		sucrose?
lucose units while amylopectin is composed	d of	(a) It hydrolyzes to fructose and glucose.
housands of D-galactose units.		[b] It is a reducing sugar.
Answer. [b]		(c) It is a monosaccharide.
	I	
mpacting Lives		

Teekay	Page 56	Educational Consult
(d) lt undergoes mutarotation in water.	(b	)Glucose
Answer. (a)	(c)	Sucrose
726. Which term describes a sugar where	e one of (d)	Galactose
the OH groups has been replaced	Ar	iswer (b)
with a hydrogen?	73	3. Glucose cannot be classified as
(a) amino sugar	(a)	a <i>hex</i> ose
(b) imino sugar	<i>(b)</i>	) an oligosaccharide
(c) dehydroxy sugar	(c)	an aldose
(d) deoxy sugar	(d	a monosaccharide
Answer. (d)	An	swer. {b}
727. Monosaccharides <i>are</i> classified accor	ding 73	4. Which of the following statements is false
to :	ab	out glucose?
(a) the number of carbon atoms in the m	olecule. (a)	it is a reducing sugar.
<i>(h)</i> whether they contain an aldehyde or all	retone <i>(b)</i>	) it is a disaccharide.
group		it has a pyranose form.
(c) their configurational relationship to	[d]	) it is a polyalcohol
glyceraldehyde		swer. {b)
(d) all of the above	73	5. Which of the following statements is false
Answer (d)	ah	out o-D-glucose?
728 Which is a monosaccharide?	(a)	tit has a nyranose ring
(a) Sucrose		it is a hemiacetal
(a) Sucrose		it shows mutarotation
(c) Galactose		it is the purest form of table sugar
(d) Cellulose		swor (d)
Answer (c)	73	6 alpha-D-Clucoso is different from B-D-glucoso
720 Which is a disaccharida?		0. alpha-D-Glucose is unlet ent nom D-D-glucose
(a) Chucoso	(d	) he could they are mirror images of each other
(b) Maltoso		because they are opentiomere
(c) Fructoso		because they are geometrical isometri
(d) Cellulose	(u	because they are geometrical isomers
(u) Cellulose		<b>Swer.</b> (d) $7 \times D$ Characterization is $c(n)$ .
Answer. [D]	/3	/. a-D-Glucopyranose is a(n) :
730. Which of the following is not a	(a)	
monosaccharide?	(D)	) nemiketai
(a) Ribose	(C)	acetal
(b) Fructose	(d	ketal
(c) Sucrose	Ar	iswer. (a)
(d) Glucose	73	8. The number of asymmetric carbon atoms in
Answer. (c)	th	e n-D-glucopyranose molecule
731. The designation D or L before the na	ame of a is:	
monosaccharide	(a)	2
(a) indicates the direction of rotation of		
polarized light.	(b)	3
<i>(b)</i> indicates the length of the carbon chai	n inthe (c)	- 4
carbohydrate.	(d	) S
(c) indicates the position of the OH group	on the Ar	iswer. (d)
carbon next to the primary	73	9. Which of the following statements is false
alcohol group.	ab	out an aldohexose?
(d) indicates the position of the asymmetry	ric (a)	It is a monosaccharide.
carbon atoms in the carbohydrate	(b	) lt contains a potential aldehyde group.
Answer. (c)	(c)	alpha-D-Glucopyranose is an
732. The principal sugar in blood is	alc	lohexose. (d) Fructose is an aldohexose.

Teekay Page 57	Educational Consult
740. All of the following monosaccharides give	Answer. (c)
the same osaT.one except	<b>747.</b> Which of the following statements is false
(a] Galactose	about sucrose?
(b) Glucose	(a) It is also called table sugar.
(c) Fructose	[b] It may be fermented by yeast to produce
(d) Mannose	alcohol.
Answer. [a]	(c) It reduces Fehling's solution.
741. Mutarotation is a term related to	(d) It does not reduce Tollens' reagent
(a) Interconversion of anomers	Answer. (c)
(b) Relationship of D- and L- families	<b>748.</b> The sugar that yields only glucose on
(r) Hydrolysis of sucrose	hydrolysis is
(d) Number of monosaccharides in a	[a] Lactose
carbohydrate	(b) Sucrose
Answer. (a)	(c) Maltose
742. The mutarotation of glucose is	(d) Fructose
characterized by :	Answer (c)
(a) a change from an aldehyde to ketone	<b>749.</b> A reducing sugar will
structure.	(a) react with Fehling's test
(b) a change of specific rotation from a (+) to a (-	(b) not react with Fehling's test
) value.	[c] have fewer calories
(c) the presence of an intramolecular bridge	(d) always be a ketone
structure.	Answer. (a)
(d) the irreversible change from a-D to the B-	<b>750.</b> Which of the following carbohydrates is not
Dform.	a reducing sugar?
Answer. (c)	(a) Glucose
<b>743.</b> Which of the following statements is false	<i>Th</i> ) Sucrose
about glyceraldehyde?	(c) Fructose
(a) Its IUPAC name 1s 1.2-dihydroxypropanal	(d) Lactose
(h) It is isomeric with 1 3-dihydroxypropanan	Answer. (b)
(c) It is optically active	<b>751</b> . Which of the following carbohydrates will
(d) It shows mutarotation	not give a <i>red</i> precipitate of $Cu20$
Answer. (d)	when heated with Benedict's solution?
744 Common table sugar is	(a) Maltose
	[h i Glucose
(a) Glucose	(c) Sucrose
(c) fructose	(d) Fructose
(d) Maltose	Answor (c)
Answer (h)	752 Which of the following compounds reduces
<b>745</b> Which of the following carbohydrates is	752. Which of the following compounds reduces
sweeter than sucross?	(a) Chicoso
(a) Chucoso	(a) Glucose
(b) Emistance	(b) Sucrose
(b) Fructose	(c) Methanol
(c) Lactose	(d) Acetic acid
(d) None of these	Answer. (0)
Answer. (b)	<b>753.</b> The reagent that can be used to
<b>746.</b> By approximately what factor is the	differentiate an aldose and a ketose is :
sweetness of saccharin greater than that	(a) Bromine water
of sugar?	(b) Fehling's solution
(a) S	(c) Tollcns' reagent
(b) S0	(d) None of these
(c) 500	Answer (a)
(d) 1000	7 <b>54.</b> Starch
Impacting Lives	

Teekay	Page 58	Educational Consult
(a) is a trisaccharide.		(c) conformational isomers
<i>(b)</i> is also called amylose.		(d) cis-trans isomers
(c) is also called amylopectin.		Answer. (a)
(d) is a mixture of amylose + amylopectin.		762. Aromatic hydrocarbons are also called :
Answer. (d)		(a) Arenes
755.The monosaccharide obtained byhydr	olysis	(b) Huckel's compounds
of starch is :	5	(c) Trienes
(a] D-Glucose		(d) Alkoxy compounds
<i>(b)</i> Maltose		Answer. (a)
(c) D-Galactose		763. Which of the following is an aromatic
(d) D-Ribose		compound?
Answer. (a)		(a) CHvCH2CH2Br
756. Which of the following statements is	false	CHa
about cellulose?		713 70 CH CH-CH-
(a) It is a polymer of glucose molecules jot	web in	
$\beta$ -1,4 linkages.		
<i>(b)</i> It is o major component of cotton.		$() () - CH_2CH_3$
(c) It is used in the manufacture of Dacron	fibres.	
(d) It iS u.scd in the manufacture of Rayon	fibres	< >−сн₃
Answer. (c)		(d)
757. Which of the following products is not		Answer. (c)
derived from cellulose?		764. Which of the following compounds is not un
(a) Rayon		aromatic compound?
<i>{b j Insulin</i>		$\bigcirc$
(c) Gun cotton		
<i>{d)</i> Paper		(a)
Answer. (b)		CH3
758. Coal-tar ts the maln source of:		
(a) Aromatic compounds		
<i>(b)</i> Aliphatic compounds	Y	
(c) Heterocyclic compounds		$(h) \approx$
(d) None of these		
Answer. {a)		
759. Which of the following fractions of coa	il-tnr	(c) $\underline{N}$
distillation contains benzene?		
(a) Light oil		C N P
(c) Heavy oll		Answer (b)
(d) Anthracene oli		765 Which of the following compounds is
760 Which of the following fractions of co	al tar	known as aniline?
distillation contains	Jal-lal	Known as annine:
nanhthalono?		
(a) Light oil		
(h) Middle oil		
(c) Heavy oil		
(d) Anthracene		
oilAnswer (b)		
761. What is the term used for isomers that	t arc	
in ranid equilibrium?	i ui c	
(a) keto-enol tautomers		
(h) constitutional isomers		
		I
· · · · ·		

Teekay	Page 59	Educational Consult
ОH		<b>768.</b> The correct lUPAC name lor <i>the</i> following
$\rightarrow$		compound is:
$( \cap )$		(a) p-Nitrobenzene methanoic acid
		(b j o-Nitrobenzoic acid
$(a) \sim$		(c) o-Nitrobenzene methanoic acid
C⊓3		(d) m-Nitrobenzoic acid
		Answer. (b)
		769. The correct IUPAC name for the following
		compound is:
( <sup><i>u</i></sup> ) NH <sub>2</sub>		(a)2-Unioro-5-muoro-1-nyaroxybenzene
		[D1-Hydroxy-2-chloro-5-hldorobenzene (cl1 Eluoro 2 hydroxy 4 chlorobenzene
$\Diamond$		(c)1-Fluoro-5-liveroxy-4-cliforobelizene (d) None of these
		Answer (d) The name should be 2-Chloro-5-
()		flijoronhenol
<b>v</b> - 1		<b>770.</b> The correct IUPAC name for the following
COOF	I	compound is:
	_	(a) 3-Bromo-4-ethyl-1-nitrobenzene
$\sim$		(b) 1-8tliyI-2-bromo-4-nitrobenzene
		[c] 2-Bromo-4-nitroto1uene
(d)		(d) 3-Bromo- 1-ethyl-4-nitrobenzene
Answer (c) [	$a_i$ is Phenol $(h)$ is Toluene and $(d)$	Answer. (d). The numbering is done to produce
is Benzoic aci	d	the smallest total. Therefore, number from the
766. Which of	the following structures represent	ethyl group toward the nitro group. List the
m-dichlorobe	nzene?	substituents in alphabetic order.
CI		771. Which of the following statements are
$\sim$		(a) Ronzono is a planar molecule
		(h) The six carbon-carbon bonds have the same
		length
Ŭ,		(c) Benzene has alternating single and double
$\bigcirc$		bonds.
		(d) (a) and <i>{b</i> )
(b) Cl		Answer (d)
ÇI		<b>772.</b> Which of the following statements is
∠ ci		incorrect about benzene?
		(a) All of the carbon atoms are <i>sp</i> hybridized.
(c)		{b j lt has delocalized electrons.
CI		(c) The carbon-carbon bond lengths are all the
$\Diamond$		(d) All twolve atoms lie in the same plane
	Y	(u) All twelve atoms he in the same plane.
Ý		<b>773</b> Which of the following statements about
a CI		henzene is correct?
Answer, <i>[h1</i>		(a) All of the carbon atoms are rn3 hybridized.
<b>767.</b> The best	name for the following compound	[b] It has no delocalized electrons.
is:	s see see some some some some some	(c) The carbon-carbon bond length is longer
(o) 3-Methyl-b	oromohcnenc	than that of ethane.
(b) 3-Bromoan	iline	(d) It is a planar molecule.
(r) 3-Methylbr	romobenzene	Answer (d)
(d) 3-Bromoto	luene	774. The delocalized u system in benzene is
Answer. (q		formed by a cyclic overlap of 6
Impacting Lives		
r		

Teekay Pa	ge 60	Educational Consult
orbitals.	-	increases the rate of H+ loss.
(s) s		(c) It serves as a Lewis base catalyst by reacting
{h } n		with C12 to generate chloride ions
$\left( c \right) p$		(d) It sorves as a Lowis acid catalyst by reacting
(d)sp		with the Cl2 and thereby activates it toward
Answer (b)		attack by hong ano's n electrons
<b>775</b> Aromotic molecules contain		allack by benz.ene s il electrons.
<b>775.</b> Aromatic molecules contain $\π$		Answer. (Q
electrons.		/B0. Which of the following is not a correct
(a) no		smtement concerning the Friedel- Crafts
$\{b\}$ 4n + 2 (with n being an integer)		acylation of benzene?
(r) 4n + 2 (with n being 0.5)		[o] An alkyl group substitutes for a hydrogen.
(d) 4n (with n an integer)		<i>tb j</i> The benzene ring attacks an acyliuin ion.
Answer. (b)		(c] The acylium ion is resonance stabilized.
<b>776.</b> Which of the structures below would be		(d) The acylium ion is often produced from an
aromatic?		acyl chloride.
+ +		A-nswer t•)
$\dot{\lambda}$ $\wedge$ $^{+}$		7B1. What is the major organic product of the
		reaction between benzene and isobutyl chloride
		in the presence of AiCl3?
		(o) tart-butylbenzene
1 11 111 IV		<i>tb j</i> isobutylbenzene
		(c) n-butylbenzene
		(d) chlorobenzene
(b) 1, 11, and 1V		Answer. (o)
(c) III and IV		7BZ. Which of the following statements are <i>false</i>
		about benzene?
Answer. (c)		(0) It 1s a planar molecule with bond angles 120°-
777. Which of the following is aromatic?		(6) It 1s lmmlsclblc with water forming the lower
(A) cyclopentadienyl cation		layer.
<i>{b)</i> 1,3-cyclohexadiene		(c) It can be converted into cyclohexane by
(C) cyclobutenyl anion		hydrogenation at 200°C in the
(d) cycloheptatrienyl cation		presence of Ni catalyst.
Answer. (d)		(d) It reacts with ethyl chloride in the presence
<b>778.</b> Which <i>of the</i> following is not a correct		of aluminium chloride
statement about the electrophilic		to form ethylbenzene.
substitution mechanism of benzene?		Answer. (6)
(a) Benzene functions as a nucleophile.		7B3. Which statement about the structure of
(b) Formation of a carbocation intermediate i	S	benzene is not true?
the rate-determining step.	-	(a) The two Kekule structures of benzene are in
(c) The carbocati on intermediate contains an	sp3	o9uil ibrium.
hybridized carbon in the ring.		<i>(b)</i> The carbon-carbon bond lengths in benzene
(d) The addition product is a frequently		are greater than the carbon-carbon
observed minor product		double bonds in aliphatic compounds.
Answer. (d)		(c) The molecular geometry of benzene is best
<b>779.</b> What purpose does FeCl3 serve in the		described as planar.
electrophilic aromatic substitution		(d) The stability of benzene ring is much greater
reaction between chlorine and benzene?		than the stability of 1,3,5-
(a) It serves as a radical initiator to produce t	he	cycloheptatriene.
chlorine radical needed to		Answer. (a). Neither of the two Kckule
propagate the chain reaction.		structures f'or benzene has been isolated. No
<i>(b)</i> It functions by destabilizing the		equilibrium exists between these two structures
carbocationic intermediate and thereby		•

Teekay	Page 61	Educational Consult
because they are resonance struc	tures, differing (c) Free	radicals
only in the positions of electron	s. (d) Unch	anged molecules
784. The carbon atoms in a benze	ene ring are : Answer.	(a]
(a) <i>sp</i> hybridized	789. Phe	nol on distillation with Zinc dust gives
<i>(b)</i> sp3 hybridized	(a) Phen	ylzine
(c) sp2 hybridised	(b) Benze	ene
(d) None of these	(c) Cyclo	phexanone
Answer (c)	(d) Benz	oic acid
<b>785</b> Which of the following com	ounds uses Answer	Jh)
only on 2 hybridized carbons for h	and formation? 700 Wh	$\left[ U \right]$
only sp2 hybridized carbons for b	onu iormation: 790. Wi	luono?
	[a] can b	e prepared by treating benkene with
$(a) \succeq$	methyl c	chloride in the presence of A1Cl3.
$\square$	<i>(b)</i> is cor	iverted to benzoic acid on refluxing with
	acid ie Kl	M nO4 solution.
	(c] on re	fluxing with concentrated H2S04 gives
/\	mixture	of ortfo and <i>para</i>
	toluenes	ulfonic acid.
$(c) \subseteq$	(d) can b	e nicratod with concentrated nitric acicl
	to give a	mixture of
())-CH3	ortho and	d meto nitrotoluene.
(d)	Answer	(d)
Answer. [b]	791 Wh	hich of the following reagents will react
<b>786.</b> The C—C bond length in benz	zene is	thy group rather than the
(a) greater than the $C-C$ bond len	gth in ethane.	ring in methylbongene?
(b) ) shorter than the $C-C$ bond let	ngth in	ring in methylbenzene:
ethylene		The in the presence of uv light
(c) Same as titat of C-C bond lengt	th In ethylene	t in the presence of AICI3
(d) intermediate between C-C box	nd length in	OCI in the presence of AlCI3
ethane and C—C hand length in a	thylene (d) Hydr	ogen in the presence of nickel
Anguor (d)	Answer	. (a)
707 Which of the following comp	792. Wh	iich of the following can be made by the
787. Which of the following comp	action of	CH3Cl on benEene in the
aromatic	presence	e of aluminium chloride?
(a)	(a) Ethyl	benzene
	(b) o-Xvl	ene
	(c) Chlor	cibenzene
	(d) m-Xy	zlene
(b) <u>N</u>	Answer	(h]
	793 For	reactions of ethylbenzene, the ethyl
	, , , , , , , , , , , , , , , , , , , ,	reactions of early idenzene, the early
	(a) ortho	director
(c)	(b) or the	nora director
		director
(d)	(d) ortho	o-meta director
	Answer.	{ <i>b</i> }
Answer $(h)$ Use the concent of a	romaticity and 794. Tol	uene reacts with bromine in the
Huckel rule to arrive at the	presence	e ot uv light to give
annast anguar	(a) m-bro	omotoluene
<i>correct</i> answer.	(b) Benz	yl bromidc
/ oo. Unaracteristic reactions of al	<i>[c]</i> o-Bro	omotoluene
nydrocarbons are initiated by	(d) Benz	oyl bromide
(a) Electrophiles	Answer	(Ď)
<i>(b)</i> Nucleophiles		- /

#### Teekay

795. Toluene reacts with chlorine in the presence of AlC13 to give
(a) o-Chlorotoluene
(b) o- plus p-Chlorotoluene
(c) m-Chlorotoluene
(d) o- plus m-Chlorotoluene
Answer. [b)
796. Toluene undergoes oxidation to give
(a) Benzyl alcohol
(b) Quinone
(c) Benzaldehyde
(d) Benzoic acid

## Answer. (d)

797. Ethylbenzene undergoes oxidation with acidic potassium dichromate to give



# Answer. (d)

798. Oxidation of cumene with acidic K2Cr207 gives

- (a) Phenylacetic acid
- *(b)* Benzaldehyde
- (c) Benzyl alcohol
- (d) Benzoic acid

# Answer. (d)

799. Oxidation of toluene with chromyl chloride gives benzaldehyde. This reaction is known as

- (a) Perkin reaction
- (b) Bcnzoin condensation
- (c) Etard's real tion
- (d) Ozonolysis

Answer. (c)

800. Oxidation of p-xylene with acidic potassium dichromate gives



801- Cresols on distillation with z.inc dust gives (a) o-Xylene [b] Benzene (c) o- plus p-Xylene (d) Toluene Answer. (d) 802. An organic compound [A] has a molecular weight 78 and an empirical formula CH. [A ] is resistant to attack by oxidizing agents but undergoes substitution and addition reactions. X-ray analysis shows that the distance between adjacent carbon atoms is 1.39d (Intermediate between single and double bond lengths). The evidence suggests that compound (A) is (a) Benzene (b) Cyclohexane (c) Toluene (d) n-Hexane Answer. (a) **803.** Benzene undergoes substitution reaction more easily than addition reaction because [*a* ] it has a cyclic structure (b) it has three double bonds (c) it has six hydrogen atoms (d) there is delocalization of electrons Answer. (d) 804. Benzene reacts with concentrated HN03 in the presence of concentrated H2S04 to give nitrobenzene. This reaction is an example of [a] Electrophilic addition

- (b) Nucleophilic addition
- (c) electrophilic substitution
- (d) Nucleophilic substitution
- Answer (c)

Teekay	Page 63	Educational Consult
805. Which of <i>the</i> following agent	ts is used in	811. In sulfonaLion of <b>benzene</b> , the attacking species
order to make benzene react with	n concentrated	is
nitric acid to give nitrobenzene?		(a} H+
(a) Concentrated H2S04		(b) 502
{b] FeCl3 catalyst		(c) 503
(c) Lindlar's catalyst		(d)
(d) Ultraviolet light		Answer. (r)
Answer. {a}		812. Consider the following reaction :
806 Which of the following agen	ts is used in	CH-CH-
order to make benzene react with	n acetyl chloride	1 2 - 3
to give acetophenone?		
(a) Ultraviolet light		HCI + CH3CH2CI + HCI
(h) AlCl3 catalyst		
(c) Platinum catalyst		The catalyst used to complete the above reaction
(d) Al 203 catalyst		
Answer (h)		$\{a\}$ LIAIH4
807 Which of the following ager	nts is used in	(b) AICI3
order to make henzene react wit	h bromine to	(c) Na
give bromubentene?		(d) KOH
(a) Illtraviolet light		Answer. (b)
<i>(a)</i> Ottraviolet light		<b>813.</b> Benzene reacts with H2 at 150°C at 30 atm
(c) Nickel catalyst		in the presence of Ni catalyst to
(d) Al203 catalyst		give
Answer (h)		(a) Cyclohexane
<b>808</b> In the Friedel-Craft acetylat	ion of an	[b) Cyclohexene
aromatic ring the role of the AlCl	3 is to	(c) n-Hexane
	5 15 10	(d) No reaction occurs
		Answer. (a)
(a) Form a $+$ ion.		814. Benzene reacts with chlorine in the
(b) Function as a Lewis base		presence of FeCl3 catalyst to form
(c) Chlorinate the aromatic ring		(a) Hexachlorobenzene
(d) Withdraw electrons from the	aromatic ring	(b) Chlorobenzene
Answer. (A)	U	(c) Hexachlorocyclohexane
<b>809.</b> The electrophile which is cons	sidered to be	(d) Benzyl chloride
the active agent in the nitration	of	Answer. {b }
benzene is		815. Benzene reacts with acetic anhydride in the
$( ) NO^{-}$		presence of A1C13 to form
(a) NO <sub>2</sub>		(a) Acetophenone
(b) NO <sup>+</sup>		(b) Benzophenone
(0) NO		(c) Phenylacetic acid
$(c) NO^{\dagger}$		(d) Phenyl acetate
(0) 1102		Answer (a)
(d) HNO+		816. Benzene reacts with benzoyl chloride in the
(d) mto :		presence of anhydrous
Answer. (c)		aluminium chloride to form
810. In chlorination of benzene, I	FeCl3 is used to	(a) Benzyl chloride
generate		(b) Benzaldehyde
$\sim C1$ :		(C) Benzal chloride
(a)		(I) Benzopnenone
(B) (I+		Answer. (a)
		817. Benzene undergoes Friedel-Crafts reaction
		with isopropyl alcohol in the
Answer. (b)		presence of H2SU4 catalyst to give

Teekay	Page 64	Educational Consult
(a) n-Propylbenzene		Answer. {b}
<i>{b</i> } Benzophenone		825. Which of the following will undergo
(c) IsopropylbenEene		substitution in the <i>ortho</i> and part
(d)Nothing happens		positions rather than in the meta position?
Answer. (d)		(a) Nitrobenzene
818- Benzene reacts with propene in	the	(b) Benzoic acid
presence of H2SD4 catalyst to give	the	(c) Acetanilide
(al n-PronylhenEene		[d] Benzaldehyde
(h) Benzonhenone		Answer (c)
(c) Cumene		826 Phenol is an <i>ortho-nara</i> director because
(d) Nothing hannens		the hydroxyl group :
Answer (c)		[a] denotes electrons that increase electron
<b>810</b> Ozonolysis of honzono givos		donsity at ortho and nara positions favoring
(a) Formis asid		nucleon hilis attack
(a) FOI IIIC actu		(b) denotes electrone that increase electron
(D) GIYOXAI		(D) utilities electrons that increase electron
(d) Chusing		electrophilic attack
		electrophilic attack.
Answer. [b]		(c) donates electrons to the ortho and para
820, Gammexane Is		positions and attracts circtrens away from meta
(a) Hexachioroethane		positions ravoring nucleophilic attack of the ring.
		(d) donates electrons to the <b>ortho</b> and para
(c) Hexachlorocyclonexane		positions and attracts electrons away from
(d) TNT		metapositions favoring electrophilic attack on
Answer. (c)		the ring.
821. Which group is an activating subst	ituent?	Answer. (b)
(a] —NO3		827. Compared to benzene, nitratraion of
(b)Cl		toluenetakes place at
(c) -0 H		(a) Same rate
(d) -F		(b) Faster rate
Answer. (c)		(c) Slower rate
<b>822.</b> Which of <i>the</i> following compounds	s reacts	(d) Can not predict
most rapidly with HN03/H2SO4?		Answer. (a)
(a] toluene		828. Electron-withdrawing groups are <i>meta</i>
(b) anisole		directors because :
(c) nitrobenzene		(a) the carbonium ion intermediate has a
(d) benzonitrile		negative charge on the meta position.
Answer. {b)		(b) the more stable resonance hybrid occurs
<b>823.</b> In electrophilic aromatic substitut	ion	with <i>meta</i> attachment of the
reactions a chlorine substituent		electrophile.
		(c) the less stable resonance hybrid occurs withmeta
(a) is a deactivator and a m-director		attachment of the
{b ) is a deactivator and an o,p-director	•	electrophile.
(c] is an activator anü a m-director		(d)the carbonium ion intermediate has a
(d) is an activator and an o,p-director		positive charge on the <i>meta</i> position.
Answer. (b)		<b>Answer.</b> <i>(b)</i> When the electrophile attaches to
824. Which of the following substitue	nts is an	the benzene ring, a carbonium ion is formed.
ortho and part director and ring		This carbonium ion is relatively stable because
deactivating?		the charge can be delocalized about the ring.
(a) -NH2		ortho or para attack results in a resonance form,
(b) —Cl		with the positive charge on the carbon bearing
(c) —0CH3		the electron-withdrawing group (highly
(d) —0H		unstable). This resonance form does not occur
		-

with <i>meta</i> placement of <i>the</i> incoming electrophile. B29. Which compound undergoes substitution reactions faster than benzene?	<ul><li>831. Which of the following compounds undergoes nitration most readily?</li><li>[a) Benzene</li><li>(b) Telegoes</li></ul>
electrophile. B29. Which compound undergoes substitution reactions faster than benzene?	undergoes nitration most readily? [a) Benzene
B29. Which compound undergoes substitution reactions faster than benzene?	[a) Benzene
reactions faster than benzene?	$(h_{1})$ T always
	<i>ip i</i> loluene
	(c) Nitrohenzene
NO <sub>2</sub>	(d) Benzoic acid
1 2	Answer (b)
	022 Which of the following compounds is most
	852. Which of the following compounds is most
(a)	(a) Demonstrated?
CH	(a) Benzene
0.13	[b] Chlorobenzene
Ĭ	(c] Toluene
$\wedge$	[d]Nitrobenzene
	Answer. (c)
	833- p-Nitrotoluene we on treatment with chlori
$(U) \sim$	in the presence of FeCl3 gives
Br	(a) m-Chlorotoluene
	(b) 2-Chloro-4-nitrotoTuene
	(c) p-Chlorutoluene
	(d) 2-Nitro-4-chlorotoluene
(c) V	Answer. {h)
NH <sub>2</sub>	<b>834</b> How many isomeric dichlorobenzenes are
	there?
	(a) 3
	(a) $(b)$
(d)	
Answer (d) Remember that activating	
(aloctron donating) groups on honzono result in	$\left( a\right) 12$
anhanced electrophilic substitution	Answer. (a)
920 Which compound undergoes substitution	835. Which of the following compounds has
reactions slower than honzono?	maximum dipole moment?
	(a) p-Dichlorobenzene
	(b) m-DichlorobenEene
	(c] o-Dichlorobenzene
	(d) Carbon tetrachloride
(a) 😒	Answer. (c)
NHCH <sub>3</sub>	836. Which of the following <i>reagents</i> is most
	suitable for preparing chlorobenzene
	from benzene?
(b)	(a) Aqueous chlorine
Ę	(b) Chlorine in the presence of ultraviolet light
	(c) Chlorine in acetic acid
	[d] Chlorine in the presence of FeC13
(c)	Answer (d)
OCH <sub>3</sub>	837 Which of the reactions between the
	following pairs of reagants proceeds by a
	nonowing pairs of reagents proceeds by a
(d) V	mechanism that may be described as an
Answer. (c). The halogen (—F) is the only	electrophilic substitution?
Reactivating group because of i1	(a) Benzene + Kr2/FeHr3
electronegativity and its low resonance	(b) Propene + Br2
potential.	(c) Acetone + NH2NH2
	(d) Ethyl bromide + <i>aq</i> KOH
	Ancwor (a)

Teekay Page 66	Educational Consult
838. Aryl halides are less reactive towards	<b>844.</b> DDT is prepared by <i>the</i> reaction of
nucleophilic substitution as compared	chlorobenzene with (in the presence of
to alkyl halides due to	conc. H2504)
(a) Inductive effect	(a) Chloral
(h) Resonance stabilization	(h) Chlorine
(c) Tautomerism	(c) Chloroform
(d) Stercoisomerism	(d) Carbon totrachlorido
Answer (h)	Anguer (2)
920 Which of the following compounds would	Allswei. (d) $\mathbf{QAF}$ Which of the following reason to some he
be unaffected by beiling acueous	<b>645.</b> Which of the following reagents can be
	used to distinguish between
(a) Chlanaethana	chiorobenzene and allyl chioride?
(b) A set al de la de	$\begin{bmatrix} a \end{bmatrix} H Z / N I$
tb) Acetaldenyde	(b) Br2 in CC14
(c) Chlorobenzene	(c) Zn/HCl
(d) Ethyl acetate	(d) NH2NH2
Answer. (c)	Answer (b]
840. Hydrolysis of benzal chloride gives	846. Which of the following reagents can be
(a) Phenol	used to distinguish between
(b) Benzaldchyde	chlorobenzene and bcnzyl chloride*
(c) Benzyl alcohol	(a) alcoholic AgNO3
(d) Benzoyl chloride	(b) Br2 in CCI4
Answer. (b)	(c) KCN
841. Hydrolysis of berizotrichloride gives	(d) Br2 in H2O
(a) Benzophenone	Answer. (a)
(b) Benzoic acid	847. An organic compound (3) was heated with
(c) Benzyl alcohol	a solution of NaOH. The resulting solution was
(d) Phenol	cooled, acidifted with dilute HN03, and then
Answer. (b)	AgN03 solution was added. This gave a white
842. Which reagent can be used to carry out the	precipitate. The compound (A) may be
following conversion?	
Cl Cl	
	(a)
$()$ $\xrightarrow{()}$ $()$	(( ))—CH2
	(b)
T T	
CI CI	( ))— COOH
(a) HN03 + H2S04	(c)
(b) NaNO2	
(c) NaN02 - HCl at 5°C	(( ))—CHI
(d) N02	(d)
Answer. (a)	Answer. (b)
<b>843</b> . The following compound is called?	848 Which of the following statements is
	false/Incorrect about nitrobenzene?
	(a) is formed by nitration of henzene at
	temperatures below 60°C
CCI:	(h) can be further nitrated to give m-
(a) Chloral	dinitrohenzene
(h) DDT	(c) can be ovidized with $KMn04$ to give honzoic
(c) Lindane	acid
(d) BHC	(d) can be reduced with En /UCI to give aniling
Answer (h)	(u) can be reduced with Sil/ not to give annine.
	AllSwel. (C)

(d) can be reduced with 5n/HCI to give aniline. Answer. (c)

Teekay	Page 67	Educational Consult
<b>849.</b> Which of <i>the</i> following compounds re	eacts	(c) can be nitrated with concentrated HNO3 to
least rapidly in electrophilic		form p-nitroaniline.
substitution reactions?		(d) gives an intense green color with a
(a) Nitrobenzene		suspension of bleaching powder in water.
(b) Phenol		Answer. {b}
(c) Bromobenzene		856. Which of the following statements is false
(d) Tolucne		about benzylamine?
Answer. {a)		(a) can be prepared by reduction of benzonitrile
<b>850.</b> Nitrobenzene reacts with Br2 in the		with LiA1[J4.
presence of FeCl3 to give		(b) cannot be prepared by reduction of
(a] o-chloronitrobenzene		benzamide with LiAlH4.
<i>(b)</i> o-bromonitrobenzene		(c] reacts with nitrous acid to give benzyl
(c) m-chloronitrobenzene		alcohol.
(d) m-bromonitrobenzene		(d) reacts with KMn04 to give benzoic acid.
Answer. (d)		Answer. {b)
<b>851.</b> Nitration of nitrobenzene with		<b>857.</b> Aniline is prepared by
HN03/H2S04 gives :		(a) the reaction of benzene with ammonia
(a) m-dinitrobenzene		(b) the reduction of nitrobenzene with 5n/HCI
(b) n-d initrobenzene		(c) the dehydrogenation of nitrobenzene
(c) p-dinitrobentene		(d) the reaction of nitrobenzene with 12/Na0H
(d) m-nitrobenzenesulfonic acid		Answer. (b)
Answer. (a)	.1	858 Benzamide reacts with Br2 and KOH to give
<b>852.</b> Nitrobenzene undergoes reduction with	th	(a) Benzene
zinc and KOH to give		(b) Benzylamine
(a) Annine (b) Hydrogoborgono		[C] Aniline
(a) Az obenzene		(a) Benzonitrile
(d) None of these		Answer. (D) 950 When ammonia IS added to an allud halida
Answer (h)		in the presence of base
<b>853</b> Reduction of n-nitrotoluene with Sn +	нсі	(a) primary amines form
gives ·		(b) amides form
(e) p-methylaniline		(c) nitrated allot halides form
<i>(b)</i> Toluene		(d) quaternary ammonium salts form
(c) m methylaniline		Answer (a). Salts of 1° amines can be prepared
(d) <i>N-M</i> ethylaniline		from ammonia and alkyl halides. 1 reating the
Answer. (a)		ammonium salt with base gives a 1° amine.
854. Which of the following statements is	false	$\widehat{NH_{0}}$ + $\widehat{R}$ $\widehat{N}$ $\widehat{NH_{0}}$ $\widehat{CI}$ $\widehat{NaOH}$ $\widehat{R}$ $\widehat{NH_{0}}$ + $\widehat{NaX}$ + $\widehat{H_{0}OH}$
about aniline		860 Aniline reacts with bromine water to form
(a)lt is a primary aromatic amine.		(a) Bromobenzene
(b) ]t is a weak base which forms .salts with	n	(h) <i>m</i> -Bromoaniline
acids.		(c) 2.4.6-Tribromoaniline
(c) It liberates nitrogen on treatment with	ice-	[d] o- plus p-Bromoaniline
cold nitrous acid.		Answer. (c)
(d) It can be prepared by the reduction of		<b>861.</b> Which of the following will give a
nitrobenzene with tin and HCl.		precipitate with excess bromine water?
Answer. (c)		(a) Bromobenzene
855. Which of the following statements is		(b) Aniline
correct about aniline?		(c) Nitrobenzene
(a) diagonations to have a set of the set of		(d) Toluene
(D) diazotizes to benzenediazonium chloride	9	Answer. (b)
when treated with ice-cold mitrous acid.		862. Aniline reacts with nitrous acid at low
		temperatures to give
Impacting Lives		

3

NH<sub>2</sub>

 $NH_2$ 

NH<sub>2</sub>

872. Which of the following compounds is most

ii

С

3

Answer. (a)

(c)

(d) NH3

basic?

Teekay (a) a N-nitrosoamine 868. Rank the following compounds in order of *{b]* a nitrile increasing base strength : (c) a diazonium salt (d) a nitrite salt NHa Answer. (c) 863. When aniline is heated with chloroform (a) 1, 2, 3, 4 and alcoholic KOH, the product is [b] 4, 2, 3, 1 (a] Benzonitrile (c) 1, 4, 3, 2 *{b]* p-Chloroaniline (d) 4, 3, 1, 2 (c) Phenylisocyanide Answer. (d). Of the aromatic amines, the (d) m-Chloroaniline weakest base is the one with the Answer. (c) electron-withdrawing substituent: (4) is the 864. Aniline reacts with acetic anhydride to give weakest followed by aniline (3). Compound (2) is the strongest base because its (a) N-Methylaniline electrons are not delocalized (b) p-Aminoacetophenone effectively into the aromatic ring, and it is (c) Acetonilide therefore a primary amine, and its pka (d) m-Aminoacetophenone is slightly lower than that of ammonia. Answer. (c) 869. Which of the following compounds is most 865. Aniline undergoes oxidation with basic? Na2Cr207/H2S04 to give (a)Aniline (a) Schiff's base (b) Benzylamine *(b)* p-Benzoquinone [c] Acetanilide (c) Phenol (d) p-nitroaniline (d) Benzoic acid Answer. (6) **Answer**. {*b* ] **870-** Which of the following is most basic? 866. Which of the following reagents does not (a) CH3NH2 react with aniline? (b) (CH3)3N (a) Acetyl chloride (c) (CH3)2NH *(b)* Acetic anhydride (c) Ammonia (d) Nitrous acid (4 Answer. (c) **Ånswer**. (c) 867. Rank the following compounds in order of **871.** Which of the following is most basic? increasing base strength : (o) (Cll3)2NII SNH, NFt CFh NN, (b)(a) 1, 2, 3

- *{b)* 3, 2, 1 {*c*} 2, 3, 1
- (d) 2, 1, 3

**Answer.** (c). The aromatic amines are weaker bases than the nonaromatic cyclic amines because the lone-pair electrons on nitrogen are delocalized over the aromatic ring. This delocalization stabilize us the compound but also decreases the ability of the nitrogen to act as a base. Compound (3) is a stronger base than aniline because of the electron-releasing substituent on the ring.
NL1y

Page 69

NH<sub>2</sub>

Educational Consult



Answer. (a) 873. Which of the following is least basic? (a) Aniline {b) p-Nitroaniline (c) Ammonia (d) Trimethylamine Answer. (b) 874. Which of the following is least basic? (a) CH3NH2 (b) (CH3)3N (c) (CH3)2NH

(d) Answer. (d)875. Which of the following compounds is least basic?

# (a)NH (b)NH<sub>2</sub> NO. (c)NH2 NO<sub>2</sub> (d)Answer. (c) 876 Which of the following compounds will be roost soluble in water at room temperature? NH2 VaNC CHC Cl. 0° (a) Phenol (b) Phenylammonium chloride (c) Benzoic acid (*d*) Benzylamine Answer. {b} 877. The diazonium salts are the reaction products of the reaction of nitrous acid with (a) Primary aliphatic amine (b) primary aromatic amines (c) secondary aliphatic amines (d) secondary aromatic amines Answer. {b} 878. Preparation of a diazonium salt from a primary aromatic amine is known as (a) Coupling reaction (b) Sandmeyer reaction (c) Diazotization (d) Corey-House synthesis Answer. (c)

**879.** Which of the following reagents is used to prepare benzenediazonium

Teekay	Page70	Educational Consult
chloride from aniline?		(a) Benzonitrile
(a) NaNO2 + HCI		(b) Benzoic acid
<i>{b]</i> LiAlH4		(c] Nitrobenzene
(c) NH2 N H2 + KOH		(d) Benzenediazonium chloride
(d) NaOH		Answer. (a)
Answer. (a)		<b>887-</b> Benzenediazonium chloride yeld with
<b>880.</b> Which of the following are optimum		hypophosphorus acid to form :
temperature conditions for making		(a) Phenol
benzenediazonium chloride from aniline?		(b) Benzaldehyde
(a) 0°C to 10°C		(c) Aniline
<i>(b)</i> 20°C to 25°C		[d] Benzene
(c) 30°C to 40°C		Answer. (d)
(d) 45°C to 50°C		888. Benzenediazonium chloride reacts with
Answer. (a)		phenol to form :
881. benzenediazonium chloride <i>reacts</i> with	1	(a) <i>p</i> -chlorophenol
warm water to give	-	(b) Chlorobenzene
(a) Aniline		(c) p-Hydroxyazobenzene
<i>{b</i> } Phenol		(d) DDT
(c) Benzene		Answer. (c)
(d) Chlorobenzene		889. Phenol is used
Answer. (h)		(a) in alcoholic beverages
<b>882.</b> Bromobenzene can be prepared by tre	eating	(h) as anesthetic
aniline with	Juling	(c) in antisentics
(al Conc HBr		(d) as moth repellant
(b) Br2/FeBr3		Answer. (c)
{c} CuBr		<b>890.</b> Which of the following compounds is
(d) Nitrous acid then CuBr		aspirin
Answer. (d)		(a) Methyl sallcylate
883. Chlorobenzene can be prepared by		(b) Salicylic acid
treatment of aniline with		(c) Phenyl salicylate
(a) Cuprous chloride		(d) Acetylsalicylic acid
(b) Chlorine in the presence of UV light		Answer. (d)
(c) Nitrous acid followed by treatment with	CuC1	891. When phenol is treated with neutral FeC13
(d) Chlorine in the presence of FeCl3		solution, it develops
Answer. (c)		(a) Violet color
<b>884.</b> lodobenzene can be prepared by		(b) Yellow color
(a) treating chlorobenzene with 12 using Fe	Cl3	(c) Green color
catalyst.		(d) Nothing happens
<i>(b)</i> treating phenol with [2 in Na0H solutio	n.	Answer. (a)
(c) treating henzenediazonium chloride wi	th Kl	892. Which of the following ccmpounds will
(d) treating benzene with CH3l using A1Cl3		react with PC15 least vigorously?
catalyst.		[a] Ethanol
Answer. (c)		[b] Acetic acid
885. Benzenediazonium chloride reacts wi	th Kl	(c) Phenol
to form :		(d] Benzoic acid
(a) Benzenediazonium iodide		Answer (c)
(b) m-Diiodobenzene		893. Sodium phenoxide reacts with CO2 at
(c) lodobenzene		126°C under 6 atm pressure to give
(d) n- plus p-Dr iodobcnzcnc		salicylic acid. This reaction is called
Answer. (c)		(a) Kolbe's reaction
886. What is the major product of the follow	wing	(b) Perkin reaction
reaction?		(c) Wurtz reaction

Impacting Lives

(d) HVZ reaction

#### (d) CH3CH2-S-S—CH2CH3 Answer. (c)

(c)(3) > (1) > (2)

Answer. (s) 894. Phenol reacts with excess bromine water to give

(a) o- plus p-Bromophenol

(b) Bromobenzene

(r) 2,4,6-Tribromophenol

(f) m-Bromopheno1

Answer. (c)

895. Resorcinol on distillation with zinc dust gives

(a) Benzene

(b) Cyclohexane

(c) Toluene

(d) m-Xyleee

Answer. (a)

896. Bakelite has the following characteristics (a) A polymer made from phenol and formaldehyde

*((b)* A thermosetting plastic

(c) Can be used as an adhesive

(d) All of the above

Answer. (d)

897. Anisole is formed when phenol is treated with

(a] CH3I/NaOH

(b) CH3CH2I/NaOH

(c) CHCI3/NaOH

(d) Acetic anhydride

#### Answer. (a)

898. Anisole on heating with concentrated HI gives(a) lodobenzene(b) Phenol + CH31(c) lodobenzene CH20 H

(c) lodobenzene + CH30 H(d) Phenol + CH30 H

Answer. {b)

Answer. {D

**899.** Which group forms the strongest H-bonds to water molecules?

(a) Alcohols

(b) Ethers

(c) Phenols

(d) All equally strong

Answer. (c)

**900.** The compound which is most capable of hydrogen bonding is :

(a) CH30CH2CH3

*{b)* CH3CH 2CH 2CH3

OH

**901.** Rank the following molecules in decreasing order of boiling points :



(d) (2) > (1) > (3)
Answer. (a]. Hydroxyl groups can more easily form intramolecular H-bonds in the ortho position than in the meta or part positions.
These intramolecular H-bonds decrease boiling point Therefore, compound (3) has a higher boiling point than compound (2). Compound (2), with two hydroxyl groups, has a still higher boiling point than phenol,

902. Rank the following compounds in order of increasing boiling point :



Teekay Page 72 906. Which of the following compounds is least acidic? (a] HCI (b) Phenol (c) Acetylene (d) Picric acid Answer. (c) **907.** Which of the following compounds is least acidic? (a) Formic acid *{b )* Ethanol (c) Acetic acid (d) Phenol Answer. (b) 908- Which of the following compounds is most acidic? (a) Formic acid *{b]* Ethanol (c) Acctic acid (d) Phenol Answer. (a) 909. Which of the following compounds is most acidic? (a] Water (b) Cyclohexanol (c) Ethanol (d) Phenol Answer. (d) 910. Which one is most acidic? (a) Hexanol *{b]* Phenol (c) Water (d) Diisopropyl ether Answer. (b) 911. Which of the following compounds is most acidic? (a) o-Cresol (b] p-Nitrophenol (c) p-Cresol (d) p-Chlorophcnol **Answer**. (b) 912. Which of the following compounds is most acidic? (a) Phenol *(b) m*-Nitrophenol (c) m-Cresol (d) m-Bromophenol Answer. (b) **913.** Which molecule is the most acidic?



(d) (2) and (3) have similar acidities
Answer. (b). Acidity is enhanced when the conjugate base is further stabilized. By placing an electron-withdrawing group on the ring, the phenoxide ion ix further stabilized when the charge spreads over more atoms (greater delocalization). The methoxy group is electron-releasing and destabilizes the phenoxide.
914. Which of the following compounds is most



## Answer. (r)

**915.** Which of the following compounds is least acidic?



# Answer. {b)

**916.** Arrange the following compounds in order of increasing acidity:



Teekay	Page73	Educational Consult
(a) phenols are soluble in NaOH, b	out alcohols are	at 180°C to give :
not		(a) Mandelic acid
<i>{b]</i> alcohols are soluble in NaOH, I	out phenols are	(b) Cinnamic acid
not	1	(c) Malonic acid
(c) phenols are soluble in NaHCO3	3, but alcohols	(d) Benzoic acid
are not	,	<b>Answer.</b> <i>{b].</i> This is Perkin reaction.
(d) alcohols arc soluble in NaHCO	3. but phenols	<b>924.</b> Which of the following reagents will give an
are not		addition reaction with
Answer. (d)		benzaldehvde?
918. Benzovl chloride reacts with	hvdrogen in	(a) NaHSO3
the presence of palladium and	, 0	(b) 12/Na0H
barium sulfate to give :		(c] H2S04
(a) Benzvl alcohol		
(b) Benzaldehyde		<b>V</b>
(c) Cinnamic acid		
(d)Benzophenone		$_{\rm CH_3}$ – C – Cl
<b>Answer.</b> (b) This is Rosenmund re	eduction.	$\begin{pmatrix} u \\ \end{pmatrix}$
<b>919.</b> Cinnamaldehyde is prepared	d by the	<b>025</b> Bonzaldohydo roacts with a mixture of
treatment uf :	5	<b>525.</b> Delizable injue reacts with a linkture of
(a) benzaldehyde with acetaldehy	de in the	acid to give
presence of NaOH		(a) n Nitrohonzaldohudo
<i>(b)</i> benzaldehyde with formaldeh	vde in the	(b) a Nitrobaaraldabyda
presence of NaOH		(c) n-Nitrobenzoic acid
(c) benzophenone with acetaldeh	yde in the	(d) m-Nitrobenzaldehyde
presence of NaOH		Answer (d)
d benzophenone with formaldeh	vde in the	926 Acetophenone reacts with Br2 in the
presence of NaOH		presence of FeCl3 to form ·
Answer. [a]		(a) m-Bromoacetonhenone
<b>920.</b> Benzaldehyde reacts with		(h) m-Dibromobe <i>ozene</i>
methylmagnesium iodide (followe	d by	(c) $n + n$ Bromoacetonhenone
hydrolysis) to form		[d] p-Dibromobenzene
(a) l-Phenytethanol		Answer (a)
<i>{b)</i> Acetophenone		927- Which of the following statements is NOT
(c) 2-Phenylethanol		correct :
(c) Benzophenone		(a) Tollens' test is given by both aliphatic and
Answer. (e)		aromatic aldehydes.
<b>921.</b> Which of the following will u	indergo Aldol	(b) Fehling's test is given by aliphatic aldehydes.
condensation?		lt is not given by aromatic
(a) Acetone		aldehydes.
(b) Benzaldehyde		(c] Iodoform test is given by methyl ketones and
(c) Benzoic acid		acetaldehydes.
(d) Benzophenone		(d) Lucas test is given by both aliphatic and
Answer. (a)	1 11	aromatic aldehydes.
922. When benzaldenyde is heate	a with	Answer (d)
(a) Denzyl alachal		928. Benzaldehyde can be distinguished from
(a) Benzyi alconol		acetophenone by using :
(c) benzoic acid		(a) tollens' reagent
(d) Benzyl alcohol + Sodium honz	oatc	(b) Lucas reagent
Answer, (d) This is Cannizzaro re	Paction	(c) Felling's reagent
923. Benzaldehvde reacts with a	retic anhydride	(d) All of these
in the presence of sodium acetate		Answer. (a)
in the presence of sourum actual	-	929. Benzaldehyde does not react with
Impacting Lives		

TEEKay	Teekav
--------	--------

(a) Tollens' reagent

(d) 4

Answer. (c)

(b) Phenylhydrazine
(c) Fehling's solution
t< NaHSO3
Answer. (c)
930. Whi¢h of the fcillpwing will give a positive
Tollens' test?
0
(b) <>= 0
(c) <u>CH</u> <u>3</u> c H <sub>2</sub> c" −H

### Answer. (c)

(d)

931. Which of the following compounds yields a yellow precipitate when warmed with a mixture of iodine In aqueous sodium hydroxide?

(a) Acetophenone

≻=O

(b) Benzaldehyde

(c) Renzophenone

(d) Propanal

Answer. (a)

932. A compound [*A*) gave a positive iodoform reaction and left a solution containing a monocarboxylacc Ion, Which of the following could {*A*} be?



#### Answer. (c)

933. how many isomers containing a benzene ring correspond to the formula CfiH4(COOH) 2

- (a) 1
- *{b)* 2
- (c) 3

934. Which of the following is not a carboxylic acid? (a) Malonic acid (b) Acetic acid (c) Picric acid (d) Adipic acid Answer. (c) 935. Phenylmagnesium bromide reacts with CO2 followed by acid-hydrolysis to form : (a) Phenol (b) Benzoic acid (c) Bromobenzene [d] Acetophenone Answer. b) 936. Benzoic acid and diphenyl ketone are both solids Insoluble in water. A mixture of these compounds can be separated if we (a) add the mixture to a solution of NaOH in water, and filter. (b) add the mixture to a solution of HCI in water, and filter. (c) dissolve the mixture in benzene and boil otI the benzene. (d) It 1s impossible to separate a mixture of two solids when both are insoluble in water. Answer. (a) 937. Which of the following reactions does not produce benzene as one of the products? (a) Treating benzenediazonium chloride with H3P02 (b) Distilling phenol with zinc dust (c) Fusing sodium benzoate with NaOH (d) Fusing sodium benzenesulfonate with NaOH Answer.(d) 938. Which of the following esters produces butyric acid and phenol on hydrolysis? (a) Phenyl butyrate *{b]* Butyl benzoate (c) Benzyl propionate

- [*d* ) Phenyl propionate **Answer**. (a)
- 939. Benzoyl chloride on basic-hyd rolysis

(Na0H/H20) gives :

(a)Benzoic acid

(b) Methyl benzoate

Teekay	Page 7S	Educational Consult
(c) Sodium benzoate		(a) o-Nitrobenzoic acid is more acidic than <i>p</i> -
(d) Ethyl benzoate		nitrophenol
Answer. (C)		(b) o-NitrobenEoic aCid is tess acidic than p-
940. Sodium benzoate on heating with s	soda-lime	nitrophenol
gives :		(c) o-Nitrobenzoic acid and p-nitrobenzoic acid
(a) Sodium phenoxide		have the same acid strength
(b) Benzene		(d) Nonc of these statements is true
(c) Benzaldehyde		Answer. (a)
(d) Benzophenone		946. The order of increasing acidity is:
Answer. (b)		(a) chloroacetic acid > formic acid > benzoic acid
941. The product of the following reacti	on is :	(b) benzoic acid $\gg$ formic acid $>$ chloroacetic acid
0		(c) all are the same
		[d] not determinable from available data
		Answer. (a)
0		947. Which of the following is the strongest
		acid?
$(a)$ $(b)$ $-b$ $-c$ $-c$ $H_2CH_3$		(a) Trichloroacetic acid
		(h) Phenol
		(c) Acetic acid
		(d) Benzoic acid
		Answer (a)
( ))—с – сн <sub>2</sub> сн <sub>3</sub>		948 Which of the following is the strongest
(c)		acids
		(a) Ethanol
		(b) Phenol
Answer (h)		(c) Anisole
942 Methyl henzoate on hydrolysis give	20	(d)Benzoic acid
(a) Acetic acid		Answer. (d)
(b) Benzoic acid		949. Wliich acid is weaker than benzoic acid?
(c) Picric acid		(a) p-Methylbenzoic acid
(d) Phenylacetic acid		<i>(b)</i> p-Chlorobenzoic acid
Answer [h]		<i>{c)</i> p-Nitrobenzoic acid
943 Aspirin is produced by heating sali	icylic acid	(d) o-ChlorobenEoic aCid
with	leyne dela	Answer (a)
(a) acetic anhydride in the presence of		950. Compound [A] when reacted with PCI5 and
nhosphoric acid		then with ammonia gave (B).
(b) benzoic anhydride in the presence of	f	Compound (B) when treated with bromine and
phosphoric acid	-	NaOH, produced (C). Compound
(c) methyl alcohol in the presence of sulfi	iric	(c) on treatment with NaN02/HCl at 0°C and
acid		then boiling with H20 produced ocresol.
(d) phenol in the presence of sulfuric act	id	Compoind (A) is
Answer (a)		[o]o-Toluic acid
944 Oil of Wintergreen (Methyl salicy)	ate) is	[b] o-Chlorotoluene
obtained by treating salicylic acid		(c) m-Toluic acid
with		(d) o-Bromotoluene
(a) Methane at 150°C		Answer (a)
(b) $CH30H$ and cone $H250A$		951 The reason why materials annear colored is
(c) Mothyl chlorida		(a) The selective absorption of spectral colors
(d) NaOH and then CH2OH		(h) The interaction between the light and the
$\left( u \right) $ NaOTI and then CHOON Answer $\int h$		electrons of the dve molecules
945 Which of the following statements	is true ·	(c) The composition of the white light from
, is, which of the following statements	10 11 40 1	various spectral colors

Teekay	Page 76	Educational Consult
(d) All of these		(b) 1,2-Dinitronaphthalene
Answer. (d)		(c) 2-Nitronaphthalene
952. Which of the following is a chromophe	ore?	(d) 1,5-Dinitronaphthalene
(a) —N02		Answer (a)
{b] —S03 H		<b>960.</b> All carbon atoms in anthracene are
(°) — OH		(a) sp hybridized
(d) -COOH		(h) sp3 hybridized
Answer, {a}		(c) sp2 hybridized
953 Which of the following is an auxochro	me?	(d) None of these
(A) $-N=0$	inc.	Answer (c)
(h) = NO2		<b>961</b> Anthracene undergoes electrophilic
(C) - N = N - C		substitution reactions mainly at
(d) - OH		(a) c-1
(u) = 011		$(a) C^{-1}$
954 The water so 1 ubility of dyes can be		(c) (c - 9)
increased by introducing		[d] C-1 and $C-2$
(a) 502Na groups		Answer (c)
(a) COOH groups		<b>062</b> Anthracono undorgoos ovidation with
$\{b\}$ COOLINE GROUPS		$\frac{902}{1200}$ Antimacene undergoes oxidation with $\frac{02}{1200}$ at $\frac{500\%}{1200}$ to give
(d) All of those		(a) Reprois acid
(u) All of these		(a) Anthroquinono
Allswei. (u)	. h	(b) Antinaquinone
955. Which dyes become linked to the fibro	e by	(d) Personhanana
chemical reaction?		(d) Benzophenone
(a) Acta ayes		Allswer. (D) 062. Which of the following is not a hotorographic
(D) Direct dyes		<b>963.</b> Which of the following is not a neterocyclic
(c) Disperse dyes		compound?
(d)none of these		
Answer. [a]		
956. All carbon atoms in naphthalene are		(a) N
(a) sp hybridized		
{b) sp2 hybridized		
(c) sp3 hybriaized		
(d)None of these		$(b) \sim N$
Answer. (b)		0
957. Naphthalene undergoes oxidation with		U.
Na2Cr207 / H2504 to form		$\frown$
(a) Phthalic acid		
(b) Phenylacetic acid		(c)
(c) Tetralin		
		l y
Answer. [a]		(d) S
958. Naphthalene undergoes reduction with	th H2	
In the presence of NI catalyst at		Answer. (c)
nigh temperature and pressure to give		964. Which uf the followingheterocyclic
(a) Phthalic acid		compounds is not aromatic?
(b) Decalin		[a] Pyridine
(c)Benzoic acid		[b]Pyrrole
(d) tetralin		(c) Furan
Allswer. (D)		(a) Piperiaine
<b>959.</b> Naphthalene undergoes nitration with		Answer (d)
HNU3/H25U4 at 60°C to give mainly		<b>965.</b> The N' atom in pyridine is
(a) i-Nitronaphthalene		(a) sp3 hybridized
Impacting Lives		

Teekay	Page 77	Educational Consult
<i>(b) sp2</i> hybridized		(b) 3-Aminopyridine
(c) sp hybridized		(c) 4-Aminopyridiee
(d) cannot be predicted		(d) None of these
Answer. (b)		Answer. (a)
966. Pyridine has a delocalized n molecula	r	<b>973.</b> Which of the following reagents will react
orbital containing		with pyrrole to form 2-
(a) 4 electrons		formylpyrrole?
(b) 6 electrons		(a) HCOOH
(c) 8 electrons		(b) CIIC18/KOI I
(d) 12 electrons		(c) H202
Answer. [b]		(d) [CH3C0)20/SnC14
967. Pyrrole is less basic than pyridine bed	ause	Answer. (b)
the lone-pair of electrons on Natom		974. Furan reacts with ammonia in the presence
in pyrrole		of alumina at 400°C to give
(a) is part of the delocalized o molecular or	bital.	(a) Pyridine
(b) is not part of the delocalized z molecular		(b) Furfural
orbital.		(c) Pyrrole
(c) resides in <i>sp2</i> hybrid orbital		(d) Furoic acid
(d) resides ;n sp hybrid orbital		Answer. (c)
Answer. ia1		<b>975.</b> Which of the following reagents will react
968. Pyridine is less basic than trimethylar	nine	with furan to form 2-furansu1fonic
because the lone-pair of electrons		acid2
on N-atom in pyridine resides in		(a) SO3 in pyridine at 100°C
[a j sp2 hybrid orbital		<i>[b]</i> Dilute H2S04 at 200°C
tb) <i>sp</i> hybrid orbital		(c) 502 at 100°C
(c) sp3 hybrid orbital		(d) Dilute H2S04 at 100°C
(d) p-orbital		Answer. (a)
Answer. (a)		976, When aniline is treated with glycerol in the
<b>969.</b> Pyridine reacts with HCI to form		presence of sulfuric acid and
(a) Pyridinium chloride		nitrobenzene, it gives quinoline. This reaction in
(b) 2-Ch1oropyridine		called
(c) 3-Chloropyridine		(a) Fischer synthesis
(d) All of these		(b) Skraup synthesis
Answer. [a]		(c) Diazotization
<b>970.</b> Pyridine undergoes electrophilic		(d} Corey-house synthesis
substitution with fuming H2SO4 at 3S0°C t	0	Answer. {b)
give		<b>977.</b> Quinoline undergoes nucleophilic
(a) 2-Pyridinesulfonic acid		substitution on heating with NaNH2 to give
<i>{b ]</i> 4-Pyridinesu1fonic acid		(a) 2-Aminoquinoline
(c) 3-Pyridinesulfonic acid		(b) 4-Arninoquirtoline
(d) None of these		(c) 3-Aminoquinoline
Answer. (c)		(d) 8-Aininoquinoline
<b>971.</b> Pyridine reacts with a mixture of KN(	)3 and	Answer. (a)
H2S04 at 800°C to give		978 Orton is prepared by the polymerization of
(a) I-Nitropyridine		(a) Vinyl cyanide
<i>(b)</i> 2-Nitropyridine		(b) Allyl alcohol
(c) 3-Nitropyridine		[c) Vinyl chloride
(a) 4-Nitropyridine		(d) Allyl chloride
Answer. (C) 072 Dumiding underge og musika sukilis		Answer. (a)
<b>972.</b> Pyriaine undergoes nucleophilic		<b>9/9.</b> which of the following statements is true
substitution with NanH2 at 100°C to form		about terpenes?
(a) 2-Minnopyriunie		

Impacting Lives

Teekay	Page78	Educational Consult
(a) They are a class of lipids that can be is	olated	Answer. (a)
from plants.		<b>987.</b> Which of the following is a thermosetting
<i>{b]</i> They contain carbon atoms in multipl	es of	polymer?
five.		(a) Bakelite
(c) They are composed of isoprene units j	oined	<i>[b]</i> Nylon-6,6
together in a head-to-tail fashion.		(c) Polyethylene
(d) All of the above		(dJ Tcflon
Answer. (d)		Answer. (a)
<b>980.</b> Ihow many isoprene units are in a		<b>988.</b> Which of the following is an example of a
sesquiterpene?		condensation polymer?
(A] 1		[a] Nylon-6,6
(b) 2		(b) Teflon
(c) 3		(c] Polypropylene
(d) 4		(d) Orton
Answer. (c)		Answer. (a)
<b>981.</b> What is the IUPAC name for isoprene	?	989. Which of the following polymers contain
(a) 1,3-pentadiene		nitrogen
<i>{b)</i> 2,4-pentadiene		(a) PVC
(c) 2,3-dimcthyl-1, 3-butadiene		(b) Teflon
(d) 2-methyl-1,3-butadiene		(c) Nylon
Answer. (d)		(d) Terylene
982. How many isoprene units are presen	lt In	Allswer. (c)
(a) 1		diaming to form
$\begin{bmatrix} a \end{bmatrix} 1$ (b) 2		(a) Bakolita
(c) 2		[h] Nylon-6 6
(d) 4		(c) Tervlene
Answer. [c]		(d) Nylon-6.8
<b>983.</b> Teflon is prepared by the polymeriz	ation of	Answer. (b)
(a) Butadiene		<b>991.</b> Ethylene glycol reacts with dimethyl
<i>(b)</i> Vinyl cyanide		terephthalate to form
(c) Vinyl chloride		(a) Nylon-6,6
(d) Tetrafluoroethylene		(b) Teflon
Answer. (d)		(c) Dacron
984. 8akelite is obtained from :		(d) Orton
(a) Phenol and formaldehyde		Answer. (c)
(b) Adipic acid and hexamethylene diamin	ne	992. Natural rubber is a polymer of
(c) Dimethyl terephthalate and ethylene	glycol	(a) Propene
(d) Neoprene		[b] Isoprene
Answer. (a)		(c) Formaldehyde
<b>985.</b> Nylon-6,6 is obtained from :		(d) Phenol
(a) Adipic acid and hexamethylene diamin	ne	<b>Answer</b> . {b)
<i>{b)</i> I etrafluoroethylene		<b>993.</b> The monomers for <i>Buna-S</i> are 1,3-
(c) Vinyl cyanide		butadiene and
(u) vinyibenzene		(a) Eurylene grycor
Allswei. (u) 996 Noopropo is a polymor of the followi	na	(b) Aupic aciu
monomer	118	(d) Canrolactum
(a) Chloroprene		Answer (c)
(h) Isoprene		994. Which of the following statements is not
(c) Isobutane		true?
(d) lsopentene		(a) Natural rubber is a hydrocarbon.
Impacting Lives	I	

(a) Carboxylic acid

(b) Aldehyde

(c) alcohol

(d) ketones

ANSWER (a)

*(b)*Natural rubber is made of isoprene units (c) Natural rubber is a polymer of I,3-Butadiene (d) Natural rubber can be vulcanized. Answer. (c) 995. Which of the following are addition polymers? 1. Polypropylcne 11. Teflon III. Nvlon (a) Only l *{b*) Only II (c) Only III (d) both I and II Answer. (d) 996- Which of the following are condensation polymers? 1. Polypropylene 11. Teflon III. Nylon (a) Only I *{b*) Only II (c) Only III (d) Both II and III Answer. (c) 997. Which one of the following is used to make Teflon? (a) Fluoroethene (b) 1,1,4,4-tctrafluorohutadienc (c) 1,2-difluoroethene (d) Tetrafluoroethylene Answer. (d) 998. Vulcanization is the process of crosslinking polymer chains in rubber using (a) Sulfur. *{b]* Formaldehyde, (c) Benzoyl peroxide. (d) Ethylene glycol. **Answer**. (*a*) **999.** What is the purpose of plasticizers? (a) Harden plastics (b) Soften plastics (c) Initiate polymerizations (d) Cross-link polymer chains **Answer**. [b] 1000. Polymers which soften on heating and harden when cooled are (a) Cross-linked polymers *{b*) Copolymers (c) Thermosetting polymers (d) Thermoplastics Answer. (d) 1001. Oxidation of Primary alcohol gives

Impacting Lives

