

LAUTECH
nano⁺ 2018

NANOTECHNOLOGY RESEARCH GROUP

...Enabling Qualitative Research at the Smallest Scale



Theme: NANOTECHNOLOGY FOR SUSTAINABLE DEVELOPMENT: Prospects for Africa

VENUE: The Hall & Laboratory of Industrial Microbiology and Nanobiotechnology, LAUTECH, Ogbomoso
DATE: 23 - 25 OCTOBER, 2018 TIME: 9:00 AMwww.nanotech.lautech.edu.ng

Binomial theory can be used when probability of success is very high

Binomial Formula $b(x, n, p) = {}^nC_x \times p^x \times (1-p)^{n-x}$
 $n! / (x! \times (n-x)!)$

b = binomial probability

x = total number of success

p = probability of a success in an individual trial

n = number of trials

$${}^nC_x = \frac{n!}{(n-x)!}$$

A coin is tossed 10 times what is probability of getting 6 heads

Probability of success $p = 0.5$ Probability of failure $1 - 0.5 = 0.5$ $n = 10$ $x = 6$

$$P_6 = {}^nC_x \times p^x \times (1-p)^{n-x}$$

$$P_6 = {}^{10}C_6 \times p^{0.5} \times (1-0.5)^{10-6}$$

$$= \frac{10!}{(10-6)! \times 6!} \times 0.5^6 \times 0.5^4$$

$$\frac{10!}{4! \times 6!} \times 0.5^6 \times 0.5^4 \quad \left| \begin{array}{l} \text{mean} = np = 10 \times 0.5 = 5 \\ \text{Variance} = npq = 10 \times 0.5 \times 0.5 = 2.5 \\ \text{Standard deviation} = \sqrt{npq} = \sqrt{10 \times 0.5 \times 0.5} \end{array} \right.$$

POISSON DISTRIBUTION

occur when we have a rare event when probability of success is very low

$$P(X, \mu) = \frac{[e^{-\mu}] [\mu^x]}{x!}$$

where X = actual number of success that can result from

$$e = 2.71828$$

μ = mean of population

Properties

mean = μ

Variance = μ

Standard deviation = $\sqrt{\mu}$

Average no of homes sold by Solomon is 2 homes per day what is the probability of selling 3 homes tomorrow

Solution

$$\mu = 2$$

$$x = 3$$

$$e =$$

$$P(X, \mu) = \frac{(e^{-\mu}) (\mu^x)}{x!} = \frac{e^{-2} \times 2^3}{3!}$$

Theme: NANOTECHNOLOGY FOR SUSTAINABLE DEVELOPMENT: Prospects for Africa

VENUE: The Hall & Laboratory of Industrial Microbiology and Nanobiotechnology, LAUTECH, Ogbomosho

DATE: 23 - 25 OCTOBER, 2018 TIME: 9:00 AM

www.nanotech.lautech.edu.ng

HYPERGEOMETRIC DISTRIBUTION

IF a population of size N contains K items of successes will have $[N-K]$ failures, then the probability of hypergeometric random variables X ,

The no of success in a random sample n

$$P(X=K) = \frac{\binom{K}{k} \binom{N-K}{n-k}}{\binom{N}{n}} = \frac{\binom{K}{k} \binom{N-K}{n-k}}{\binom{N}{n}}$$

N = size from Population

n = size from sample

K = success from population

k = success from sample.

A deck of cards contain 20 cards, 6 red cards and 14 black cards. 5 cards are drawn randomly without replacement probability of exact 4 red cards are drawn

Solution

Exactly = 4

$$N - K = 20 - 6 = 14$$


$N = 20$

$$n - k = 5 - 1 = 4$$

$n = 5$

$$\frac{\binom{6}{4} \binom{14}{1}}{\binom{20}{5}} = \frac{\binom{6}{4} \binom{14}{1}}{\binom{20}{5}}$$

NORMAL DISTRIBUTION



$$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

$Z =$ Size

$\bar{x} =$ mean of Sample

$\mu =$ mean of Population

$\sigma =$ Standard deviation of Population

$n =$ Size of Sample

ABE 301

Course Content

Probability & Statistics.

Probability space, theories, Conditional probability and independency, random variables, discrete and continuous distributions.

Mean and variance, Binomial, Poisson hyper geometry.

Exponential and normal distribution and their characteristic

Center limit theorem.

Elementary Sample theorem for normal population Statistical inference on mean proportion and Variance.

- Simple linear regression & Engineering application
- Chi-Square.

Probability & Statistics. Likelihood.

Probability is the measure of the ~~likelyhood~~ that a particular event will occur in any one trial or experiment carried out in a pre-scribed condition

Notation

P - Probability

The probability that a certain event of A will occur is denoted by $P(A)$ or $p(A)$. It is denoted by $(P \text{ Success})$

Success or Failure.

When an event occurs in any one trial, it's called Success but when it fails it's called failure. In N ^{of} any trials there are x (successes) there will ~~also~~ be $(N-x)$ failure.

$$\frac{x}{N} = P(\text{Success})$$

$$\frac{N-x}{N} = P(\text{Failure}) \rightarrow \left(\frac{N-x}{N} \right)$$

~~(N-x)~~

$$\frac{2}{120} = \frac{120-2}{120}$$

$$P(A) + P(\text{Not } A) = 1$$

$$P(A) + P(\bar{A}) = 1$$

$$\frac{x}{N} + \frac{N-x}{N} = 1$$

Type of probability

Determination of Probability may be undertaken from two processes:-

- 1) Empirical or Experimental.
- 2) Classical or Theoretical.

Empirical or Experimental.

It's based on the ^{previous} known result.

→ Expectation. It's defined as the product of No. of trials and probability that event A will occur in a number of trials.

Classical

The classical approach to probability is based on application of theoretical number of ways in which it's possible for an event to occur.

Classical probability (P) of an event (A) occurring is defined by

$$P(A) = \frac{\text{Number of ways in which event A occurs}}{\text{Total Number of all possible outcomes}}$$

2	2	2	2	2	2
1	2	3	4	5	6
1	1	1	1	1	1

$$= \frac{2}{6}$$

Mutually Exclusive and Mutually Non-Exclusive Event

Mutually Exclusive Event - This cannot occur together...

Mutually Non-Exclusive Event's are those which happen simultaneously eg a pair of events like 2 dice casted.

or - Add
both - Multiplication

Additional Law of Probability

For $\rightarrow$ Given event (A) and (B) are Mutually Exclusive
Since either

IF A and B are Mutually Exclusive, Probability will be either A or B but not both.

$$P(A \text{ or } B) = P(A) + P(B)$$

Independent Event and Dependent Event $\rightarrow$ (Check)

When the occurrence of one event doesn't affect the probability of occurrence of the second event $\rightarrow$ Independent event (Replaced)

Event are dependent when the occurrence of one event affect the probability of occurrence of the second event (Not replaced)

Oranges

red orange = 12

White orange = 8

Blue orange = 10

Independent

dependent $\Rightarrow$

White $\frac{2}{8}$

$$\frac{2}{30} \rightarrow A \quad \frac{10}{30} \rightarrow B$$

Conditional probability of an event ^(B) given occurring when given an event (A) has already taken place
 vice versa. ^{but} In case of an event ^(B) stated above

$$P(B/A) = \frac{5/30}{2/30}$$



$$\left. \begin{aligned} P(\text{picking 2 white orange}) &= \frac{2}{30} \text{ (A)} \\ P(\text{picking 5 blue orange}) &= \frac{5}{30} \text{ (B)} \end{aligned} \right\} \text{Independent.}$$

$$P(\text{picking 5 red orange}) = \frac{5}{28} \text{ } \left. \vphantom{\frac{5}{28}} \right\} \text{dependent.}$$

Binomial Distribution.

Binomial theorem can be used when the probability of Success is very high.

Binomial formula.

$$b(x; n, p) = {}^n C_x \times p^x \times (1-p)^{n-x}.$$

Where

b = Binomial Probability.

x = total No. of successes

p = Probability of success in an individual trial.

n = Number of trials.

$${}^n C_x = \frac{n!}{(n-x)! x!}$$

Example: A coin is tossed 10 times what is the probability of getting exactly 6x?

$$p = 0.5$$

$$q = 1 - 0.5 = 0.5$$

$$n = 10$$

$$x = 6$$

Not less than 6 - 6, 7, 8, 9, 10
 Not more than 6 - 6, 5, 4, 3, 2, 1, 0.

$$P_6 = {}^nC_x \cdot p^x \cdot (1-p)^{n-x}$$

$$= {}^{10}C_6 \times p^6 \times (1-0.5)^{10-6}$$

$$= \frac{10!}{(10-6)! 6!} \times 0.5^6 \times 0.5^4$$

$$= \frac{10!}{4! 6!} \times 0.5^6 \times 0.5^4$$

$$\text{Mean} = nP = 10 \times 0.5 = 5$$

$$\text{Variance} = nPq = 10 \times 0.5 \times 0.5 = 2.5$$

$$\text{Standard deviation} = \sqrt{nPq}$$

Poisson Distribution

occurs When we have a rare event, the probability of Success is very low.

$$P(x; \mu) = \frac{e^{-\mu} (\mu^x)}{x!}$$

Where

x = actual number of successes that result from experiment.

$\bar{x}$ = mean^s of sample.

s = standard deviation of sample.

σ = standard deviation of population

μ = mean of population.

$$e = 2.71828$$

Properties.

$$\text{mean} = \mu$$

$$\text{Variance} = \mu$$

$$\text{Standard deviation} = \sqrt{\mu}$$

Example

Average number of homes sold by Solomon Company is 2 homes per day. What is the probability that exactly three (3) homes would be sold tomorrow.

Solution

$\mu = 2$, Since 2 homes are sold per day.

$$x = 3$$

$$e = ?$$

$$P(x; \mu) = \frac{(e^{-\mu}) (\mu^x)}{x!} = \frac{e^{-2} \times 2^3}{3!}$$

$$\text{Ans} = 0.180$$

A num ^{Success.} of female from a Population (male & female)

Hyper Geometric

When one takes a certain number from a sample drawn from a population.

If a Population of size "N" contains 'k' items of successes, failure will be (N-k), then the probability of the hypergeometric random variable (X), the no. of successes in a random sample of size (n) is

$$P(X=k) = \frac{\binom{K}{k} \binom{N-k}{n-k}}{\binom{N}{n}}$$

$$= \frac{\binom{K}{k} \binom{N-k}{n-k}}{\binom{N}{n}}$$

K - Population. (Success)

k = Sample. (Success)

Examples

A deck of cards containing 20 cards, 6 red cards and 14 black cards. 5 cards are drawn at random without replacement. What is the probability that exactly 4 red cards are drawn?

Solution

$$N = 20$$

$$n = 5$$

$$N - K = 20 - 6 = 14$$



$$n - k = 5 - 4 = 1$$

$$= \frac{\binom{6}{4} \binom{14}{1}}{\binom{20}{5}}$$

$$= \frac{\binom{6}{4} \binom{14}{1}}{\binom{20}{5}}$$

Normal Distribution

Approach a given point without touching it e.g. Sin 90.




$Z = \text{Size}$

$\bar{x} = \text{mean of sample}$

$\sigma = \text{standard deviation of Population}$

$n = \text{size of sample}$

$\mu = \text{mean of population}$



Normal distribution table

$$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

	Probability →
Normal size ↓	