

- 15.7** Find the Laplace transforms of the following functions:

- $2\delta(3t) + 6u(2t) + 4e^{-2t} - 10e^{-3t}$
- $te^{-t}u(t-1)$
- $\cos 2(t-1)u(t-1)$
- $\sin 4t[u(t) - u(t-\pi)]$

- 15.8** Determine the Laplace transforms of these functions:

- $f(t) = (t-4)u(t-2)$
- $g(t) = 2e^{-4t}u(t-1)$
- $h(t) = 5 \cos(2t-1)u(t)$
- $p(t) = 6[u(t-2) - u(t-4)]$

- 15.9** In two different ways, find the Laplace transform of

$$g(t) = \frac{d}{dt}(te^{-t} \cos t)$$

- 15.10** Find  $F(s)$  if:

- $f(t) = 6e^{-t} \cosh 2t$
- $f(t) = 3te^{-2t} \sinh 4t$
- $f(t) = 8e^{-3t} \cosh tu(t-2)$

- 15.11** Calculate the Laplace transform of the function in Fig. 15.49.




Figure 15.49 For Prob. 15.11.

- 15.12** Find the Laplace transform of the function in Fig. 15.50.




Figure 15.50 For Prob. 15.12.

- 15.13** Obtain the Laplace transform of  $f(t)$  in Fig. 15.51.




Figure 15.51 For Prob. 15.13.

- 15.14** Determine the Laplace transforms of the function in Fig. 15.52.




Figure 15.52 For Prob. 15.14.

- 15.15** Obtain the Laplace transforms of the functions in Fig. 15.53.




Figure 15.53 For Prob. 15.15.

- 15.16** Calculate the Laplace transform of the train of unit impulses in Fig. 15.54.




Figure 15.54 For Prob. 15.16.

**15.27** Calculate the inverse Laplace transform of:

$$(a) \frac{6(s-1)}{s^4-1} \quad (b) \frac{se^{-\pi s}}{s^2+1} \quad (c) \frac{8}{s(s+1)^3}$$

**15.28** Find the time functions that have the following Laplace transforms:

$$(a) F(s) = 10 + \frac{s^2 + 1}{s^2 + 4}$$

$$(b) G(s) = \frac{e^{-s} + 4e^{-2s}}{s^2 + 6s + 8}$$

$$(c) H(s) = \frac{(s+1)e^{-2s}}{s(s+3)(s+4)}$$

**15.29** Obtain  $f(t)$  for the following transforms:

$$(a) F(s) = \frac{(s+3)e^{-6s}}{(s+1)(s+2)}$$

$$(b) F(s) = \frac{4 - e^{-2s}}{s^2 + 5s + 4}$$

$$(c) F(s) = \frac{se^{-s}}{(s+3)(s^2 + 4)}$$

**15.30** Obtain the inverse Laplace transforms of the following functions:

$$(a) X(s) = \frac{1}{s^2(s+2)(s+3)}$$

$$(b) Y(s) = \frac{1}{s(s+1)^2}$$

$$(c) Z(s) = \frac{1}{s(s+1)(s^2 + 6s + 10)}$$

**15.31** Obtain the inverse Laplace transforms of these functions:

$$(a) \frac{12e^{-2s}}{s(s^2+4)}$$

$$(b) \frac{2s+1}{(s^2+1)(s^2+9)}$$

$$(c) \frac{9s^2}{(s^2+4s+13)}$$

**15.32** Find  $f(t)$  given that:

$$(a) F(s) = \frac{s^2 + 4s}{s^2 + 10s + 26}$$

$$(b) F(s) = \frac{5s^2 + 7s + 29}{s(s^2 + 4s + 29)}$$

\***15.33** Determine  $f(t)$  if:

$$(a) F(s) = \frac{2s^3 + 4s^2 + 1}{(s^2 + 2s + 17)(s^2 + 4s + 20)}$$

$$(b) F(s) = \frac{s^2 + 4}{(s^2 + 9)(s^2 + 6s + 3)}$$

## Section 15.5 Application to Circuits

**15.34** Determine  $i(t)$  in the circuit of Fig. 15.59 by means of the Laplace transform.




Figure 15.59 For Prob. 15.34.

**15.35** Find  $v_o(t)$  in the circuit in Fig. 15.60.




Figure 15.60 For Prob. 15.35.

**15.36** Find the input impedance  $Z_{in}(s)$  of each of the circuits in Fig. 15.61.




Figure 15.61 For Prob. 15.36.

**15.37** Obtain the mesh currents in the circuit of Fig. 15.62.




Figure 15.62 For Prob. 15.37.

**15.38** Find  $v_o(t)$  in the circuit in Fig. 15.63.




Figure 15.63 For Prob. 15.38.

- 16.2** Determine the period of these periodic functions:
- $f_1(t) = 4 \sin 5t + 3 \sin 6t$
  - $f_2(t) = 12 + 5 \cos 2t + 2 \cos(4t + 45^\circ)$
  - $f_3(t) = 4 \sin^2 600\pi t$
  - $f_4(t) = e^{j10t}$

- 16.3** Give the Fourier coefficients  $a_0$ ,  $a_n$ , and  $b_n$  of the waveform in Fig. 16.47. Plot the amplitude and phase spectra.




Figure 16.47 For Prob. 16.3.

- 16.4** Find the Fourier series expansion of the backward sawtooth waveform of Fig. 16.48. Obtain the amplitude and phase spectra.




Figure 16.48 For Probs. 16.4 and 16.50.

- \***16.5** A voltage source has a periodic waveform defined over its period as

$$v(t) = t(2\pi - t) \text{ V}, \quad 0 < t < 2\pi$$

Find the Fourier series for this voltage.

- 16.6** A periodic function is defined over its period as

$$h(t) = \begin{cases} 10 \sin t, & 0 < t < \pi \\ 20 \sin(t - \pi), & \pi < t < 2\pi \end{cases}$$

Find the Fourier series of  $h(t)$ .

- 16.7** Find the quadrature (cosine and sine) form of the Fourier series

$$f(t) = 2 + \sum_{n=1}^{\infty} \frac{10}{n^3 + 1} \cos \left( 2nt + \frac{n\pi}{4} \right)$$

- 16.8** Express the Fourier series


$$f(t) = 10 + \sum_{n=1}^{\infty} \frac{4}{n^2 + 1} \cos 10nt + \frac{1}{n^3} \sin 10nt$$

- in a cosine and angle form,
- in a sine and angle form.


- 16.9** The waveform in Fig. 16.49(a) has the following Fourier series:

$$v_1(t) = \frac{1}{2} - \frac{4}{\pi^2} \left( \cos \pi t + \frac{1}{9} \cos 3\pi t + \frac{1}{25} \cos 5\pi t + \dots \right) \text{ V}$$

Obtain the Fourier series of  $v_2(t)$  in Fig. 16.49(b).



(a)



(b)


Figure 16.49 For Probs. 16.9 and 16.52.

### Section 16.3 Symmetry Considerations


- 16.10** Determine if these functions are even, odd, or neither.

- $1 + t$
- $t^2 - 1$
- $\cos n\pi t \sin n\pi t$
- $\sin^2 \pi t$
- $e^{-t}$

- 16.11** Determine the fundamental frequency and specify the type of symmetry present in the functions in Fig. 16.50.




(a)




(b)

\*An asterisk indicates a challenging problem.




**Figure 16.50** For Probs. 16.11 and 16.48.

- 16.12** Obtain the Fourier series expansion of the function in Fig. 16.51.




**Figure 16.51** For Prob. 16.12.

- 16.13** Find the Fourier series for the signal in Fig. 16.52. Evaluate  $f(t)$  at  $t = 2$  using the first three nonzero harmonics.




**Figure 16.52** For Probs. 16.13 and 16.51.

- 16.14** Determine the trigonometric Fourier series of the signal in Fig. 16.53.




**Figure 16.53** For Prob. 16.14.

- 16.15** Calculate the Fourier coefficients for the function in Fig. 16.54.



**Figure 16.54** For Prob. 16.15.


- 16.16** Find the Fourier series of the function shown in Fig. 16.55.



**Figure 16.55** For Prob. 16.16.

- 16.17** In the periodic function of Fig. 16.56,
- find the trigonometric Fourier series coefficients  $a_2$  and  $b_2$ ,
  - calculate the magnitude and phase of the component of  $f(t)$  that has  $\omega_n = 10 \text{ rad/s}$ ,
  - use the first four nonzero terms to estimate  $f(\pi/2)$ ,
  - show that

$$\frac{\pi}{4} = \frac{1}{1} - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \dots$$



**Figure 16.56** For Prob. 16.17.

- 16.18** Determine the Fourier series representation of the function in Fig. 16.57.




Figure 16.57 For Prob. 16.18.

- 16.19** Find the Fourier series representation of the signal shown in Fig. 16.58.




Figure 16.58 For Prob. 16.19.

- 16.20** For the waveform shown in Fig. 16.59 below,
- specify the type of symmetry it has,
  - calculate  $a_3$  and  $b_3$ ,
  - find the rms value using the first five nonzero harmonics.

- 16.21** Obtain the trigonometric Fourier series for the voltage waveform shown in Fig. 16.60.




Figure 16.60 For Prob. 16.21.

- 16.22** Determine the Fourier series expansion of the sawtooth function in Fig. 16.61.




Figure 16.61 For Prob. 16.22.

## Section 16.4 Circuit Applications

- 16.23** Find  $i(t)$  in the circuit of Fig. 16.62 given that

$$i_s(t) = 1 + \sum_{n=1}^{\infty} \frac{1}{n^2} \cos 3nt \text{ A}$$




Figure 16.62 For Prob. 16.23.

- 16.24** Obtain  $v_o(t)$  in the network of Fig. 16.63 if

$$v(t) = \sum_{n=1}^{\infty} \frac{10}{n^2} \cos \left( nt + \frac{n\pi}{4} \right) \text{ V}$$




Figure 16.63 For Prob. 16.24.




Figure 16.59 For Prob. 16.20.