

Statistics, the exploration and Analysis of Data

- Roxy Peck & Jay Devore
Schaum's Series.

ABE 302: Engineering Statistics (3)

Course Content

(1) Probability & Statistics

- Probability Space theory
- ~~Probability~~
- Conditional probability & independence
- Random variables, discrete and continuous distributions
- Mean and Variance
- Binomial, Poisson, hypergeometry
- Exponential and Normal distributions and their characteristics
- Centre limit theory
- Elementary Sample theory for normal population.
- Statistical Inference on mean and Variance
- Simple linear regression and general applications
- Chi-Square test

Probability & Statistics

Probability is the measure of the likelihood that a particular event will occur in any one trial or experiment carried out in a prescribed condition.

Notations

The probability that a certain event A will occur is denoted by $P(A)$ or $p(A)$. It is also denoted by $(P \text{ success})$.

Success or Failure

When an event occurs in any one trial, it is called Success, but when it fails to occur, it is called failure.

In n ^{and} ~~many~~ trials, there are x successes, there will also be $(N-x)$ failures

$$P(\text{success}) = \frac{x}{N}$$

$$P(\text{failure}) = \frac{(N-x)}{N}$$

$$P(A) + P(\text{Not } A) = 1$$

$$\text{or } P(A) + P(\bar{A}) = 1$$

$$\text{Therefore, } \frac{x}{N} + \frac{N-x}{N} = 1$$

Types of Probabilities :

Determination of probability can be undertaken from two perspectives.

- (1) Empirical or Experimental
- (2) Classical or Theoretical

(1) Empirical is based on the previous known result

Expectation

This is defined as the product of the number of trials and probability that event A will occur in a number of trials

(2) The classical approach to probability is

based on the application of theoretical ^{number} ~~known~~ of ways in which it is possible for an event to occur.

Classical probability P of an event (A) occurring is defined by

$$P(A) = \frac{\text{number of ways in which event A occurs}}{\text{Total number of all possible outcomes}}$$

Mutually Exclusive and Mutually Non-Exclusive Events

Mutually exclusive events are events which cannot occur together

Mutually non-exclusive events are those which happen simultaneously, e.g. a pair of events like 2 dice casted.

Additional Law of probabilities

For a given event, A and B are mutually exclusive if either A or B occurs, but

or = Addition
but = Multiplication

not both.

$$P(A \text{ or } B) = P(A) + P(B)$$

Independent Events and Dependent Events

Events are independent when the occurrence of one event does not affect the probability of occurrence of the second event.

Events are dependent when one event just affects the probability of occurrence of a second event.

Independent — Replacement of events
Dependent — Non-replacement.

red orange = 12

blue orange = 10

white orange = 8

~~white orange~~

$$\left. \begin{aligned} P(\text{picking 2 white orange}) &= \frac{2}{30} \text{ (A)} \\ P(\text{picking 5 blue orange}) &= \frac{5}{30} \text{ (B)} \end{aligned} \right\} \text{Independent}$$
$$P(\text{picking 5 red orange}) = \frac{5}{28} \text{ } \left. \right\} \text{Dependent}$$

Conditional probability

The conditional probability of an event B occurring when given an event A has already taken place vice-versa

In case of the event B above

$$P\left(\frac{B}{A}\right) = \frac{5/30}{2/30}$$

19-05-2021

Binomial Distribution

This can be used when the probability of

success of an event is very high.

The Binomial Formula = $b(x; n, p) = {}^nC_x \cdot p^x \cdot (1-p)^{n-x}$

$$b(x; n, p) = {}^nC_x \cdot p^x \cdot (1-p)^{n-x}$$

where b = binomial probability

x = total number of successes

n = total number of trials.

n = number of trials.

Example : A coin is tossed 10 times.

What is the probability of getting exactly 6 heads

$$P = 0.5$$

$$q = 1 - 0.5$$

$$n = 10$$

$$x = 6$$

$$P_6 = {}^{10}C_6 \cdot p^6 \cdot (1-p)^{10-6}$$

$$= {}^{10}C_6 \cdot 0.5^6 \cdot (1-0.5)^{10-6}$$

$$= \frac{10!}{(10-6)!6!} \times 0.5^6 \times 0.5^4$$

$$= 210 \times \frac{1}{64} \times \frac{1}{16}$$

$$P_6 = 0.2051$$

μ - population mean

$\bar{x}$ - Sample mean

σ - SD of sample

δ = SD of population

$$\text{Mean} = np$$

$$\text{Variance} = npq$$

$$SD = \sqrt{npq}$$

When we have a rare event, the probability is very low

Poisson Distribution

$$P(x; \mu) = \frac{e^{-\mu} \mu^x}{x!}$$

where x = Actual number of successes that result from a no. of trials

$$\mu = 2.71828$$

μ = mean of population.

Properties of Poisson Distribution

$$\text{Mean} = \mu$$

$$\text{Variance} = \mu$$

$$SD = \sqrt{\mu}$$

Example

The average number of homes sold by a company is 2 homes per day. What is

the probability that exactly 3 homes will be sold tomorrow?

Solution

$\mu = 2$ since 2 homes are sold on average per day
 $X = 3$

$$P(x: \mu) = \frac{e^{-\mu} (\mu^x)}{x!}$$

$$P(x: \mu) = \frac{(e^{-2}) (2^3)}{3!}$$

$$P(x: \mu) = \frac{1.083}{6} = \underline{\underline{0.180}}$$

Hypergeometric Distribution.

When taking an exact number from a

sample of a population.

If a population of size "N" contains "K" items of success, failure = $(N - K)$,

then the probability of the hypergeometric random variable (X),
The number of successes in a random

sample size (n) is

$$P(X = k) = \frac{\binom{K}{k} \binom{N-K}{n-k}}{\binom{N}{n}}$$

$$P(X = k) = \frac{{}^K C_k \cdot {}^{(N-K)} C_{(n-k)}}{{}^N C_n}$$

K = success picked from population
k = success picked from sample

Example.

A deck of cards contains 20 cards - 6 red cards and 14 black cards. 5 cards are drawn at random, without replacement. What is the probability that exactly 4 red cards are drawn?

Solution

Exactly = 4 $\Rightarrow k$

$N = 20$

$k = 6$

$n = 5$

$N - k = 20 - 6 = 14$

$n - k = 5 - 4 = 1$


$$\frac{\binom{6}{4} \binom{14}{1}}{\binom{20}{5}}$$

$$= \frac{\binom{6}{4} \binom{14}{1}}{20C5}$$

$$= \frac{15 \times 14}{15504}$$

$$= 0.0135$$

Normal Distribution



$$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$


$Z = \text{size}$

$\bar{x}$ = Mean of sample

μ = mean of population

σ = Standard deviation of population

n = size of sample



non-parametric tests - based on observations

26-05-2021

Chi-Square (χ^2)

The chi-square test is a type of non-parametric test

$$\chi^2 = (\text{Observation} - \text{Expected})^2$$

$$\chi^2 = \frac{(O - E)^2}{E}$$

	Expected	
	E	E
ABE	5	5
CVE	6	5
EFE	7	5

Level of significance = 0.05

$$\sum \chi^2 = \frac{(5-5)^2}{5} + \frac{(6-5)^2}{5} + \frac{(7-5)^2}{5}$$

Degree of freedom = n - 1 = 3 - 1 = 2

no. of samples n = 3

$$0 + \frac{1^2}{5} + \frac{2^2}{5}$$

$$= 0 + 0.2 + 0.8 = 1$$

$$\chi^2 = 1$$

Observation

	Small	Medium	Large	Σ
White	10	12	8	30
Red	5	13	10	28
Black	2	5	12	19
Σ	17	30	30	77

Expected

	Small	Medium	Large
White	$\frac{30 \times 17}{77}$	$\frac{30 \times 30}{77}$	$\frac{30 \times 30}{77}$
Red	$\frac{28 \times 17}{77}$	$\frac{28 \times 30}{77}$	$\frac{28 \times 30}{77}$
Black	$\frac{19 \times 17}{77}$	$\frac{19 \times 30}{77}$	$\frac{19 \times 30}{77}$

White 6.62 11.68 11.68

Red 6.18 10.91 10.91

Black 4.19 7.40 7.40

7 12 12

6 11 11

4 7 7

$$\chi^2 = \frac{\sum (O - E)^2}{E} = \frac{(10-7)^2}{7} + \frac{(12-12)^2}{12} + \frac{(8-12)^2}{12}$$

+ ...



White, Red, Black
Small, Medium, Big

$$\text{degree of freedom} = (3-1)(3-1) = 4$$

Measures of Association

The study of association has to do with bivariate (2 variables), whereby 2 or more variables are considered as a means of viewing the relationship existing between them. When an increase in variable leads to an increase in another variable, we have a positive association of the relations. However when an increase in a variable leads to a decrease in the other, then we have a negative association.

Types of Statistical Relationship



Karl Pearson's Product - Moment Correlation Coefficient.

This is the most popular mathematical method of quantifying or measuring correlation. It is always denoted by the symbol r , and is given by

$$r = \frac{\text{Cov}(x, y)}{\sqrt{\text{Var}(x) \cdot \text{Var}(y)}} = \frac{\text{Cov}(x, y)}{\sigma_x \cdot \sigma_y}$$

$$= \frac{\sum xy}{M \sigma_x \cdot \sigma_y}$$

where σ_{xy} = standard deviation of x and y respectively.

M = population size

Find the regression line

$n =$ number of sample

$$y = a + bx_1 + cx_2 \quad \text{--- (1)}$$

$$y = a_n + b \sum x_1 + c \sum x_2 \quad \text{--- (2)}$$

$$\sum x_1 y = a \sum x_1 + b \sum x_1^2 + c \sum x_1 x_2 \quad \text{--- (3)}$$

$$\sum x_2 y = a \sum x_2 + b \sum x_1 x_2 + c \sum x_2^2 \quad \text{--- (4)}$$

x_1	y	$x_1 y$	x_1^2	y^2
4	5	20	16	25
6	4	24	36	16
7	5	35	49	25
8	6	48	64	36

$$n = 4$$

$$\sum x_1 = 25$$

$$\sum y = 20$$

$$\sum x_1 y = 127$$

$$\sum x_1^2 = 165$$

$$\sum y^2 =$$

$$b = \frac{\sum (x_1 - \bar{x})(y - \bar{y})}{\sum (x_1 - \bar{x})^2}$$

$$a = \bar{y} - b\bar{x}$$

$$b = \frac{\sum x_1 y - \frac{\sum x_1 \sum y}{n}}{\sum x_1^2 - \frac{(\sum x_1)^2}{n}}$$

$$a = \frac{\sum y - \frac{\sum x_1 \sum y}{n}}{n}$$

$$r = \frac{\sum x_1 y}{\sqrt{\sum x_1^2 \sum y^2}} \quad \text{--- (1)}$$

$$r = \frac{N \sum x_1 y - \sum x_1 \sum y}{\sqrt{(N \sum x_1^2 - (\sum x_1)^2)(N \sum y^2 - (\sum y)^2)}}$$

02/04/2021

Linear Regression

It is used for prediction of dependent variable X and for testing the strength of independent variable Y .

$$Y = a + bx \quad \text{--- (1)}$$

$$\sum y = a_n + b \sum x \quad \text{--- (2)}$$

$$\sum x y = a \sum x + b \sum x^2 \quad \text{--- (3)}$$

where

$a, b =$ constants

$$y = 5 - 2x$$

when $x = 40$,
what is y ?

$$y = 8.16 \times 0.2x$$

$$\Sigma y = a_n + b \Sigma x$$

$$20 = 4a + 25b \quad \text{--- (1)}$$

$$\Sigma xy = a \Sigma x + b \Sigma x^2$$

$$127 = 25a + 165b \quad \text{--- (2)}$$

Solving Simultaneously,

$$a = 3.6$$

$$b = 0.2$$

~~Ms. Foyse~~

Analysis of Variance: One Way Test

Let us consider sample data presented below.

2, 4, 4, 3, 4, 8, 7, 8, 9, 9, 2

$$t \text{ test} = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}$$

$$\frac{s}{\sqrt{n}}$$

μ = mean of population (given data)

s = SD of sample

n = no. of sample

$$\bar{x} = \text{mean}$$

$$\text{degree of freedom} = 11 - 1 = 10$$

μ = mean of entire population



Statistics SI 2-100

t-table

female - 4, 3, 4, 3, 5, 7, 6, 9, 2, 5, 3

female - larger variance

Smaller variance

$$\frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}} \pm \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}$$

male

female

Eng. Dr. Mrs. Foyse

Statistical Inference

2 types of statistics

1. Descriptive

2. Inferential

There are 3 main underlying ideas in

inference

1. Sample is likely to be a good representation of the population, if the sample is well collected.

Types of Sampling

(1) Random Sampling

Every member of the population is given the equal chance to be selected in a situation. e.g. to determine the quality of petrol kernels for sale.

Random Sampling is used to eliminate bias, and most theorems in statistics are based on random sampling.

Disadvantage - Random Sampling may not be a good representation of the population.

(2) Stratified Sampling

The population is first divided into 2 or more groups (called strata) and random sampling is then conducted.

Advantage - ~~that~~ They are usually more representative of the population.

Sampling distribution
Confidence Interval
Hypothesis testing.

(3) Cluster Sampling

Units of population exist already in particular groups. Samples are usually representative of the population.

Disadvantage - It may be difficult to locate cluster values.

(4)

~~Systematic~~ ^{Systematic} Sampling

Elements are selected from a pre-determined interval in population frame.

Sampling/Samples collected may or may not be a good representation of population.

To determine the extent of uncertainty in a sample.

dividing the population into non-overlapping subgroups, called clusters of interest.

09-06-2024

SAMPLING DISTRIBUTION

Parameter $\rightarrow$ Statistic

A parameter is a quantity that measures or describes the population we want to study

A statistic is a portion, or part of the sample

Population mean μ

Population variance σ

Population Standard deviation

Population proportion

Sample

Sample mean $\bar{x}$

Sample variance

Sample Standard deviation

Sample proportion

17

21

21

60

$$\bar{x}_1 = \frac{60}{3} = 20$$

18

19


25

62

$$\bar{x}_2 = \frac{62}{3} = 20.6$$

Picking 3 samples from a population of 10

$$\binom{10}{3} = {}^{10}C_3 = 120$$



The more the number of samples, the more your distribution tends to a normal distribution

If $n \geq 30$, it tends to a normal distribution

For Normal distribution,

$$\text{Mean} = \frac{\sum x}{n}$$

$$\text{Variance} = \frac{\sum (x - \bar{x})^2}{n}$$

$$\text{S.D} = \sqrt{\frac{\sum (x - \bar{x})^2}{n}} = \sqrt{V}$$

For Sampling distribution

$$\text{Variance} = \frac{\sigma^2}{n} \Rightarrow \text{standard error of sampling distribution.}$$

$$\text{Proportion } (\bar{p}) = \sqrt{\frac{p(1-p)}{n}}$$

$$\sqrt{\frac{p(1-p)}{n}}$$

$$\textcircled{1} \mu_1 = \mu$$

$$\textcircled{2} \sigma_x = \frac{\sigma}{\sqrt{n}}$$

Properties of Sampling Distribution of Mean

1. For any number of sample (N) , the center of the mean distribution $(\bar{x})$ will also coincide with the mean population being sampled by the spread of the mean (S.D).
2. Distribution will decrease as N increases

Conditions for Properties of Sampling distribution of Sample mean.

$\bar{x}$ = Mean of the observation in a random

Sample of size n .

From a population having mean (μ) and standard deviation σ , the mean value of the $\bar{x}$ distribution of (μ) and the

Standard deviation of the $\bar{x}$ distribution by σ_x . The following rules hold

Rule 1 : $\mu_1 = \mu$

Rule 2 : $\sigma_x = \frac{\sigma}{\sqrt{n}}$

Rule 3

When the population distribution is normal, the sampling distribution is also normal for any sample size n

Rule 4

Central Limit Theory states that when 'n' is sufficiently large, the sampling distribution is well approximated by a normal curve even when the population distribution is not itself normal.

Sample Distribution for Sample Proportion

Parameter

Statistic

π = Population proportion

p $\Rightarrow$ Sample Proportion

$P = \frac{\text{No of successes in the sample}}{n}$

Properties of Sample Proportion in Sample Distribution

(1) The sampling distribution of p is centered at $\bar{x}$. i.e. $\mu_p = \pi$. Therefore, p is an unbiased

Statistic for estimating π

(2) The standard deviation of $\hat{p} = \sqrt{\frac{\pi(1-\pi)}{n}}$

as long as n is large and $n\pi \geq 10$

$n = \text{no of sample}$ $n(1-\pi) \geq 10$
 $\pi = \text{proportion}$

11/01/2021

Confidence Interval

Introduction to confidence interval for one

mean (sigma(σ) known)

Standard deviation

statistics

When sampling from a normally distributed population with a known value of σ

$\bar{x}$ is a point estimator of μ - population mean

Sample mean

~~that close is the~~

The confidence interval shows or

relates how close the sample mean

is to the population mean

→

point estimate - single value.

Confidence interval is of the form $\bar{x} \pm \text{Margin of Error}$

Some assumptions will be made, and then

an appropriate margin of error is determined.

(1) A simple random sample from a population of interest

(2) A normally distributed population.

(3) The population standard deviation (σ) must be known.

Under these assumptions, the confidence interval formula

$(1-\alpha)100\%$ confidence interval for μ

$$\bar{x} \pm \frac{Z_{\alpha} \cdot \sigma}{\sqrt{n}}$$

where $\frac{Z_{\alpha} \cdot \sigma}{\sqrt{n}}$ is the margin of error.

The $(1-\alpha)100\%$ is the confidence level, often chosen to be 95%, 99% or 99.9%

95%
90%
99.9%
Standard Intervals


The appropriate Z value is based on the confidence level.

for $(1-\alpha) 100\%$

The area under the entire curve is 1

The remaining area is α .


split it evenly into both tails



The appropriate Z value of any given confidence level will have to be found from the standard normal distribution table.

for a 95% confidence interval,

$$(1-\alpha) 100\% = 95\%$$



$$Z\left(\frac{\alpha}{2}\right)$$

$$= Z(0.025) = 1.96 \text{ from Normal dist. table.}$$


$$\bar{X} \pm 1.96 \times \frac{s}{\sqrt{n}}$$

for a 90% confidence interval,

$$(1-\alpha) 100\% = 90\%$$

$$1 - 0.90 = 0.10$$

$$\frac{\alpha}{2} = \frac{0.10}{2} = 0.05$$



$$Z\left(\frac{\alpha}{2}\right) = Z(0.05) = 1.645$$

Example

$$n = 135$$

$$\bar{x} = 0.988$$

find 95% confidence interval for the population mean

(suppose it is known that $\sigma = 0.028$)

$$n = 135$$

$$\bar{x} = 0.988$$

$$\alpha = 1 - 0.95 = 0.05$$

$$\sigma = 0.028 \quad \frac{\alpha}{2} = \frac{0.05}{2} = 0.025$$

$$\bar{x} \pm Z\left(\frac{\alpha}{2}\right) \times \frac{\sigma}{\sqrt{n}} \quad Z(0.025) = 1.96$$

$$= 0.988 \pm Z(0.025) \times \frac{0.028}{\sqrt{135}}$$

$$= 0.988 \pm 1.96 \times \frac{0.028}{\sqrt{135}}$$

$$= 0.988 \pm 0.0047$$


$$= [0.988 + 0.0047, 0.988 - 0.0047]$$

$$= (0.983, 0.993)$$

$$(1 - \alpha) \times 100 = 95\%$$
$$1 - 0.05 = 0.95$$

Therefore we are 95% confident that the population mean lies in this interval.

Video 2 - Confidence Intervals and margin of error - Khan Academy.



P = population proportion
 $\hat{p}$ = sampling proportion

$$\sigma_{\hat{p}} = \sqrt{\frac{P(1-P)}{n}}$$

There is a 95% probability that the population proportions P is within 2

standard deviations of $\hat{p}$

Sample proportion $\hat{p} = 0.54$, $n = 100$


Sample Standard Error of sample proportion

$$\sigma_{\hat{p}} = \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

$$0.54 - (0.40), \quad 0.54 + (0.40)$$

$$= \sqrt{\frac{0.54 \cdot 0.46}{100}}$$

$$\approx 0.05$$



Population of likely voters

P = proportion that supports candidate A

Sample 1: $n = 100$

Sample proportion $\hat{p} = 0.54$

Sample distribution of the sample proportion
for $n = 100$



The mean of the sampling distribution = Population of the proportion

$\sigma_{\hat{p}}$ = standard deviation of the sample proportion.

23/06/2021 SAMPLING DISTRIBUTION

The average male drinks 2L of water when 0.7L active outdoors (with a standard deviation of 0.1L).

You are planning a full day nature trip for 50 men and will bring 110L of water. What is the probability that you will run out

σ (average / mean) = $\frac{\text{population mean } \mu}{n}$

$$\sigma = 0.7L$$

$$n = 50$$

$n > 30$ for normal distribution

$$\text{Sample mean } \bar{x} = \frac{110}{50} = 2.2L \quad (\text{Sample mean})$$

Population mean $\mu = 2L$

Sample mean = $\bar{x} = 2.2L$

$$\text{Variance } (\sigma_x^2) = \frac{\sigma^2}{n} \quad \sigma_x = \frac{\sigma}{\sqrt{n}}$$

$$\sigma_x = \frac{\sigma}{\sqrt{n}} = \frac{0.7}{\sqrt{50}} = 0.099$$

$$Z = \frac{\bar{x} - \mu}{\sigma} = \frac{202 - 200}{0.99} = 2.02$$

from Z table, $P(2.02) = 0.9783$



$$P(\text{run out of water}) = 1 - 0.9783 = 0.0217$$


$$= 2.2\%$$

(2) Suppose samples of 100 items were drawn from a population with a mean of 200 and standard deviation 30. What proportion of the samples will have mean

(a) greater than 202

(b) < 199

(c) between 198.5 and 203



$$n = 100$$

$$\mu = 200 \quad (\text{population mean})$$

$$\sigma = 30$$

(i) $\bar{x} = 202$

$$Z = \frac{\bar{x} - \mu}{\sigma}$$

$$n = 100$$

$$\sigma_{\bar{x}}$$

$$Z = \frac{202 - 200}{\frac{30}{\sqrt{100}}}$$

$$= \frac{30}{10} = 3$$

$$Z = \frac{2}{3} = 0.67$$

$$P(Z = 0.67)$$

positive tables
 $Z = 0.7486$

$$= 1 - 0.7486 = (0.2514)$$

(ii)

$$Z = \frac{\bar{x} - \mu}{\sigma_{\bar{x}}}$$

$$x = 199$$

$$Z = \frac{199 - 200}{\frac{30}{\sqrt{100}}} = -\frac{1}{3} = -0.33 \quad (\text{negative tables})$$


$$= 0.3707 \approx 37.07\%$$

(iii) Between 198.5 and 203

$$Z = \frac{198.5 - 200}{\frac{30}{\sqrt{100}}}$$


$$= -\frac{1.5}{3} = -0.50$$

$$Z(-0.5) = 0.3085$$



$$Z = \frac{203 - 250}{3} = \frac{3}{3} = 1.00$$

$$Z_{1.00} = 0.8413$$



$\bar{x}$ is a point estimator of μ . $\bar{x}$ takes $\alpha = 0.8413 - 0.3085 = 0.5328 = 53.28\%$ single value.

Confidence Interval

A sample of 135 women yielded an average of 20/48 ratio of 0.988.

What is a 95% confidence interval for the population mean?

(Suppose it is known that $\sigma = 0.028$)

$$0.95 = 1.6$$

$$0.90 = 2.58$$

30-06-2021


Confidence Interval

let $P =$ Sample proportion

let $\bar{p} =$ the population proportion of the defective nails

$P =$ sample proportion = no of defective in the sample

$$\sigma_P = \sqrt{\frac{P(1-P)}{n}}$$



$$Z(0.025) \Rightarrow (0.025 + 0.95) = 0.9750$$

The variance of bolts joined in an industry is 100 while the mean height of ~~selected~~ 16 bolts collected from the

industry is 185
find a 95% confidence interval for the
population mean and make a confidence
statement.

Solution

Both height (σ^2) - population variance = 100

$$S.D \pm 0 = \sqrt{100} = 10, \quad n = 16, \quad \bar{x} = 185.$$

$$\bar{x} \pm z\left(\frac{\alpha}{2}\right) \frac{\sigma}{\sqrt{n}}$$

$$1 - 0.95 = 0.05$$

$$\frac{\alpha}{2} = 0.025$$

$$z(0.95 + 0.025) = 0.975$$

$$\approx 1.96$$

$$185 \pm 1.96 \times \frac{10}{\sqrt{16}}$$

$$\text{upper limit} = 189.15$$

$$\text{lower limit} = 180.10.$$

Confidence statement - The result shows that

at 95% confidence level (probability), the
interval 180.10 and 189.5 contains the
mean height of the both population.

~~Obj~~ If the purposes of the survey on
affordable action discussed

Example 9.1

$P =$ no of successes in the sample
 n

$$P = \frac{531}{1013} = 0.530 - \text{Point estimate of proportion.}$$

1. Assume a confidence level first. (95%)

$$\bar{p} \pm z\left(\frac{\alpha}{2}\right) \sqrt{\frac{p(1-p)}{n}} \quad np \geq 10$$

$$= 0.530 \pm 1.96 \sqrt{\frac{0.530(1-0.530)}{n}}$$

$$= (0.499, 0.561)$$

Confidence statement - ...

Confidence interval ~~for~~ the population mean (μ) when σ is unknown.

$$\sigma_x = \frac{\sigma}{\sqrt{n}}$$

population S.D

$$z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

$$-1.96 < \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}} < 1.96$$

$$\bar{x} - 1.96\left(\frac{\sigma}{\sqrt{n}}\right) < \mu < \bar{x} + 1.96\left(\frac{\sigma}{\sqrt{n}}\right)$$

If S.D is unknown, use the sample data to determine it.

Then t distribution


$$t = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}$$

* Signif unknown

* Sample size small

Use T -table

Curve for t table
Similar to the z table



Properties

The t -value corresponding to any number of degree of freedom is

Degree of freedom = $n - 1$

If $n = 30$, $df = (30 - 1) = 29$.

Example

(Population SD unknown).

From a sample of 45 students, the mean mathematics score in a school was found to be 55%, with a SD of 15.

Find the 99% confidence interval for the mean score of mathematics in the school.

SUMMARY NOTES

$$\bar{x} = 55.9$$

$$n = 49$$

$$d = 49 - 1 = 48$$

$$\text{confidence level} = 99\%$$

$$S = 15$$

$$t(df) = t_{95\%}^{(48)} = 2.70$$

A solved problem from your book on t-distribution. ✓

$$x = a'y + b$$

$$a' = \frac{n \sum xy - (\sum x)(\sum y)}{n \sum y^2 - (\sum y)^2} \checkmark$$

$$b' = \bar{x} - a'y$$

Types of Statistics

(1) Descriptive Statistics

This summarizes data using graphs and summary values like means and interquartile range. Descriptive statistics help to realize relationships and patterns but do not draw conclusions beyond the data we have at hand.

(2) Inferential Statistics

This analysis helps us to draw conclusions beyond the data we have from the population from which it was drawn. ✓

Statistical Inference is the process of drawing conclusions about population parameters based on a sample taken from the population. The data we collect from the population is called a sample.

* The Sampling distribution of a statistic is the probability distribution of that statistic

* Statistic comes from sample

* Parameter comes from a population.

* A parameter is a measurement or quantity that ~~comes~~ ^{describes} a population

* A statistic is the measurement or quantity that helps to describe the sample

Examples of Parameter

① Population mean

② Population Variance

③ Population Standard Deviation

Examples of Statistic

① Sample mean

② Sample Variance

③ Sample Standard Deviation

* The Sampling distribution of the statistic is the subject of Statistical Inference. It is the distribution of the statistic if we are to repeatedly draw samples from the population.

A large sample is a sample with more than 30 items i.e. $n > 30$

The Standard Error - Standard deviation of a Sampling distribution.

Standard Error of Sample mean = $\frac{\sigma}{\sqrt{n}}$

Standard Error of Proportions

$$\sqrt{\frac{P(1-P)}{n}}$$

* Statistic - Any quantity computed from values in a sample is called a statistic.

* Sample Proportion - The fraction of individuals or objects in a sample that have some characteristics of interest.

(OR) - The ^{Proportion} ~~population~~ of individuals in a sample that possesses a population property.

* Unbiased Statistic

A statistic whose mean value is equal to the value of the population characteristic being estimated is said to be an unbiased statistic

A statistic that is not unbiased is said to be biased.

* Among several unbiased statistics the best statistic to use is the one with the smallest standard deviation.

* For unbiased statistic, the best statistic for obtaining a point estimate of the population mean μ is the sample mean $\bar{x}$

* Point Estimate: The point estimate of a population characteristic is a single number that is based on sample data and represents a plausible value of the characteristic.

Population - The entire collection of individuals or objects about which an information is desired

Sample - A subset of the population selected for study

A bowl contains 5 red balls, 4 white and 3 green balls. 2 balls are drawn one after the other without replacement. What is the probability of drawing a red ball first?

Sol. ~~Let a white ball is drawn first~~

$$(i) P(R,W) + P(R,R) + P(R,G)$$

$$= \text{Total} = 5 + 4 + 3 = 12 \text{ balls}$$

$$P(R,W) = \frac{5}{12} \times \frac{4}{11} = \frac{5}{33}$$

$$P(R,R) = \frac{5}{12} \times \frac{4}{11} = \frac{5}{33}$$

$$P(R,G) = \frac{5}{12} \times \frac{3}{11} = \frac{5}{44}$$

$$= \frac{5}{33} + \frac{5}{33} + \frac{5}{44} = \frac{5}{12}$$

(ii) The probability of picking a white ball the second time, given that a red ball was drawn the first time.

$$P(R,W) = \frac{5}{12} \times \frac{4}{11} = \frac{5}{33}$$

(iii) Drawing a green ball and white ball in that order

$$P(G,W) = \frac{3}{12} \times \frac{4}{11} = \frac{1}{11}$$

28-07-2021

- 2 types of t-distribution
- When to use z-tables and t-tables
- Notes, Examples, Assignments

A statement made/claim made that is to be tested - Null Hypothesis eg The average height of ^{students} ~~future~~ ^{is} 1.7m.

Alternative Hypothesis - Hypothesis against the null-hypothesis

* When the population standard deviation is not known, the t-distribution is used.

* When the standard deviation is ~~not~~ known, use z-table $z = n - 1$

$$\frac{s}{\sqrt{n}} \text{ — Standard error}$$

$$t_D = \frac{n - 1}{s}$$

$$s = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n - 1}}$$

Example

The breaking strength of cables produced by a manufacturer has a mean of $\mu = 1800$ lb and standard deviation of 160 lb. By a new

$$\begin{aligned}
 90\% &- 1.645 \\
 95\% &- 1.96 \\
 99\% &- 2.58 \\
 99.73 &- 3
 \end{aligned}$$

technique in manufacturing, it is claimed that the strength is increased. To test this claim, a sample of 50 cables is tested. It was found out that the mean is 1850 lb. Can we support this claim?

Solution

$$\mu = 1800 \text{ lb}$$


$$\sigma = 100 \text{ lb}$$

$$n = 50$$

$$\bar{x} = 1850 \text{ lb}$$

$$H_0: \mu = 1800 \text{ lb}$$

$$H_a: \mu > 1800 \text{ lb}$$



This test is a one-tailed test.


Select a significance level

Significance level = 1 - confidence level.

Use significance level of $\alpha = 0.01$ (99%) for this question.


test statistic = z

$$\frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}} = \frac{1850 - 1800}{\frac{100}{\sqrt{50}}} = 3.54$$



Based on the significance level, go to the z table and check the p-value

$$0.01 \Rightarrow 1 - 0.01 = 0.99$$



Using the 2-tailed table, $\Rightarrow$

$$[2.33]$$

For a right-hand tailed test, use 2.33 for 0.01 significant level (99%)

Since p-value = 2.33, is less than 2.58, the null hypothesis is accepted

If z-score is > the value of true p-value, the H_0 is not correct, H_a is accepted so the claim is supported.